

Lattice QCD with light Wilson quarks

Martin Lüscher

CERN — Theory Division

Wilson fermions

- ★ Conceptually well-founded
- ★ Non-perturbative improvement & renormalization
- ★ Fully worked out in qQCD

However, q QCD simulations tend to be exceedingly “expensive”

SESAM & T χ L '98, UKQCD '99, CP-PACS '00, JLQCD '03, . . .

With Wilson quarks we need, say,

$$a = 0.1 \rightarrow 0.04 \text{ fm}$$

$$L = 2 \rightarrow 5 \text{ fm}$$

$$m_\pi = 500 \rightarrow 200 \text{ MeV}$$

⇒ large lattices & solver iteration numbers

$$48 \cdot 24^3, 64 \cdot 32^3, 96 \cdot 48^3, \dots$$

$$10^3 - 10^4 \text{ applications of } \mathcal{D} \text{ per source field}$$

Algorithms should be designed to work well in this regime!

Domain-decomposition methods

- ◇ Schwarz-preconditioned solver
- ◇ HMC on a block-decomposed lattice

M.L. '03 [JHEP 0305 (2003) 052; CPC 156 (2004) 209; CPC 165 (2005) 199]

Del Debbio, Giusti, M.L., Petronzio & Tantalò '05 [in preparation]

Hasenbusch acceleration

- ◇ HMC & PHMC with polynomial preconditioner


Hasenbusch '01 [PLB 519 (2001) 177]; Hasenbusch & Jansen '03 [NPB 659 (2003) 299];

Urbach, Jansen, Shindler & Wenger '05 [hep-lat/0506011] → parallel talk by Urbach

Domain decomposition

In general the benefits are

- ★ *Parallel efficiency*
- ★ *Scale separation*
- ★ *Softer scaling behaviour*



The classical DD method is the *Schwarz Alternating Procedure*



Hermann Amandus Schwarz 1870:

Dirichlet problem in complicated domains

Widely used as a preconditioner in engineering (fluid dynamics etc.)

... and now also in lattice QCD!

see Y. Saad: *Methods for Sparse Linear Systems*, SIAM 2003, for example


The properties of QCD we shall exploit are

(a) *Asymptotic freedom*

Dirichlet b.c. on the blocks imply


$$q \geq \pi/l > 1 \text{ GeV}$$

⇒ easy to simulate at all quark masses



(b) *Ellipticity of the Dirac operator*

$$\frac{\delta^2 S_{\text{eff}}}{\delta A_{\mu}^a(x) \delta A_{\nu}^b(y)} = 2 \text{tr} \{ T^a \gamma_{\mu} S(x, y) T^b \gamma_{\nu} S(y, x) \}$$
$$\sim |x - y|^{-6}$$



⇒ blocks are approximately decoupled

Block decomposition of the Dirac operator

black blocks: Ω


white blocks: Ω^*

exterior boundaries: $\partial\Omega, \partial\Omega^*$

$$D = \frac{1}{2} \{ \gamma_\mu (\nabla_\mu^* + \nabla_\mu) - \nabla_\mu^* \nabla_\mu \} + m_0$$

$$= D_\Omega + D_{\Omega^*} + D_{\partial\Omega} + D_{\partial\Omega^*}$$

$$D_\Omega = \sum_{\text{black } \Lambda} D_\Lambda, \quad D_{\Omega^*} = \sum_{\text{white } \Lambda} D_\Lambda$$



The quark determinant factorizes into

$$\det D = \prod_{\text{blocks } \Lambda} \det \hat{D}_\Lambda \times \det R$$

\uparrow
 eo preconditioned, Dirichlet b.c.

where the block interaction is given by


$$R : \mathcal{H}_{\partial\Omega^*} \rightarrow \mathcal{H}_{\partial\Omega^*}$$

$$R = 1 - P_{\partial\Omega^*} D_\Omega^{-1} D_{\partial\Omega} D_{\Omega^*}^{-1} D_{\partial\Omega^*}$$

Note that

$$R^{-1} = 1 - P_{\partial\Omega^*} D^{-1} D_{\partial\Omega^*}$$

$\mathcal{H}_{\partial\Omega^*}$: quark fields on $\partial\Omega^*$




For $N_f = 2$ flavours

$$S_{\text{pf}} = \sum_{\text{blocks } \Lambda} \|\hat{D}_\Lambda^{-1} \phi_\Lambda\|^2 + \|R^{-1} \chi\|^2$$

where χ is defined on $\partial\Omega^*$ and ϕ_Λ on the even sites in Λ

We now

- evolve only the *active* link variables in the blocks and
- translate the gauge field after each trajectory by a random vector



Fermion forces

$$\frac{d}{dt} U(x, \mu) = \Pi(x, \mu) U(x, \mu)$$


$$\frac{d}{dt} \Pi(x, \mu) = -F_G(x, \mu) - F_\Lambda(x, \mu) - F_R(x, \mu)$$

Example

$32 \cdot 16^3$ lattice, 8^4 blocks

$\beta = 5.3$, $c_{sw} = 1.9095$

$a \sim 0.1$ fm, $m \sim 14 - 70$ MeV




This behaviour is not unexpected since

(a) F_Λ is protected by the b.c. (*asymptotic freedom*)

(b) F_R is suppressed by propagators (*ellipticity*)

$$F_R(x, \mu) =$$

$$2\text{Re} \left(R^{-1} \chi, D^{-1} \delta_{x, \mu}^U D D^{-1} D_{\partial\Omega^*} \chi \right)$$




Leap-frog integration

Sexton & Weingarten '92; Peardon & Sexton '02

Choose integration step-sizes $\varepsilon_0, \varepsilon_1, \varepsilon_2$ such that

$$\varepsilon_0 \|F_G\| \simeq \varepsilon_1 \|F_\Lambda\| \simeq \varepsilon_2 \|F_R\|$$

For example



$$\varepsilon_1 = 4\varepsilon_0, \quad \varepsilon_2 = 5\varepsilon_1$$

$\Rightarrow F_R$ is the most “expensive” but least-often computed force

Simulations

Del Debbio, Giusti, M.L., Petronzio, Tantalò [CERN – Tor Vergata]

$N_f = 2$, Wilson action

Trajectory length $\tau = 0.5$

$\Rightarrow \langle \text{link path length} \rangle = 0.53$

Schwarz-preconditioned solver

Residues $10^{-7} \dots 10^{-11}$

\Rightarrow reversibility $|(U' - U)_{ij}| < 10^{-9}$

lattice	β	κ	$\sim m/m_s$	block size	τ/ε_2	N_{traj}	P_{acc}
$32 \cdot 24^3$	5.6	0.15750	0.93	$8 \cdot 6^2 \cdot 12$	5	6400	0.80
		0.15800	0.48		6	9500	0.80
		0.15825	0.30		10	9400	0.86
		0.15835	0.17		16	5000	0.91

Simulations performed on 8 nodes (16 Xeon processors) at the ITP Bern

Large-lattice runs

lattice	β	κ	$\sim m/m_s$	block size	τ/ε_2	N_{traj}	P_{acc}
$64 \cdot 32^3$	5.8	0.15410	0.75	$16 \cdot 8^3$	8	5000	0.86
		0.15440	0.38		10	5050	0.89

Simulations performed on $32 + 32$ nodes at the Fermi Institute

The lattice spacing is

$$\left. \begin{array}{l} a \sim 0.080 \text{ fm} \quad \text{at} \quad \beta = 5.6 \\ a \sim 0.064 \text{ fm} \quad \text{at} \quad \beta = 5.8 \end{array} \right\} L \simeq 2 \text{ fm}$$

(Setting $r_0 = 0.5 \text{ fm}$ at the point where $r_0 m_\pi = 1.26$)

Chiral behaviour of m_π and F_π

SU(2) ChPT predicts

$$m_\pi^2 = M^2 R_\pi, \quad M^2 = 2Bm$$

$$R_\pi = 1 + \frac{M^2}{32\pi^2 F^2} \ln(M^2/\Lambda_\pi^2) + \dots$$



where, in real-world QCD,

$$\ln(\Lambda_\pi^2/M^2) \Big|_{M=140 \text{ MeV}} \simeq 2.9 \pm 2.4$$

Gasser & Leutwyler '84

$\Rightarrow R_\pi \simeq \text{constant} = 0.956(8)$ in the range $M = 200 - 500 \text{ MeV}$

32 · 24³ lattice




For F_π we expect

$$F_\pi = F - \frac{M^2}{16\pi^2 F} \ln(M^2/\Lambda_F^2) + \dots$$

$$\ln(\Lambda_F^2/M^2)|_{M=140 \text{ MeV}} \simeq 4.6 \pm 0.9$$

This fits the last three points

$$\Rightarrow F_\pi|_{M=140 \text{ MeV}} = 80(7) \text{ MeV}$$






*Within errors and up to $m_\pi \sim 500$ MeV,
the data are compatible with 1-loop ChPT*


Large-lattice experiences

- ★ Similar step numbers as on the small lattices
- ★ Runs are very stable
- ★ Long effective-mass plateaus

$64 \cdot 32^3$ lattice, $m_\pi \simeq 429$ MeV



$m_\pi \simeq 606$ MeV



Performance figures & timings

A comparison of algorithms is non-trivial since

- *their efficiency depends on the lattice parameters, the program and the computer*
- *there are many tunable parameters*
- *it is generally difficult to determine the relevant auto-correlation times reliably*

A purely algorithmic cost figure is

$$\nu = 10^{-3} (2N_2 + 3) \tau_{\text{int}}[P], \quad N_2 = \tau/\varepsilon_2$$

$\nu \simeq 5 - 29$ in previous HMC simulations [SESAM, UKQCD, CP-PACS, ...]


lattice	β	c_{sw}	am	$\tau_{\text{int}}[P]$	ν
32 · 16 ³	5.3	1.9095	0.035	56(26)	0.84(39)
			0.021	14(4)	0.27(8)
			0.011	21(6)	0.40(11)
			0.007	17(5)	0.39(12)
32 · 24 ³	5.6	0.0	0.027	53(22)	0.69(29)
			0.014	33(11)	0.50(17)
			0.009	27(10)	0.62(23)
			0.006	21(5)	0.74(18)
64 · 32 ³	5.8	0.0	0.019	16(3)	0.30(6)
			0.011	13(2)	0.30(5)

} preliminary

Timings

Accepted trajectories per day using 8 and 32 nodes

Note that $\frac{64 \cdot 32^3}{32 \cdot 24^3} = 4.74$



Conclusions

Numerical simulations of LQCD with light Wilson quarks are much less “expensive” than previously estimated!

⇒ it is now possible to reach the chiral regime on large lattices

Example

$96 \cdot 48^3$ lattice, $12^2 \cdot 8^2$ blocks

$a = 0.06$ fm, $am = 0.0035$, $m_\pi = 270$ MeV

To simulate this lattice, a (current) PC cluster with 288 nodes should be sufficient

The Schwarz-preconditioned HMC algorithm

- ★ works out at $a \leq 0.1$ fm and on large lattices, now also with $O(a)$ improvement
- ★ scales favourably & is highly parallel
- ★ easily extends to gauge actions with 6-link terms (Iwasaki, Symanzik)

A wide range of physics questions may now be addressed, in a conceptually solid framework
