

Course 311, Part IV: Galois Theory Problems Hilary Term 2006

1. Use Eisenstein's criterion to verify that the following polynomials are irreducible over \mathbb{Q} :—

$$(i) \ t^2 - 2; \quad (ii) \ t^3 + 9t + 3; \quad (iii) \ t^5 + 26t + 52.$$

2. Let p be a prime number. Use the fact that the binomial coefficient $\binom{p}{k}$ is divisible by p for all integers k satisfying $0 < k < p$ to show that if $tf(t) = (t+1)^p - 1$ then the polynomial f is irreducible over \mathbb{Q} .

The *cyclotomic polynomial* $\Phi_p(t)$ is defined by $\Phi_p(t) = 1 + t + t^2 + \cdots + t^{p-1}$ for each prime number p . Show that $t\Phi_p(t+1) = (t+1)^p - 1$, and hence show that the cyclotomic polynomial Φ_p is irreducible over \mathbb{Q} for all prime numbers p .

3. The Fundamental Theorem of Algebra ensures that every non-constant polynomial with complex coefficients factors as a product of polynomials of degree one. Use this result to show that a non-constant polynomial with real coefficients is irreducible over the field \mathbb{R} of real numbers if and only if it is either a polynomial of the form $at + b$ with $a \neq 0$ or a quadratic polynomial of the form $at^2 + bt + c$ with $a \neq 0$ and $b^2 < 4ac$.
4. A complex number z is said to be *algebraic* if there $f(z) = 0$ for some non-zero polynomial f with rational coefficients. Show that $z \in \mathbb{C}$ is algebraic if and only if $\mathbb{Q}(z) : \mathbb{Q}$ is a finite extension. Then use the Tower Law to prove that the set of all algebraic numbers is a subfield of \mathbb{C} .
5. Let L be a splitting field for a polynomial of degree n with coefficients in K . Prove that $[L : K] \leq n!$.
6. (a) Show that $\mathbb{Q}(\sqrt{2}, \sqrt{3}) = \mathbb{Q}(\sqrt{2} + \sqrt{3})$ and $[\mathbb{Q}(\sqrt{2}, \sqrt{3}), \mathbb{Q}] = 4$. What is the degree of the minimum polynomial of $\sqrt{2} + \sqrt{3}$ over \mathbb{Q} ?
(b) Show that $\sqrt{2} + \sqrt{3}$ is a root of the polynomial $t^4 - 10t^2 + 1$, and thus show that this polynomial is an irreducible polynomial whose splitting field over \mathbb{Q} is $\mathbb{Q}(\sqrt{2}, \sqrt{3})$.
(c) Find all \mathbb{Q} -automorphisms of $\mathbb{Q}(\sqrt{2}, \sqrt{3})$, and show that they constitute a group of order 4 isomorphic to a direct product of two cyclic groups of order 2.

7. Let K be a field of characteristic p , where p is prime.
- (a) Show that $f \in K[t]$ satisfies $Df = 0$ if and only if $f(t) = g(t^p)$ for some $g \in K[t]$.
- (b) Let $h(t) = a_0 + a_1t + a_2t^2 + \cdots + a_nt^n$, where $a_0, a_1, \dots, a_n \in K$. Show that $(h(t))^p = g(t^p)$, where $g(t) = a_0^p + a_1^p t + a_2^p t^2 + \cdots + a_n^p t^n$.
- (c) Now suppose that Frobenius monomorphism of K is an automorphism of K . Show that $f \in K[t]$ satisfies $Df = 0$ if and only if $f(t) = (h(t))^p$ for some $h \in K[t]$. Hence show that $Df \neq 0$ for any irreducible polynomial f in $K[t]$.
- (d) Use these results to show that every algebraic extension $L: K$ of a finite field K is separable.
8. For each positive integer n , let ω_n be the primitive n th root of unity in \mathbb{C} given by $\omega_n = \exp(2\pi i/n)$, where $i = \sqrt{-1}$.
- (a) Show that the field extensions $\mathbb{Q}(\omega_n):\mathbb{Q}$ and $\mathbb{Q}(\omega_n, i):\mathbb{Q}$ are normal field extensions for all positive integers n .
- (b) Show that the minimum polynomial of ω_p over \mathbb{Q} is the *cyclotomic polynomial* $\Phi_p(t)$ given by $\Phi_p(t) = 1 + t + t^2 + \cdots + t^{p-1}$. Hence show that $[\mathbb{Q}(\omega_p):\mathbb{Q}] = p - 1$ if p is prime.
- (c) Let p be prime and let $\alpha_k = \omega_{p^2}\omega_p^k = \exp(2\pi i(1 + kp)/p^2)$ for all integers k . Note that $\alpha_0 = \omega_{p^2}$ and $\alpha_k = \alpha_l$ if and only if $k \equiv l \pmod{p}$. Show that if θ is an automorphism of $\mathbb{Q}(\omega_{p^2})$ which fixes $\mathbb{Q}(\omega_p)$ then there exists some integer m such that $\theta(\alpha_k) = \alpha_{k+m}$ for all integers k . Hence show that $\alpha_0, \alpha_1, \dots, \alpha_{p-1}$ all belong to the orbit of ω_{p^2} under the action of the Galois group $\Gamma(\mathbb{Q}(\omega_{p^2}):\mathbb{Q}(\omega_p))$. Use this result to show that $[\mathbb{Q}(\omega_{p^2}):\mathbb{Q}(\omega_p)] = p$ and $[\mathbb{Q}(\omega_{p^2}):\mathbb{Q}] = p(p - 1)$.
9. Show that the field $\mathbb{Q}(\xi, \omega)$ is a splitting field for the polynomial $t^5 - 2$ over \mathbb{Q} , where $\omega = \omega_5 = \exp(2\pi i/5)$ and $\xi = \sqrt[5]{2}$. Show that $[\mathbb{Q}(\xi, \omega):\mathbb{Q}] = 20$ and the Galois $\Gamma(\mathbb{Q}(\xi, \omega):\mathbb{Q})$ consists of the automorphisms $\theta_{r,s}$ for $r = 1, 2, 3, 4$ and $s = 0, 1, 2, 3, 4$, where $\theta_{r,s}(\omega) = \omega^r$ and $\theta_{r,s}(\xi) = \omega^s \xi$.

10. Let f be a monic polynomial of degree n with coefficients in a field K . Then

$$f(t) = (t - \alpha_1)(t - \alpha_2) \cdots (t - \alpha_n),$$

where $\alpha_1, \alpha_2, \dots, \alpha_n$ are the roots of f in some splitting field L for f over K . The *discriminant* of the polynomial f is the quantity δ^2 , where δ is the product $\prod_{1 \leq i < j \leq n} (\alpha_j - \alpha_i)$ of the quantities $\alpha_j - \alpha_i$ taken over all pairs of integers i and j satisfying $1 \leq i < j \leq n$.

Show that the quantity δ changes sign whenever α_i is interchanged with α_{i+1} for some i between 1 and $n - 1$. Hence show that $\theta(\delta) = \delta$ for all automorphisms θ in the Galois group $\Gamma(L:K)$ that induce even permutations of the roots of f , and $\theta(\delta) = -\delta$ for all automorphisms θ in $\Gamma(L:K)$ that induce odd permutations of the roots. Then apply the Galois correspondence to show that the discriminant δ^2 of the polynomial f belongs to the field K containing the coefficients of f , and the field $K(\delta)$ is the fixed field of the subgroup of $\Gamma(L:K)$ consisting of those automorphisms in $\Gamma(L:K)$ that induce even permutations of the roots of f . Hence show that $\delta \in K$ if and only if all automorphisms in the Galois group $\Gamma(L:K)$ induce even permutations of the roots of f .

11. (a) Show that the discriminant of the quadratic polynomial $t^2 + bt + c$ is $b^2 - 4c$.

(b) Show that the discriminant of the cubic polynomial $t^3 - pt - q$ is $4p^2 - 27q^2$.

12. Let $f(t) = t^3 - pt - q$ be a cubic polynomial with complex coefficients p and q , and let the complex numbers α , β and γ be the roots of f .

(a) Give formulae for the coefficients p and q of f in terms of the roots α , β and γ of f , and verify that $\alpha + \beta + \gamma = 0$ and

$$\alpha^3 + \beta^3 + \gamma^3 = 3\alpha\beta\gamma = 3q$$

(b) Let $\lambda = \alpha + \omega\beta + \omega^2\gamma$ and $\mu = \alpha + \omega^2\beta + \omega\gamma$, where ω is the complex cube root of unity given by $\omega = \frac{1}{2}(-1 + \sqrt{3}i)$. Verify that $1 + \omega + \omega^2 = 0$, and use this result to show that

$$\alpha = \frac{1}{3}(\lambda + \mu), \quad \beta = \frac{1}{3}(\omega^2\lambda + \omega\mu), \quad \gamma = \frac{1}{3}(\omega\lambda + \omega^2\mu).$$

(c) Let K be the subfield $\mathbb{Q}(p, q)$ of \mathbb{C} generated by the coefficients of the polynomial f , and let M be a splitting field for the polynomial f over $K(\omega)$. Show that the extension $M:K$ is normal, and is thus a Galois extension. Show that any automorphism in the Galois group $\Gamma(M:K)$ permutes the roots α, β and γ of f and either fixes ω or else sends ω to ω^2 .

(d) Let $\theta \in \Gamma(M:K)$ be a K -automorphism of M . Suppose that

$$\theta(\alpha) = \beta, \quad \theta(\beta) = \gamma, \quad \theta(\gamma) = \alpha.$$

Show that if $\theta(\omega) = \omega$ then $\theta(\lambda) = \omega^2\lambda$ and $\theta(\mu) = \omega\mu$. Show also that if $\theta(\omega) = \omega^2$ then $\theta(\lambda) = \omega\mu$ and $\theta(\mu) = \omega^2\lambda$. Hence show that $\lambda\mu$ and $\lambda^3 + \mu^3$ are fixed by any automorphism in $\Gamma(M:K)$ that cyclically permutes α, β, γ . Show also that the quantities $\lambda\mu$ and $\lambda^3 + \mu^3$ are also fixed by any automorphism in $\Gamma(M:K)$ that interchanges two of the roots of f whilst leaving the third root fixed. Hence prove that $\lambda\mu$ and $\lambda^3 + \mu^3$ belong to the field K generated by the coefficients of f and can therefore be expressed as rational functions of p and q .

(e) Show by direct calculation that $\lambda\mu = 3p$ and $\lambda^3 + \mu^3 = 27q$. Hence show that λ^3 and μ^3 are roots of the quadratic polynomial $t^2 - 27qt + 27p^3$. Use this result to verify that the roots of the cubic polynomial $t^3 - pt - q$ are of the form

$$\sqrt[3]{\frac{q}{2} + \sqrt{\frac{q^2}{4} - \frac{p^3}{27}}} + \sqrt[3]{\frac{q}{2} - \sqrt{\frac{q^2}{4} - \frac{p^3}{27}}}$$

where the two cube roots must be chosen so as to ensure that their product is equal to $\frac{1}{3}p$.