

Module MA2321: Analysis in Several Real
Variables
Michaelmas Term 2018
Part III (Section 7)

D. R. Wilkins

Copyright © David R. Wilkins 2015–2018

Contents

7	Differentiation of Functions of One Real Variable	77
7.1	Interior Points and Open Sets in the Real Line	77
7.2	Differentiable Functions of a Single Real Variable	77
7.3	Rolle’s Theorem	78
7.4	The Mean Value Theorem	79
7.5	Concavity and the Second Derivative	80
7.6	Taylor’s Theorem	81

7 Differentiation of Functions of One Real Variable

7.1 Interior Points and Open Sets in the Real Line

Definition Let D be a subset of the set \mathbb{R} of real numbers, and let s be a real number belonging to D . We say that s is an *interior point* of D if there exists some strictly positive number δ such that $x \in D$ for all real numbers x satisfying $s - \delta < x < s + \delta$. The *interior* of D is then the subset of D consisting of all real numbers belonging to D that are interior points of D .

It follows from the definition of open sets in Euclidean spaces that a subset D of the set \mathbb{R} of real numbers is an open set in \mathbb{R} if and only if every point of D is an interior point of D .

Let s be a real number. We say that a function $f: D \rightarrow \mathbb{R}$ is defined *around* s if the real number s is an interior point of the domain D of the function f . It follows that the function f is defined around s if and only if there exists some strictly positive real number δ such that $f(x)$ is defined for all real numbers x satisfying $s - \delta < x < s + \delta$.

7.2 Differentiable Functions of a Single Real Variable

We recall basic results of the theory of differentiable functions.

Definition Let s be some real number, and let f be a real-valued function defined around s . The function f is said to be *differentiable* at s , with *derivative* $f'(s)$, if and only if the limit

$$f'(s) = \lim_{h \rightarrow 0} \frac{f(s+h) - f(s)}{h}$$

is well-defined. We denote by f' , or by $\frac{df}{dx}$ the function whose value at s is the derivative $f'(s)$ of f at s .

Let s be some real number, and let f and g be real-valued functions defined around s that are differentiable at s . The basic rules of differential calculus then ensure that the functions $f+g$, $f-g$ and $f \cdot g$ are differentiable at s (where

$$(f+g)(x) = f(x) + g(x), \quad (f-g)(x) = f(x) - g(x)$$

and

$$(f \cdot g)(x) = f(x)g(x)$$

for all real numbers x at which both $f(x)$ and $g(x)$ are defined), and

$$(f + g)'(s) = f'(s) + g'(s), \quad (f - g)'(s) = f'(s) - g'(s).$$

$$(f \cdot g)'(s) = f'(s)g(s) + f(s)g'(s) \quad (\text{Product Rule}).$$

If moreover $g(s) \neq 0$ then the function f/g is differentiable at s (where $(f/g)(x) = f(x)/g(x)$ where both $f(x)$ and $g(x)$ are defined), and

$$(f/g)'(s) = \frac{f'(s)g(s) - f(s)g'(s)}{g(s)^2} \quad (\text{Quotient Rule}).$$

Moreover if h is a real-valued function defined around $f(s)$ which is differentiable at $f(s)$ then the composition function $h \circ f$ is differentiable at $f(s)$ and

$$(h \circ f)'(s) = h'(f(s))f'(s) \quad (\text{Chain Rule}).$$

Derivatives of some standard functions are as follows:—

$$\frac{d}{dx}(x^m) = mx^{m-1}, \quad \frac{d}{dx}(\sin x) = \cos x, \quad \frac{d}{dx}(\cos x) = -\sin x,$$

$$\frac{d}{dx}(\exp x) = \exp x, \quad \frac{d}{dx}(\log x) = \frac{1}{x} \quad (x > 0).$$

7.3 Rolle's Theorem

Theorem 7.1 (Rolle's Theorem) *Let $f: [a, b] \rightarrow \mathbb{R}$ be a real-valued function defined on some interval $[a, b]$. Suppose that f is continuous on $[a, b]$ and is differentiable on (a, b) . Suppose also that $f(a) = f(b)$. Then there exists some real number s satisfying $a < s < b$ which has the property that $f'(s) = 0$.*

Proof First we show that if the function f attains its minimum value at u , and if $a < u < b$, then $f'(u) = 0$. Now the difference quotient

$$\frac{f(u+h) - f(u)}{h}$$

is non-negative for all sufficiently small positive values of h ; therefore $f'(u) \geq 0$. On the other hand, this difference quotient is non-positive for all sufficiently small negative values of h ; therefore $f'(u) \leq 0$. We deduce therefore that $f'(u) = 0$.

Similarly if the function f attains its maximum value at v , and if $a < v < b$, then $f'(v) = 0$. (Indeed the result for local maxima can be deduced from the corresponding result for local minima simply by replacing the function f by $-f$.)

Now the function f is continuous on the closed bounded interval $[a, b]$. It therefore follows from the Extreme Value Theorem that there must exist real numbers u and v in the interval $[a, b]$ with the property that $f(u) \leq f(x) \leq f(v)$ for all real numbers x satisfying $a \leq x \leq b$ (see Theorem 4.21). If $a < u < b$ then $f'(u) = 0$ and we can take $s = u$. Similarly if $a < v < b$ then $f'(v) = 0$ and we can take $s = v$. The only remaining case to consider is when both u and v are endpoints of the interval $[a, b]$. In that case the function f is constant on $[a, b]$, since $f(a) = f(b)$, and we can choose s to be any real number satisfying $a < s < b$. ■

7.4 The Mean Value Theorem

Rolle's Theorem can be generalized to yield the following important theorem.

Theorem 7.2 (The Mean Value Theorem) *Let $f: [a, b] \rightarrow \mathbb{R}$ be a real-valued function defined on some interval $[a, b]$. Suppose that f is continuous on $[a, b]$ and is differentiable on (a, b) . Then there exists some real number s satisfying $a < s < b$ which has the property that*

$$f(b) - f(a) = f'(s)(b - a).$$

Proof Let $g: [a, b] \rightarrow \mathbb{R}$ be the real-valued function on the closed interval $[a, b]$ defined by

$$g(x) = f(x) - \frac{b-x}{b-a}f(a) - \frac{x-a}{b-a}f(b).$$

Then the function g is continuous on $[a, b]$ and differentiable on (a, b) . Moreover $g(a) = 0$ and $g(b) = 0$. It follows from Rolle's Theorem (Theorem 7.1) that $g'(s) = 0$ for some real number s satisfying $a < s < b$. But

$$g'(s) = f'(s) - \frac{f(b) - f(a)}{b - a}.$$

Therefore $f(b) - f(a) = f'(s)(b - a)$, as required. ■

A number of basic principles of single variable calculus follow as immediate consequences of the Mean Value Theorem (Theorem 7.2). A number of such consequences are presented in the following corollaries.

Corollary 7.3 Let $f: [a, b] \rightarrow \mathbb{R}$ be a real-valued function defined on some interval $[a, b]$. Suppose that f is continuous on $[a, b]$ and is differentiable on (a, b) and that $f'(x) > 0$ for all real numbers x satisfying $a < x < b$. Then $f(b) > f(a)$.

Corollary 7.4 Let $f: [a, b] \rightarrow \mathbb{R}$ be a real-valued function defined on some interval $[a, b]$. Suppose that f is continuous on $[a, b]$ and is differentiable on (a, b) and that $f'(x) = 0$ for all real numbers x satisfying $a < x < b$. Then $f(x) = f(a)$ for all $x \in [a, b]$.

Corollary 7.5 Let $f: [a, b] \rightarrow \mathbb{R}$ be a real-valued function defined on some interval $[a, b]$, and let M be a real number. Suppose that f is continuous on $[a, b]$ and is differentiable on (a, b) and that $f'(x) \leq M$ for all real numbers x satisfying $a < x < b$. Then $f(x) \leq f(a) + M(x - a)$ for all $x \in [a, b]$.

Corollary 7.6 Let $f: [a, b] \rightarrow \mathbb{R}$ be a real-valued function defined on some interval $[a, b]$, and let M be a real number. Suppose that f is continuous on $[a, b]$ and is differentiable on (a, b) and that $|f'(x)| \leq M$ for all real numbers x satisfying $a < x < b$. Then $|f(b) - f(a)| \leq M(b - a)$.

7.5 Concavity and the Second Derivative

Proposition 7.7 Let s and h be real numbers, and let f be a twice differentiable real-valued function defined on some open interval containing s and $s + h$. Then there exists a real number θ satisfying $0 < \theta < 1$ for which

$$f(s + h) = f(s) + hf'(s) + \frac{1}{2}h^2 f''(s + \theta h).$$

Proof Let I be an open interval, containing the real numbers 0 and 1, chosen to ensure that $f(s + th)$ is defined for all $t \in I$, and let $q: I \rightarrow \mathbb{R}$ be defined so that

$$q(t) = f(s + th) - f(s) - thf'(s) - t^2(f(s + h) - f(s) - hf'(s)).$$

for all $t \in I$. Differentiating, we find that

$$q'(t) = hf'(s + th) - hf'(s) - 2t(f(s + h) - f(s) - hf'(s))$$

and

$$q''(t) = h^2 f''(s + th) - 2(f(s + h) - f(s) - hf'(s)).$$

Now $q(0) = q(1) = 0$. It follows from Rolle's Theorem, applied to the function q on the interval $[0, 1]$, that there exists some real number φ satisfying $0 < \varphi < 1$ for which $q'(\varphi) = 0$.

Then $q'(0) = q'(\varphi) = 0$, and therefore Rolle's Theorem can be applied to the function q' on the interval $[0, \varphi]$ to prove the existence of some real number θ satisfying $0 < \theta < \varphi$ for which $q''(\theta) = 0$. Then

$$0 = q''(\theta) = h^2 f''(s + \theta h) - 2(f(s + h) - f(s) - hf'(s)).$$

Rearranging, we find that

$$f(s + h) = f(s) + hf'(s) + \frac{1}{2}h^2 f''(s + \theta h),$$

as required. ■

Corollary 7.8 *Let $f: (s - \delta_0, s + \delta_0)$ be a twice-differentiable function throughout some open interval $(s - \delta_0, s + \delta_0)$ centred on a real number s . Suppose that $f''(s + h) > 0$ for all real numbers h satisfying $|h| < \delta_0$. Then*

$$f(s + h) \geq f(s) + hf'(s)$$

for all real numbers h satisfying $|h| < \delta_0$.

It follows from Corollary 7.8 that if a twice-differentiable function has positive second derivative throughout some open interval, then it is concave upwards throughout that interval. In particular the function has a local minimum at any point of that open interval where the first derivative is zero and the second derivative is positive.

Corollary 7.9 *Let $f: D \rightarrow \mathbb{R}$ be a twice-differentiable function defined over a subset D of \mathbb{R} , and let s be a point in the interior of D . Suppose that $f'(s) = 0$ and that $f''(x) > 0$ for all real numbers x belonging to some sufficiently small neighbourhood of s . Then s is a local minimum for the function f .*

7.6 Taylor's Theorem

The result obtained in Proposition 7.7 is a special case of a more general result. That more general result is a version of Taylor's Theorem with remainder. The proof of this theorem will make use of the following lemma.

Lemma 7.10 *Let s and h be real numbers, let f be a k times differentiable real-valued function defined on some open interval containing s and $s + h$, let c_0, c_1, \dots, c_{k-1} be real numbers, and let*

$$p(t) = f(s + th) - \sum_{n=0}^{k-1} c_n t^n.$$

for all real numbers t belonging to some open interval D for which $0 \in D$ and $1 \in D$. Then $p^{(n)}(0) = 0$ for all integers n satisfying $0 \leq n < k$ if and only if

$$c_n = \frac{h^n f^{(n)}(s)}{n!}$$

for all integers n satisfying $0 \leq n < k$.

Proof On setting $t = 0$, we find that $p(0) = f(s) - c_0$, and thus $p(0) = 0$ if and only if $c_0 = f(s)$.

Let the integer n satisfy $0 < n < k$. On differentiating $p(t)$ n times with respect to t , we find that

$$p^{(n)}(t) = h^n f^{(n)}(s + th) - \sum_{j=n}^{k-1} \frac{j!}{(j-n)!} c_j t^{j-n}.$$

Then, on setting $t = 0$, we find that only the term with $j = n$ contributes to the value of the sum on the right hand side of the above identity, and therefore

$$p^{(n)}(0) = h^n f^{(n)}(s) - n!c_n.$$

The result follows. ■

Theorem 7.11 (Taylor's Theorem) *Let s and h be real numbers, and let f be a k times differentiable real-valued function defined on some open interval containing s and $s + h$. Then*

$$f(s + h) = f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + \frac{h^k}{k!} f^{(k)}(s + \theta h)$$

for some real number θ satisfying $0 < \theta < 1$.

Proof Let D be an open interval, containing the real numbers 0 and 1, chosen to ensure that $f(s + th)$ is defined for all $t \in D$, and let $p: D \rightarrow \mathbb{R}$ be defined so that

$$p(t) = f(s + th) - f(s) - \sum_{n=1}^{k-1} \frac{t^n h^n}{n!} f^{(n)}(s)$$

for all $t \in D$. A straightforward calculation shows that $p^{(n)}(0) = 0$ for $n = 0, 1, \dots, k-1$ (see Lemma 7.10). Thus if $q(t) = p(t) - p(1)t^k$ for all $s \in [0, 1]$ then $q^{(n)}(0) = 0$ for $n = 0, 1, \dots, k-1$, and $q(1) = 0$. We can therefore apply Rolle's Theorem (Theorem 7.1) to the function q on the interval $[0, 1]$

to deduce the existence of some real number t_1 satisfying $0 < t_1 < 1$ for which $q'(t_1) = 0$. We can then apply Rolle's Theorem to the function q' on the interval $[0, t_1]$ to deduce the existence of some real number t_2 satisfying $0 < t_2 < t_1$ for which $q''(t_2) = 0$. By continuing in this fashion, applying Rolle's Theorem in turn to the functions $q'', q''', \dots, q^{(k-1)}$, we deduce the existence of real numbers t_1, t_2, \dots, t_k satisfying $0 < t_k < t_{k-1} < \dots < t_1 < 1$ with the property that $q^{(n)}(t_n) = 0$ for $n = 1, 2, \dots, k$. Let $\theta = t_k$. Then $0 < \theta < 1$ and

$$0 = \frac{1}{k!}q^{(k)}(\theta) = \frac{1}{k!}p^{(k)}(\theta) - p(1) = \frac{h^k}{k!}f^{(k)}(s + \theta h) - p(1),$$

hence

$$\begin{aligned} f(s + h) &= f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + p(1) \\ &= f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + \frac{h^k}{k!} f^{(k)}(s + \theta h), \end{aligned}$$

as required. ■

Corollary 7.12 *Let $f: D \rightarrow \mathbb{R}$ be a k -times continuously differentiable function defined over an open subset D of \mathbb{R} and let $s \in \mathbb{R}$. Then given any strictly positive real number ε , there exists some strictly positive real number δ such that*

$$\left| f(s + h) - f(s) - \sum_{n=1}^k \frac{h^n}{n!} f^{(n)}(s) \right| < \varepsilon |h|^k$$

whenever $|h| < \delta$.

Proof The function f is k -times continuously differentiable, and therefore its k th derivative $f^{(k)}$ is continuous. Let some strictly positive real number ε be given. Then there exists some strictly positive real number δ that is small enough to ensure that $s + h \in D$ and $|f^{(k)}(s + h) - f^{(k)}(s)| < k!\varepsilon$ whenever $|h| < \delta$. If h is a real number satisfying $|h| < \delta$, and if θ is a real number satisfying $0 < \theta < 1$, then $s + \theta h \in D$ and $|f^{(k)}(s + \theta h) - f^{(k)}(s)| < k!\varepsilon$. Now it follows from Taylor's Theorem (Theorem 7.11) that, given any real number h satisfying $|h| < \delta$ there exists some real number θ satisfying $0 < \theta < 1$ for which

$$f(s + h) = f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + \frac{h^k}{k!} f^{(k)}(s + \theta h).$$

Then

$$\begin{aligned} \left| f(s+h) - f(s) - \sum_{n=1}^k \frac{h^n}{n!} f^{(n)}(s) \right| &= \frac{|h|^k}{k!} |f^{(k)}(s+\theta h) - f^{(k)}(s)| \\ &< \varepsilon |h|^k, \end{aligned}$$

as required. ■

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function on a closed interval $[a, b]$. We say that f is *continuously differentiable* on $[a, b]$ if the derivative $f'(x)$ of f exists for all x satisfying $a < x < b$, the one-sided derivatives

$$\begin{aligned} f'(a) &= \lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h}, \\ f'(b) &= \lim_{h \rightarrow 0^-} \frac{f(b+h) - f(b)}{h} \end{aligned}$$

exist at the endpoints of $[a, b]$, and the function f' is continuous on $[a, b]$.

If $f: [a, b] \rightarrow \mathbb{R}$ is continuous, and if $F(x) = \int_a^x f(t) dt$ for all $x \in [a, b]$ then the one-sided derivatives of F at the endpoints of $[a, b]$ exist, and

$$\lim_{h \rightarrow 0^+} \frac{F(a+h) - F(a)}{h} = f(a), \quad \lim_{h \rightarrow 0^-} \frac{F(b+h) - F(b)}{h} = f(b).$$

One can verify these results by adapting the proof of the Fundamental Theorem of Calculus.

Proposition 7.13 *Let f be a continuously differentiable real-valued function on the interval $[a, b]$. Then*

$$\int_a^b \frac{df(x)}{dx} dx = f(b) - f(a)$$

Proof Define $g: [a, b] \rightarrow \mathbb{R}$ by

$$g(x) = f(x) - f(a) - \int_a^x \frac{df(t)}{dt} dt.$$

Then $g(a) = 0$, and

$$\frac{dg(x)}{dx} = \frac{df(x)}{dx} - \frac{d}{dx} \left(\int_a^x \frac{df(t)}{dt} dt \right) = 0$$

for all x satisfying $a < x < b$, by the Fundamental Theorem of Calculus. Now it follows from the Mean Value Theorem (Theorem 7.2) that there exists some s satisfying $a < s < b$ for which $g(b) - g(a) = (b-a)g'(s)$. We deduce therefore that $g(b) = 0$, which yields the required result. ■

Corollary 7.14 (Integration by Parts) *Let f and g be continuously differentiable real-valued functions on the interval $[a, b]$. Then*

$$\int_a^b f(x) \frac{dg(x)}{dx} dx = f(b)g(b) - f(a)g(a) - \int_a^b g(x) \frac{df(x)}{dx} dx.$$

Proof This result follows from Proposition 7.13 on integrating the identity

$$f(x) \frac{dg(x)}{dx} = \frac{d}{dx} (f(x)g(x)) - g(x) \frac{df(x)}{dx}. \quad \blacksquare$$

Corollary 7.15 (Integration by Substitution) *Let $u: [a, b] \rightarrow \mathbb{R}$ be a continuously differentiable monotonically increasing function on the interval $[a, b]$, and let $c = u(a)$ and $d = u(b)$. Then*

$$\int_c^d f(x) dx = \int_a^b f(u(t)) \frac{du(t)}{dt} dt.$$

for all continuous real-valued functions f on $[c, d]$.

Proof Let F and G be the functions on $[a, b]$ defined by

$$F(x) = \int_c^{u(x)} f(y) dy, \quad G(x) = \int_a^x f(u(t)) \frac{du(t)}{dt} dt.$$

Then $F(a) = 0 = G(a)$. Moreover $F(x) = H(u(x))$, where

$$H(s) = \int_c^s f(y) dy,$$

and $H'(s) = f(s)$ for all $s \in [a, b]$. Using the Chain Rule and the Fundamental Theorem of Calculus, we deduce that

$$F'(x) = H'(u(x))u'(x) = f(u(x))u'(x) = G'(x)$$

for all $x \in (a, b)$. On applying the Mean Value Theorem (Theorem 7.2) to the function $F - G$ on the interval $[a, b]$, we see that $F(b) - G(b) = F(a) - G(a) = 0$. Thus $H(d) = F(b) = G(b)$, which yields the required identity. \blacksquare

Proposition 7.16 (Taylor's Theorem with Integral Remainder) *Let s and h be real numbers, and let f be a function whose first k derivatives are continuous on an interval containing s and $s + h$. Then*

$$f(s + h) = f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + \frac{h^k}{(k-1)!} \int_0^1 (1-t)^{k-1} f^{(k)}(s + th) dt.$$

Proof Let

$$r_m(s, h) = \frac{h^m}{(m-1)!} \int_0^1 (1-t)^{m-1} f^{(m)}(s+th) dt$$

for $m = 1, 2, \dots, k-1$. Then

$$r_1(s, h) = h \int_0^1 f'(s+th) dt = \int_0^1 \frac{d}{dt} f(s+th) dt = f(s+h) - f(s).$$

Let m be an integer between 1 and $k-2$. It follows from the rule for Integration by Parts (Corollary 7.14) that

$$\begin{aligned} r_{m+1}(s, h) &= \frac{h^{m+1}}{m!} \int_0^1 (1-t)^m f^{(m+1)}(s+th) dt \\ &= \frac{h^m}{m!} \int_0^1 (1-t)^m \frac{d}{dt} (f^{(m)}(s+th)) dt \\ &= \frac{h^m}{m!} [(1-t)^m f^{(m)}(s+th)]_0^1 \\ &\quad - \frac{h^m}{m!} \int_0^1 \frac{d}{dt} ((1-t)^m) f^{(m)}(s+th) dt \\ &= -\frac{h^m}{m!} f^{(m)}(s) + \frac{h^m}{(m-1)!} \int_0^1 (1-t)^{m-1} f^{(m)}(s+th) dt \\ &= r_m(s, h) - \frac{h^m}{m!} f^{(m)}(s). \end{aligned}$$

Thus

$$r_m(s, h) = \frac{h^m}{m!} f^{(m)}(s) + r_{m+1}(s, h)$$

for $m = 1, 2, \dots, k-1$. It follows by induction on k that

$$\begin{aligned} f(s+h) &= f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + r_k(s, h) \\ &= f(s) + \sum_{n=1}^{k-1} \frac{h^n}{n!} f^{(n)}(s) + \frac{h^k}{(k-1)!} \int_0^1 (1-t)^{k-1} f^{(k)}(s+th) dt, \end{aligned}$$

as required. ■