MA232A—Euclidean and Non-Euclidean Geometry School of Mathematics, Trinity College Michaelmas Term 2017 Section 7: Möbius Transformations of the Riemann Sphere

D. R. Wilkins

Copyright © David R. Wilkins 2015–2018

Contents

7	Möł	oius Transformations of the Riemann Sphere	76
	7.1	The Riemann Sphere	76
	7.2	Möbius Transformations	76
	7.3	Inversions of the Riemann Sphere	80
	7.4	Möbius Transformations representing Rotations	82
	7.5	The Action of Möbius Transformations on the Riemann Sphere	87

7 Möbius Transformations of the Riemann Sphere

7.1 The Riemann Sphere

The *Riemann sphere* \mathbb{P}^1 may be defined as the set $\mathbb{C} \cup \{\infty\}$ obtained by augmenting the system \mathbb{C} of complex numbers with an additional element, denoted by ∞ , where ∞ is not itself a complex number, but is an additional element added to the set, with the additional conventions that

$$z + \infty = \infty$$
, $\infty \times \infty = \infty$, $\frac{z}{\infty} = 0$ and $\frac{\infty}{z} = \infty$

for all complex numbers z, and

$$z \times \infty = \infty$$
, and $\frac{z}{0} = \infty$

for all non-zero complex numbers z. The symbol ∞ cannot be added to, or subtracted from, itself. Also 0 and ∞ cannot be divided by themselves.

Note that, because the sum of two elements of \mathbb{P}^1 is not defined for every single pair of elements of \mathbb{P}^1 , this set cannot be regarded as constituting a group under the operation of addition. Similarly its non-zero elements cannot be regarded as constituting a group under multiplication. In particular, the Riemann sphere cannot be regarded as constituting a field.

The following proposition follows directly from Proposition 6.1.

Proposition 7.1 Let $\sigma: \mathbb{P}^1 \to \mathbb{R}^3$ be the mapping from the Riemann sphere \mathbb{P}^1 to \mathbb{R}^3 defined such that $\sigma(\infty) = (0, 0, -1)$ and

$$\sigma(x+y\sqrt{-1}) = \left(\frac{2x}{1+x^2+y^2}, \frac{2y}{1+x^2+y^2}, \frac{1-x^2-y^2}{1+x^2+y^2}\right)$$

for all real numbers x and y. Then σ maps \mathbb{P}^1 injectively and surjectively onto the unit sphere S^2 in \mathbb{R}^3 . Moreover if (u, v, w) is a point of S^2 distinct from (0, 0, -1) then $(u, v, w) = \sigma(x + y\sqrt{-1})$, where

$$x = \frac{u}{w+1}$$
 and $y = \frac{v}{w+1}$.

7.2 Möbius Transformations

Definition Let a, b, c and d be complex numbers satisfying $ad - bc \neq 0$. The *Möbius transformation* $\mu_{a,b,c,d}: \mathbb{P}^1 \to \mathbb{P}^1$ with coefficients a, b, c and d is defined to be the function from the Riemann sphere \mathbb{P}^1 to itself determined by the following properties:

$$\mu_{a,b,c,d}(z) = \frac{az+b}{cz+d}$$

for all complex numbers z for which $cz + d \neq 0$; $\mu_{a,b,c,d}(-d/c) = \infty$ and $\mu_{a,b,c,d}(\infty) = a/c$ if $c \neq 0$; $\mu_{a,b,c,d}(\infty) = \infty$ if c = 0.

Note that the requirement in the above definition of a Möbius transformation that its coefficients a, b, c and d satisfy the condition $ad - bc \neq 0$ ensures that there is no complex number for which az + b and cz + d are both zero.

Let A be a non-singular 2×2 matrix whose coefficients are complex numbers, and let

$$A = \left(\begin{array}{cc} a & b \\ c & d \end{array}\right).$$

We denote by μ_A the Möbius transformation $\mu_{a,b,c,d}$ with coefficients a, b, c, d, defined so that

$$\mu_A(z) = \begin{cases} \frac{az+b}{cz+d} & \text{if } cz+d \neq 0; \\ \infty & \text{if } c \neq 0 \text{ and } z = -d/c; \end{cases}$$
$$\mu_A(\infty) = \begin{cases} \frac{a}{c} & \text{if } c \neq 0; \\ \infty & \text{if } c = 0. \end{cases}$$

The following result exemplifies the reason for representing the coefficients of a Möbius transformation in the form of a matrix.

Proposition 7.2 The composition of two Möbius transformations is a Möbius transformation. Specifically let A and B be non-singular 2×2 matrices with complex coefficients, and let μ_A and μ_B be the corresponding Möbius transformations of the Riemann sphere. Then the composition $\mu_A \circ \mu_B$ of these Möbius transformations is the Möbius transformation μ_{AB} of the Riemann sphere determined by the product AB of the matrices A and B.

Proof Let

$$A = \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix},$$

and let

$$AB = \left(\begin{array}{cc} a_3 & b_3 \\ c_3 & d_3 \end{array}\right).$$

Then

$$a_3 = a_1a_2 + b_1c_2, \quad b_3 = a_1b_2 + b_1d_2,$$

 $c_3 = c_1a_2 + d_1c_2 \quad \text{and} \quad d_3 = c_1b_2 + d_1d_2$

The definitions of Möbius transformations determined by non-singular 2×2 matrices ensure that

$$\mu_A(z) = \frac{a_1 z + b_1}{c_1 z + d_1}$$

whenever $c_1 z + d_1 \neq 0$ and

$$\mu_B(z) = \frac{a_2 z + b_2}{c_2 z + d_2}$$

whenever $c_2 z + d_2 \neq 0$.

First suppose that z is a complex number for which $c_2 z + d_2 \neq 0$. Then

$$(a_1\mu_B(z) + b_1)(c_2z + d_2) = a_1(a_2z + b_2) + b_1(c_2z + d_2)$$

= $a_3z + b_3$,
 $(c_1\mu_B(z) + d_1)(c_2z + d_2) = c_1(a_2z + b_2) + d_1(c_2z + d_2)$
= $c_3z + d_3$.

It follows that if $c_2 z + d_2 \neq 0$ and $c_1 \mu_B(z) + d_1 \neq 0$ then

$$\mu_A(\mu_B(z)) = \frac{a_1\mu_B(z) + b_1}{c_1\mu_B(z) + d_1} = \frac{a_3z + b_3}{c_3z + d_3} = \mu_{AB}(z).$$

If $c_2 z + d_2 \neq 0$ but $c_1 \mu_B(z) + d_1 = 0$ then $c_3 z + d_3 = 0$ and

$$\mu_A(\mu_B(z)) = \infty = \mu_{AB}(z).$$

We conclude that $\mu_A(\mu_B(z)) = \mu_{AB}(z)$ for all complex numbers z satisfying $c_2 z + d_2 \neq 0$.

Next suppose that z is a complex number for which $c_2z + d_2 = 0$. Now the definition of Möbius transformations requires that $a_2d_2 - b_2c_2 \neq 0$. It follows that c_2 and d_2 cannot both be equal to zero. Thus if $c_2z + d_2 = 0$ then either $z = d_2 = 0$ and $c_2 \neq 0$ or else z, c_2 and d_2 are all non-zero. Thus, in all cases where $c_2z + d_2 = 0$, the coefficient c_2 of the Möbius transformation is non-zero and $z = -d_2/c_2$. Also the equations $a_2z + b_2 = 0$ and $c_2z + d_2 = 0$ cannot both be satisfied, because $a_2d_2 - b_2c_2 \neq 0$, and therefore $a_2z + b_2 \neq 0$. Now the equations determining a_3 , b_3 , c_3 and d_3 ensure that if $c_2 z + d_2 = 0$ then

$$\begin{aligned} c_2(a_3z+b_3) &= -d_2a_3 + c_2b_3 \\ &= c_2(a_1b_2 + b_1d_2) - d_2(a_1a_2 + b_1c_2) \\ &= a_1(b_2c_2 - a_2d_2) \\ &= a_1c_2(a_2z + b_2) \\ c_2(c_3z+d_3) &= -d_2c_3 + c_2d_3 \\ &= c_2(c_1b_2 + d_1d_2) - d_2(c_1a_2 + d_1c_2) \\ &= c_1(b_2c_2 - a_2d_2) \\ &= c_1c_2(a_2z + b_2), \end{aligned}$$

and therefore

$$a_3z + b_3 = a_1(a_2z + b_2)$$
 and $c_3z + d_3 = c_1(a_2z + b_2)$,

Thus if $c_2 z + d_2 = 0$ and $c_1 \neq 0$ then $c_3 z + d_3 \neq 0$ and

$$\mu_{AB}(z) = \frac{a_3 z + b_3}{c_3 z + d_3} = \frac{a_1}{c_1} = \mu_A(\infty) = \mu_A(\mu_B(z)).$$

And if $c_2 z + d_2 = 0$ and $c_1 = 0$ then $c_3 z + d_3 = 0$ and

$$\mu_{AB}(z) = \infty = \mu_A(\infty) = \mu_A(\mu_B(z)).$$

Thus $\mu_{AB}(z) = \mu_A(\mu_B(z))$ in all cases for which $c_2 z + d_2 = 0$.

It remains to show that $\mu_{AB}(\infty) = \mu_A(\mu_B(\infty))$. If $c_2 \neq 0$ (so that $\mu_B(\infty) = a_2/c_2$) and $c_1\mu_B(\infty) + d_2 \neq 0$ then

$$\mu_A(\mu_B(\infty)) = \frac{a_1\mu_B(\infty) + b_1}{c_1\mu_B(\infty) + d_1} = \frac{a_1a_2 + b_1c_2}{c_1a_2 + d_1c_2} = \frac{a_3}{c_3} = \mu_{AB}(\infty).$$

If $c_2 \neq 0$ and $c_1\mu_B(\infty) + d_2 = 0$ then $c_3 = c_1a_2 + d_1c_2 = 0$, because $\mu_B(\infty) = a_2/c_2$, and therefore

$$\mu_A(\mu_B(\infty)) = \infty = \mu_{AB}(\infty).$$

If $c_1 = c_2 = 0$ then $\mu_B(\infty) = \infty$ and therefore

$$\mu_A(\mu_B(\infty)) = \mu_A(\infty) = \infty = \mu_{AB}(\infty).$$

If $c_2 = 0$ and $c_1 \neq 0$ then $a_3 = a_1a_2$, $c_3 = c_1a_2$ and $a_2 \neq 0$ (because $a_2d_2 - b_2c_2 \neq 0$), and therefore

$$\mu_A(\mu_B(\infty)) = \mu_A(\infty) = \frac{a_1}{c_1} = \frac{a_3}{c_3} = \mu_{AB}(\infty).$$

We conclude that $\mu_A(\mu_B(\infty)) = \mu_{AB}(\infty)$ in all cases. This completes the proof.

Corollary 7.3 Let a, b, c and d be complex numbers satisfying $ad - bc \neq 0$, and let $\mu_{a,b,c,d}: \mathbb{P}^1 \to \mathbb{P}^1$ denote the Möbius transformation of the Riemann sphere \mathbb{P}^1 defined such that $\mu_{a,b,c,d}(z) = \frac{az+b}{cz+d}$ if $z \in \mathbb{C}$ and $cz+d\neq 0$, $\mu_{a,b,c,d}(-d/c) = \infty$ and $\mu_{a,b,c,d}(\infty) = a/c$ if $c\neq 0$, and $\mu_{a,b,c,d}(\infty) = \infty$ if c = 0. Then the mapping $\mu_{a,b,c,d}: \mathbb{P}^1 \to \mathbb{P}^1$ is invertible, and its inverse is the Möbius transformation $\mu_{d,-b,-c,a}: \mathbb{P}^1 \to \mathbb{P}^1$, where $\mu_{d,-b,-c,a}(z) = \frac{dz-b}{a-cz}$ if $z \in \mathbb{C}$ and $a - cz \neq 0$, $\mu_{d,-b,-c,a}(a/c) = \infty$ and $\mu_{d,-b,-c,a}(\infty) = -d/c$ if $c\neq 0$, and $\mu_{d,-b,-c,a}(\infty) = \infty$ if c = 0.

Proof If the coefficients a, b, c and d of a Möbius transformation are all multiplied by a non-zero complex number then this does not change the Möbius transformation represented by those coefficients. It follows that we may assume, without loss of generality, that ad - bc = 1. Let

$$A = \left(\begin{array}{cc} a & b \\ c & d \end{array}\right),$$

where ad - bc = 1. Then

$$A^{-1} = \left(\begin{array}{cc} d & -b \\ -c & a \end{array}\right).$$

The result therefore follows directly on applying Proposition 7.2.

7.3 Inversions of the Riemann Sphere

Let S^2 denote the unit sphere in \mathbb{R}^3 , defined so that

$$S^2 = \{(u,v,w) \in \mathbb{R}^3 : u^2 + v^2 + w^2 = 1\},$$

and let us refer to the points (0, 0, 1) and (0, 0, -1) as the North Pole and South Pole respectively. Let E denote the Equatorial Plane in \mathbb{R}^3 , consisting of those points whose Cartesian coordinates are of the form (x, y, 0), where x and y are real numbers.

Stereographic projection from the South Pole maps each point (u, v, w)of the unit sphere S^2 distinct from the South Pole to the point (x, y, 0) of the equatorial plane E for which

$$x = \frac{u}{w+1}$$
 and $y = \frac{v}{w+1}$.

Moreover a point (x, y, 0) of the Equatorial Plane E is the image under stereographic projection from the South Pole of the point (u, v, w) of the unit sphere S^2 for which

$$u = \frac{2x}{1 + x^2 + y^2}, \quad v = \frac{2y}{1 + x^2 + y^2}, \quad w = \frac{1 - x^2 - y^2}{1 + x^2 + y^2}.$$

We can also stereographically project from the North Pole. Note that, given a point in the Equatorial Plane, reflection in that Equatorial Plane will interchange the points of the sphere corresponding to it under stereographic projection from the North and South Poles. Thus a point (u, v, w) of the unit sphere S^2 distinct from the North Pole corresponds under stereographic projection to the point (x, y, 0) of the Equatorial Plane E for which

$$x = \frac{u}{1-w}$$
 and $y = \frac{v}{1-w}$.

In the other direction, a point (x, y, 0) of the Equatorial Plane E corresponds under stereographic projection from the North Pole to the point (u, v, w) of the unit sphere S^2 for which

$$u = \frac{2x}{1+x^2+y^2}, \quad v = \frac{2y}{1+x^2+y^2}, \quad w = \frac{x^2+y^2-1}{1+x^2+y^2}.$$

Proposition 7.4 Let O denote the origin (0,0,0) of the Equatorial Plane E, where

$$E = \{ (x, y, z) \in \mathbb{R}^3 : z = 0 \},\$$

and let A be a point (x, y, 0) of E distinct from the origin O. Let C be the point on the unit sphere S^2 that corresponds to A under stereographic projection from the North Pole (0, 0, 1), and let B be the point of the Equatorial Plane E that corresponds to C under stereographic projection from the South Pole. Then B = (p, q, 0), where

$$p = \frac{x}{x^2 + y^2}$$
 and $q = \frac{y}{x^2 + y^2}$.

Thus the points O, A and B are collinear, and the points A and B lie on the same side of the origin O. Also the distances |OA| and |OB| of the points A and B from the origin satisfy $|OA| \times |OB| = 1$.

Proof Let (x, y, 0) be a point of the Equatorial plane E distinct from the origin. This point is the image, under stereographic projection from the North Pole (0, 0, 1) of the point (u, v, w) of the unit sphere S^2 for which

$$u = \frac{2x}{1+x^2+y^2}, \quad v = \frac{2y}{1+x^2+y^2}, \quad w = \frac{x^2+y^2-1}{1+x^2+y^2},$$

This point then gets mapped under stereographic projection from the South Pole to the point (p, q, 0) of the Equatorial Plane E for which

$$p = \frac{u}{w+1}$$
 and $q = \frac{v}{w+1}$

Now

$$w + 1 = \frac{2(x^2 + y^2)}{1 + x^2 + y^2}.$$

It follows that

$$p = \frac{x}{x^2 + y^2}$$
 and $q = \frac{y}{x^2 + y^2}$.

Finally we note that O, A and B are collinear, where 0 = (0, 0, 0), A = (x, y, 0) and B = (p, q, 0), and the points A and B lie on the same side of the origin O. Also

$$|OA| = \sqrt{x^2 + y^2}$$
, and $|OB| = \frac{1}{\sqrt{x^2 + y^2}}$,

and therefore $|OA| \times |OB| = 1$, as required.

7.4 Möbius Transformations representing Rotations

Let a and b be complex numbers satisfying the equation $|a|^2 + |b|^2 = 1$, and let $\mu: \mathbb{P}^1 \to \mathbb{P}^1$ be the Möbius Transformation of the Riemann sphere defined such that

$$\mu(z) = \frac{az+b}{-\overline{b}z+\overline{a}}$$

when $z \neq \overline{a}/\overline{b}$, $\mu(\overline{a}/\overline{b}) = \infty$ and $\mu(\infty) = -a/\overline{b}$. (Here \overline{a} and \overline{b} denote the complex conjugates of the complex numbers a and b respectively.) We denote the complex number $\sqrt{-1}$ by i, as is customary.

Let u_0 , v_0 and w_0 are real numbers satisfying $u_0^2 + v_0^2 + w_0^2 = 1$ and $w_0 \neq -1$. Then the point (u_0, v_0, w_0) of the unit sphere in \mathbb{R}^3 corresponds, under stereographic projection from (0, 0, -1), to the complex number z_0 for which

$$z_0 = \frac{u_0 + iv_0}{w_0 + 1}.$$

Let $z_1 = \mu(z_0)$. Then

$$z_1 = \frac{a(u_0 + iv_0) + b(w_0 + 1)}{-\overline{b}(u_0 + iv_0) + \overline{a}(w_0 + 1)}.$$

Now there exists a point (u_1, v_1, w_1) of the unit sphere in \mathbb{R}^3 which corresponds under stereographic projection from the point (0, 0, -1) to the complex number z_1 . The Cartesian coordinates u_1, v_1 and w_1 of this point satisfy the equation $u_1^2 + v_1^2 + w_1^2 = 1$ and their relationship to the complex number z_1 is expressed by the following equations:

$$z_1 = \frac{u_1 + iv_1}{w_1 + 1},$$
$$u_1 = \frac{2\operatorname{Re}[z_1]}{|z_1|^2 + 1}, \quad v_1 = \frac{2\operatorname{Im}[z_1]}{|z_1|^2 + 1}, \quad w_1 = \frac{2}{|z_1|^2 + 1} - 1.$$

We seek to express the values of u_1 , v_1 and w_1 in terms of u_0 , v_0 and w_0 . Now

$$|a(u_0 + iv_0) + b(w_0 + 1)|^2$$

= $(a(u_0 + iv_0) + b(w_0 + 1))(\overline{a}(u_0 - iv_0) + \overline{b}(w_0 + 1))$
= $|a|^2(u_0^2 + v_0^2) + |b|^2(w_0 + 1)^2 + 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)](w_0 + 1).$

But $u_0^2 + v_0^2 = 1 - w_0^2 = (w_0 + 1)(1 - w_0)$. It follows that

$$|a(u_0 + iv_0) + b(w_0 + 1)|^2$$

= $(|a|^2(1 - w_0) + |b|^2(w_0 + 1) + 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)])(w_0 + 1)$
= $(1 - (|a|^2 - |b|^2)w_0 + 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)])(w_0 + 1).$

Similarly

$$|-\overline{b}(u_0+iv_0)+\overline{a}(w_0+1)|^2$$

= $(1+(|a|^2-|b|^2)w_0-2\operatorname{Re}[a\overline{b}(u_0+iv_0)])(w_0+1).$

It follows from these identities that

$$|z_1|^2 = \frac{1 - (|a|^2 - |b|^2)w_0 + 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)]}{1 + (|a|^2 - |b|^2)w_0 - 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)]},$$

and thus

$$|z_1|^2 + 1 = \frac{2}{1 + (|a|^2 - |b|^2)w_0 - 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)]}$$

Also

$$z_{1} = \frac{a(u_{0} + iv_{0}) + b(w_{0} + 1)}{-\overline{b}(u_{0} + iv_{0}) + \overline{a}(w_{0} + 1)}$$

=
$$\frac{(a(u_{0} + iv_{0}) + b(w_{0} + 1))(-b(u_{0} - iv_{0}) + a(w_{0} + 1))}{|-\overline{b}(u_{0} + iv_{0}) + \overline{a}(w_{0} + 1)|^{2}}$$

Moreover

$$(a(u_0 + iv_0) + b(w_0 + 1))(-b(u_0 - iv_0) + a(w_0 + 1))$$

= $(a^2(u_0 + iv_0) - b^2(u_0 - iv_0))(w_0 + 1)$
+ $ab((w_0 + 1)^2 - u_0^2 - v_0^2)$
= $((a^2 - b^2)u_0 + i(a^2 + b^2)v_0 + 2abw_0)(w_0 + 1)$

It follows that

$$z_1 = \frac{(a^2 - b^2)u_0 + i(a^2 + b^2)v_0 + 2abw_0}{1 + (|a|^2 - |b|^2)w_0 - 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)]},$$

and thus

$$u_1 + iv_1 = \frac{2z_1}{|z_1|^2 + 1}$$

= $(a^2 - b^2)u_0 + i(a^2 + b^2)v_0 + 2abw_0$

and

$$w_1 = \frac{2}{|z_1|^2 + 1} - 1$$

= $(|a|^2 - |b|^2)w_0 - 2\operatorname{Re}[a\overline{b}(u_0 + iv_0)]$

Now $|a|^2 + |b|^2 = 1$ and $b \neq 0$. It follows that |a| < 1, and thus $\operatorname{Re}[a] = \cos \frac{1}{2}\theta$ for some real number θ satisfying $0 < \theta < 2\pi$. Let ℓ , m and n be real numbers determined so that

$$a = \cos \frac{1}{2}\theta + in \sin \frac{1}{2}\theta$$
 and $b = (m - i\ell) \sin \frac{1}{2}\theta$.

Then

$$1 = |a|^{2} + |b|^{2} = \cos^{2} \frac{1}{2}\theta + (\ell^{2} + m^{2} + n^{2})\sin^{2} \frac{1}{2}\theta,$$

and therefore

$$\ell^2 + m^2 + n^2 = 1.$$

Then

$$\begin{aligned} |a|^2 - |b|^2 &= \cos^2 \frac{1}{2}\theta + (n^2 - \ell^2 - m^2) \sin^2 \frac{1}{2}\theta \\ &= \cos \theta + n^2 (1 - \cos \theta), \\ a^2 - b^2 &= \cos^2 \frac{1}{2}\theta + (\ell^2 - m^2 - n^2 + 2i\ell m) \sin^2 \frac{1}{2}\theta + 2in \sin \frac{1}{2}\theta \cos \frac{1}{2}\theta \\ &= \cos \theta + (\ell^2 + i\ell m) (1 - \cos \theta) + in \sin \theta \\ a^2 + b^2 &= \cos^2 \frac{1}{2}\theta + (m^2 - \ell^2 - n^2 - 2i\ell m) \sin^2 \frac{1}{2}\theta + 2in \sin \frac{1}{2}\theta \cos \frac{1}{2}\theta \end{aligned}$$

$$= \cos \theta + (m^2 - i\ell m)(1 - \cos \theta) + in \sin \theta$$

$$2a\overline{b} = 2(m + i\ell) \sin \frac{1}{2}\theta \cos \frac{1}{2}\theta - 2(\ell - im)n \sin^2 \frac{1}{2}\theta$$

$$= (m + i\ell) \sin \theta - (\ell - im)n(1 - \cos \theta)$$

$$2ab = 2(m - i\ell) \sin \frac{1}{2}\theta \cos \frac{1}{2}\theta + 2(\ell + im)n \sin^2 \frac{1}{2}\theta$$

$$= (m - i\ell) \sin \theta + (\ell + im)n(1 - \cos \theta)$$

It follows that

$$u_{1} = \operatorname{Re}[a^{2} - b^{2}]u_{0} - \operatorname{Im}[a^{2} + b^{2}]v_{0} + 2\operatorname{Re}[ab]w_{0}$$

$$= (mw_{0} - nv_{0})\sin\theta + u_{0}\cos\theta + (\ell u_{0} + mv_{0} + nw_{0})\ell(1 - \cos\theta)$$

$$v_{1} = \operatorname{Im}[a^{2} - b^{2}]u_{0} + \operatorname{Re}[a^{2} + b^{2}]v_{0} + 2\operatorname{Im}[ab]w_{0}$$

$$= (nu_{0} - \ell w_{0})\sin\theta + v_{0}\cos\theta + (\ell u_{0} + mv_{0} + nw_{0})m(1 - \cos\theta)$$

$$w_{1} = (|a|^{2} - |b|^{2})w_{0} - 2\operatorname{Re}[a\overline{b}(u_{0} + iv_{0})]$$

$$= (\ell v_{0} - mu_{0})\sin\theta + w_{0}\cos\theta + (\ell u_{0} + mv_{0} + nw_{0})n(1 - \cos\theta)$$

Let

$$\mathbf{r}_0 = (u_0, v_0, w_0), \quad \mathbf{r}_1 = (u_1, v_1, w_1) \text{ and } \mathbf{L} = (\ell, m, n).$$

Then the vectors \mathbf{r}_0 , \mathbf{r}_1 and \mathbf{L} are of unit length, and

$$\begin{aligned} \mathbf{r}_1 &= \ \cos\theta \, \mathbf{r}_0 + (\mathbf{r}_0 \, . \, \mathbf{L})(1 - \cos\theta) \, \mathbf{L} + \sin\theta \mathbf{L} \times \mathbf{r}_0 \\ &= \ (\mathbf{r}_0 \, . \, \mathbf{L}) \mathbf{L} + \cos\theta \, (\mathbf{r}_0 - (\mathbf{r}_0 \, . \, \mathbf{L}) \mathbf{L} + \sin\theta \, \mathbf{L} \times \mathbf{r}_0. \end{aligned}$$

Interpreting this formula geometrically, we see that the point (u_1, v_1, w_1) of the unit sphere in \mathbb{R}^3 is the image of the point (u_0, v_0, w_0) under a rotation through an angle θ about the axis passing through the origin in the direction of the vector **L**.

Proposition 7.5 Let a, b, c and d be complex numbers satisfying ad-bc = 1, and let $\mu: \mathbb{P}^1 \to \mathbb{P}^1$ be the Möbius transformation of the Riemann sphere defined such that

$$\mu(z) = \frac{az+b}{cz+d} \quad when \ cz+d \neq 0,.$$

 $\mu(-d/c) = \infty$ and $\mu(\infty) = a/c$ in the case $c \neq 0$ and $\mu(\infty) = \infty$ in the case c = 0. Then the Möbius transformation represents a rotation of the unit sphere in \mathbb{R}^3 if and only if $d = \overline{a}$ and $c = -\overline{b}$.

Proof The calculations undertaken in this subsection show that if $b \neq 0$, $d = \overline{a}$ and $c = -\overline{b}$ then the Möbius transformation $\mu: \mathbb{P}^1 \to \mathbb{P}^1$ corresponds to a rotation of the unit sphere in \mathbb{R}^3 . The same is true in the case when b = 0. Indeed in that case the conditions $d = \overline{a}$ and $c = -\overline{b}$ ensure that b = c = 0, |a| = 1, and these conditions ensure that the Möbius μ implements a rotation of the unit sphere about the direction (0, 0, 1) through an angle θ , where

$$a = \cos \frac{1}{2}\theta + \sqrt{-1}\sin \frac{1}{2}\theta.$$

Thus, in all cases where $d = \overline{a}$, $c = -\overline{b}$ and $|a|^2 + |b|^2 = 1$, the Möbius transformation μ of the Riemann sphere corresponds to a rotation of the unit sphere in \mathbb{R}^3 .

Now every rotation about the origin in \mathbb{R}^3 is a rotation about a fixed axis through a given angle. It follows that, given any rotation of the sphere, there are values of the complex numbers a and b for which the Möbius transformation μ implements the given rotation with $d = \overline{a}$ and $c = -\overline{b}$. Moreover a Möbius transformation with coefficients a', b', c', d' implements the same transformation of the sphere as the Möbius transformation μ if and only if either a', b', c' and d' are respectively equal to a, b, c and d or else a', b', c'and d' are respectively equal to -a, -b, -c and -d. The result follows.

Let

$$\operatorname{SU}(2) = \left\{ \left(\begin{array}{cc} a & b \\ -\overline{b} & \overline{a} \end{array} \right) : a, b \in \mathbb{C} \text{ and } |a|^2 + |b|^2 = 1 \right\}.$$

Then the 2 × 2 matrices belonging to the set SU(2) constitute a group with respect to the operation of matrix multiplication. Morever a 2 × 2 matrix with complex coefficients belongs to the group SU(2) if and only if it is a unitary matrix whose determinant is equal to one. Proposition 7.5 ensures that every 2 × 2 unitary matrix with determinant equal to one determines a corresponding rotation of three-dimensional space \mathbb{R}^3 . We obtain in this way a two-to-one homomorphism from the group SU(2) to the rotation group SO(3) of 3-dimensional space.

Remark The homomorphism from SU(2) to SO(3) whose existence is guaranteed by Proposition 7.5 can also be described using properties of quaternions. Independently of one another, Arthur Cayley and Sir William Rowan Hamilton discovered how to represent rotations of three-dimensional space using quaternions. (Cayley's account appeared in print in 1845 before Hamilton's account, read at a meeting of the Royal Irish Academy in 1844 but published in 1847.)

The homomorphism between these matrix groups gives rise to the fundamental properties of spin in quantum mechanics, where the traditional account is expressed in terms of *Pauli matrices*.

7.5 The Action of Möbius Transformations on the Riemann Sphere

Proposition 7.6 Let $\zeta_1, \zeta_2, \zeta_3$ be distinct points of the Riemann sphere \mathbb{P}^1 , and let $\omega_1, \omega_2, \omega_3$ also be distinct points of \mathbb{P}^1 . Then there exists a Möbius transformation $\mu: \mathbb{P}^1 \to \mathbb{P}^1$ of the Riemann sphere with the property that $\mu(\zeta_j) = \omega_j$ for j = 1, 2, 3.

Proof The composition of two Möbius transformations of the Riemann sphere is itself a Möbius transformation of the Riemann sphere (Proposition 7.2). Also the inverse of any Möbius transformation of the Riemann sphere is itself a Möbius transformation (Corollary 7.3). It follows that the Möbius transformations of the Riemann sphere constitute a group under the operation of composition of transformations.

Next we note that permution of the elements 0, 1 and ∞ of the Riemann sphere can be effected by a suitable Möbius transformation. Indeed the Möbius transformation $z \mapsto 1-z$ transposes 0 and 1 whilst fixing ∞ , and the Möbius transformation $z \mapsto -1/(z-1)$ cyclicly permutes 0, 1 and ∞ . It follows that any permutation of 0, 1 and ∞ may be effected by the action of some Möbius transformation.

Next we show that there exists a Möbius transformation $\mu_1: \mathbb{P}^1 \to \mathbb{P}_2$ with the property that $\mu_1(\zeta_1) = 0$, $\mu_1(\zeta_2) = 1$ and $\mu_1(\zeta_3) = \infty$. Suppose first that at least one of the distinct points $\zeta_1, \zeta_2, \zeta_3$ of \mathbb{P}^1 is the point ∞ . Because we have shown that there exist Möbius transformations permuting 0, 1 and ∞ amongst themselves, we may assume in this case, without loss of generality, that $\zeta_3 = \infty$. Let $\zeta_0 = z_0$ and $\zeta_1 = z_1$, where z_0 and z_1 are complex numbers, and let

$$\mu_1(z) = \frac{z - z_0}{z_1 - z_0}.$$

Then $\mu_1(\zeta_0) = \mu_1(z_0) = 0$, $\mu_1(\zeta_1) = \mu_1(z_1) = 1$ and $\mu_1(\infty) = \infty$. The existence of the Möbius transformation μ_1 has thus been verified in the case where at least one of $\zeta_1, \zeta_2, \zeta_3$ is the point ∞ of the Riemann sphere.

Next we consider the case where $\zeta_j = z_j$ for j = 1, 2, 3, where z_1, z_2, z_3 are complex numbers. In this case let μ_1 be the Möbius transformation defined so that

$$\mu_1(z) = \frac{(z-z_1)(z_2-z_3)}{(z-z_3)(z_2-z_1)}$$

for all complex numbers z. Then $\mu_1(\zeta_1) = \mu_1(z_1) = 0$, $\mu_1(\zeta_2) = \mu_1(z_2) = 1$ and $\mu_1(\zeta_3) = \mu_1(z_3) = \infty$. We conclude therefore that, given any distinct points $\zeta_1, \zeta_2, \zeta_3$ of the Riemann sphere, there exists a Möbius transformation μ_1 of the Riemann sphere for which $\mu_1(\zeta_1) = 0$, $\mu_1(\zeta_2) = 1$ and $\mu_1(\zeta_3) = \infty$. Let $\omega_1, \omega_2, \omega_3$ also be distinct points of the Riemann sphere. Then there exists a Möbius transformation μ_2 with the property that $\mu_2(\omega_1) = 0, \mu_2(\omega_2) = 1$ and $\mu_2(\omega_3) = \infty$. Let $\mu: \mathbb{P}^1 \to \mathbb{P}^1$ be the Möbius transformation of the Riemann sphere that is the composition $\mu_2^{-1} \circ \mu_1$ of μ_1 followed by the inverse of μ_2 . Then $\mu(\zeta_j) = \omega_j$ for j = 1, 2, 3, as required.

Proposition 7.7 Let $\rho: \mathbb{P}^1 \to \mathbb{P}^1$ denote the Möbius transformation defined so that $\rho(z) = 1/z$ for all non-zero complex numbers z. Then the mapping ρ preserves the angles between circles and straight lines. contained in the set $\mathbb{C} \setminus \{0\}.$

Proof Let z be a non-zero complex number, and let h be a non-zero complex number satisfying |h| < |z|, and let t be a real number satisfying $-1 \le t \le 1$. Then $z + th \ne 0$, and

$$\left(1+\frac{th}{z}\right)\left(1-\frac{th}{z}\right) = 1 - \frac{t^2h^2}{z^2}.$$

It follows that

$$\frac{1}{z} - \frac{th}{z^2} = \frac{1}{z+th} - \frac{t^2h^2}{z^2(z+th)}$$

and thus

$$\frac{1}{z+th} = \frac{1}{z} - \frac{th}{z^2} + \frac{t^2h^2}{z^2(z+th)}$$

Let θ be a real number, let $k = h(\cos\theta + i\sin\theta)$. Then the directions represented by the complex numbers h and k are at an angle θ to each other. Let $\alpha: (-1, 1) \to \mathbb{C}$ and $\beta: (-1, 1) \to \mathbb{C}$ be the curves defined such that

$$\alpha(t) = \rho(z + th) = \frac{1}{z + th}$$

and

$$\beta(t) = \rho(z + tk) = \frac{1}{z + tk}$$

for all real numbers t satisfying -1 < t < 1. Then the tangent directions to the curve $t \mapsto \alpha(t)$ and $t \mapsto \beta(t)$ at t = 0 are in the directions determined by the complex numbers

$$-\frac{h}{z^2}$$
 and $-\frac{k}{z^2}$.

Moreover

$$-\frac{k}{z^2} = -\frac{h}{z^2}(\cos\theta + i\sin\theta)$$

and therefore the tangent directions to the curves $t \mapsto \alpha(t)$ and $t \mapsto \beta(t)$ at t = 0 are also at an angle θ to each other. The result follows.

Proposition 7.8 Any Möbius transformation of the Riemann sphere maps straight lines and circles to straight lines and circles, and also preserves angles between lines and circles.

Proof Proposition 7.7 ensures that the Möbius transformation that sends z to 1/z for all non-zero complex numbers z is angle-preserving.

The equation of a line or circle in the complex plane can be expressed in the form

$$g|z|^2 + 2\operatorname{Re}[\overline{b}z] + h = 0,$$

where g and h are real numbers, and b is a complex number. Moreover a locus of points in the complex plane satisfying an equation of this form is a circle if $g \neq 0$ and is a line if g = 0.

Let g and h be real constants, let b be a complex constant, and let z = 1/w, where $w \neq 0$ and w satisfies the equation

$$g|w|^2 + 2\operatorname{Re}[\overline{b}w] + h = 0,$$

Then

$$|g|w|^2 + \overline{b}w + b\overline{w} + h = 0,$$

Then

$$g + 2\operatorname{Re}[bz] + h|z|^{2} = g + \overline{b}\overline{z} + bz + h|z|^{2}$$
$$= \frac{1}{|w|^{2}} \left(g|w|^{2} + \overline{b}w + b\overline{w} + h\right) = 0.$$

We deduce from this that the Möbius transformation that sends z to 1/z for all non-zero complex numbers z maps lines and circles to lines and circles.

Let $\mu: \mathbb{P}^1 \to \mathbb{P}^1$ be a Möbius transformation of the Riemann sphere. Then there exist complex numbers a, b, c and d satisfying $ad - bc \neq 0$ such that

$$\mu(z) = \frac{az+b}{cz+d}$$

for all complex numbers z for which $cz + d \neq 0$. The result is immediate when c = 0. We therefore suppose that $c \neq 0$. Then

$$\mu(z) = \frac{az+b}{cz+d} = \frac{a}{c} - \frac{ad-bc}{c} \times \frac{1}{cz+d}$$

when $cz + d \neq 0$. The Möbius transformation μ is thus the composition of three maps that each send circles and straight lines to circles and straight lines and preserve angles between lines and circles, namely the maps

$$z \mapsto cz + d$$
, $z \mapsto \frac{1}{z}$ and $z \mapsto \frac{a}{c} - \frac{(ad - bc)z}{c}$.

Thus the Möbius transformation μ must itself map circles and straight line to circles and straight lines and also preserve angles between lines and circles, as required.