Module MA3427 Algebraic Topology I

D. R. Wilkins

Michaelmas Term 2014

Topological Spaces.

Notions of Continuity.

The concept of continuity is fundamental in large parts of contemporary mathematics. In the nineteenth century, precise definitions of continuity were formulated for functions of a real or complex variable, enabling mathematicians to produce rigorous proofs of fundamental theorems of real and complex analysis, such as the Intermediate Value Theorem, Taylor's Theorem, the Fundamental Theorem of Calculus, and Cauchy's Theorem.

In the early years of the Twentieth Century, the concept of continuity was generalized so as to be applicable to functions between metric spaces, and subsequently to functions between topological spaces.

Topological Spaces.

Definition.

A topological space X consists of a set X together with a collection of subsets, referred to as open sets, such that the following conditions are satisfied:—

- (i) the empty set \emptyset and the whole set X are open sets,
- (ii) the union of any collection of open sets is itself an open set,
- (iii) the intersection of any *finite* collection of open sets is itself an open set.

The collection consisting of all the open sets in a topological space X is referred to as a *topology* on the set X.

Remark.

If it is necessary to specify explicitly the topology on a topological space then one denotes by (X, τ) the topological space whose underlying set is X and whose topology is τ . However if no confusion will arise then it is customary to denote this topological space simply by X.

Subsets of Euclidean Space.

Let X be a subset of *n*-dimensional Euclidean space \mathbb{R}^n . The *Euclidean distance* $|\mathbf{x} - \mathbf{y}|$ between two points \mathbf{x} and \mathbf{y} of X is defined as follows:

$$|\mathbf{x} - \mathbf{y}| = \sqrt{\sum_{i=1}^{n} (x_i - y_i)^2},$$

where

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

and

$$\mathbf{y}=(y_1,y_2,\ldots,y_n).$$

4

The Euclidean distances between any three points x, y and z of X satisfy the *Triangle Inequality*:

$$|\mathbf{x} - \mathbf{z}| \le |\mathbf{x} - \mathbf{y}| + |\mathbf{y} - \mathbf{z}|.$$

A subset V of X is said to be *open* in X if, given any point v of V, there exists some $\delta > 0$ such that

 $\{\mathbf{x} \in X : |\mathbf{x} - \mathbf{v}| < \delta\} \subset V.$

The empty set is also considered to be open in X.

Both \emptyset and X are open sets in X. Also it is not difficult to show that any union of open sets in X is open in X, and that any finite intersection of open sets in X is open in X. (This will be proved in more generality for open sets in metric spaces.) Thus the collection of open sets in a subset X of a Euclidean space \mathbb{R}^n satisfies the topological space axioms. Thus every subset X of \mathbb{R}^n is a topological space with these open sets. This topology on a subset X of \mathbb{R}^n is referred to as the usual topology on X, generated by the Euclidean distance function.

In particular \mathbb{R}^n is itself a topological space.

Open Sets in Metric Spaces.

Definition.

A metric space (X, d) consists of a set X together with a distance function $d: X \times X \rightarrow$ $[0, +\infty)$ on X satisfying the following axioms:

(i)
$$d(x,y) \ge 0$$
 for all $x, y \in X$,

(ii)
$$d(x,y) = d(y,x)$$
 for all $x, y \in X$,

(iii)
$$d(x,z) \leq d(x,y) + d(y,z)$$
 for all $x,y,z \in X$,

(iv) d(x,y) = 0 if and only if x = y.

The quantity d(x, y) should be thought of as measuring the *distance* between the points x and y. The inequality $d(x, z) \leq d(x, y) +$ d(y, z) is referred to as the *Triangle Inequality*. The elements of a metric space are usually referred to as *points* of that metric space.

An *n*-dimensional Euclidean space \mathbb{R}^n is a metric space with with respect to the *Eu*clidean distance function d, defined by

$$d(\mathbf{x}, \mathbf{y}) = |\mathbf{x} - \mathbf{y}| = \sqrt{\sum_{i=1}^{n} (x_i - y_i)^2}$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$. Any subset X of \mathbb{R}^n may be regarded as a metric space whose distance function is the restriction to X of the Euclidean distance function on \mathbb{R}^n defined above.

8

Definition.

Let (X, d) be a metric space. Given a point xof X and $r \ge 0$, the *open ball* $B_X(x, r)$ of radius r about x in X is defined by

$$B_X(x,r) \equiv \{ x' \in X : d(x',x) < r \}.$$

Definition.

Let (X, d) be a metric space. A subset V of X is said to be an *open set* if and only if the following condition is satisfied:

• given any point v of V there exists some $\delta > 0$ such that $B_X(v, \delta) \subset V$.

By convention, we regard the empty set \emptyset as being an open subset of X. (The criterion given above is satisfied vacuously in this case.)

Lemma 1.1 Let X be a metric space with distance function d, and let x_0 be a point of X. Then, for any r > 0, the open ball $B_X(x_0,r)$ of radius r about x_0 is an open set in X.

Proof.

Let $x \in B_X(x_0, r)$. We must show that there exists some $\delta > 0$ such that $B_X(x, \delta) \subset B_X(x_0, r)$. Now $d(x, x_0) < r$, and hence $\delta > 0$, where $\delta = r - d(x, x_0)$. Moreover if $x' \in B_X(x, \delta)$ then

$$d(x', x_0) \le d(x', x) + d(x, x_0) < \delta + d(x, x_0) = r,$$

by the Triangle Inequality, hence $x' \in B_X(x_0, r)$. Thus

$$B_X(x,\delta) \subset B_X(x_0,r),$$

showing that $B_X(x_0, r)$ is an open set, as required.

Proposition 1.2 Let X be a metric space. The collection of open sets in X has the following properties:—

- (i) the empty set \emptyset and the whole set X are both open sets;
- (ii) the union of any collection of open sets is itself an open set;
- (iii) the intersection of any finite collection of open sets is itself an open set.

Proof.

The empty set \emptyset is an open set by convention. Moreover the definition of an open set is satisfied trivially by the whole set X. Thus (i) is satisfied.

Let \mathcal{A} be any collection of open sets in X, and let U denote the union of all the open sets belonging to \mathcal{A} . We must show that U is itself an open set. Let $x \in U$. Then $x \in V$ for some open set V belonging to the collection \mathcal{A} . Therefore there exists some $\delta > 0$ such that $B_X(x, \delta) \subset V$. But $V \subset U$, and thus $B_X(x, \delta) \subset U$. This shows that U is open. Thus (ii) is satisfied.

Finally let $V_1, V_2, V_3, \ldots, V_k$ be a *finite* collection of open sets in X, and let

$$V = V_1 \cap V_2 \cap \cdots \cap V_k.$$

Let $x \in V$. Now $x \in V_j$ for all j, and therefore there exist strictly positive real numbers $\delta_1, \delta_2, \ldots, \delta_k$ such that

$$B_X(x,\delta_j) \subset V_j$$

for j = 1, 2, ..., k. Let δ be the minimum of $\delta_1, \delta_2, ..., \delta_k$. Then $\delta > 0$. (This is where we need the fact that we are dealing with a finite collection of open sets.)

12

Moreover

$B_X(x,\delta) \subset B_X(x,\delta_j) \subset V_j$

for j = 1, 2, ..., k, and thus $B_X(x, \delta) \subset V$. This shows that the intersection V of the open sets $V_1, V_2, ..., V_k$ is itself open. Thus (iii) is satisfied.

Any metric space may be regarded as a topological space. Indeed let X be a metric space with distance function d. We recall that a subset V of X is an *open set* if and only if, given any point v of V, there exists some $\delta > 0$ such that $\{x \in X : d(x,v) < \delta\} \subset V$. Proposition 1.2 shows that the topological space axioms are satisfied by the collection of open sets in any metric space. We refer to this collection of open sets as the *topology* generated by the distance function d on X.

Further Examples of Topological Spaces.

Example.

Given any set X, one can define a topology on X where every subset of X is an open set. This topology is referred to as the *discrete topology* on X.

Example.

Given any set X, one can define a topology on X in which the only open sets are the empty set \emptyset and the whole set X.

Closed Sets.

Definition.

Let X be a topological space. A subset F of X is said to be a *closed set* if and only if its complement $X \setminus F$ is an open set.

We recall that the complement of the union of some collection of subsets of some set X is the intersection of the complements of those sets, and the complement of the intersection of some collection of subsets of X is the union of the complements of those sets. The following result therefore follows directly from the definition of a topological space.

Proposition 1.3 Let X be a topological space. Then the collection of closed sets of X has the following properties:—

- (i) the empty set \emptyset and the whole set X are closed sets,
- (ii) the intersection of any collection of closed sets is itself a closed set,
- (iii) the union of any finite collection of closed sets is itself a closed set.

Hausdorff Spaces.

Definition.

A topological space X is said to be a *Haus*dorff space if and only if it satisfies the following *Hausdorff Axiom*:

• if x and y are distinct points of X then there exist open sets U and V such that $x \in U, y \in V$ and $U \cap V = \emptyset$.

Lemma 1.4 All metric spaces are Hausdorff spaces.

Proof.

Let X be a metric space with distance function d, and let x and y be points of X, where $x \neq y$. Let $\varepsilon = \frac{1}{2}d(x,y)$. Then the open balls $B_X(x,\varepsilon)$ and $B_X(y,\varepsilon)$ of radius ε centred on the points x and y are open sets (see Lemma 1.1). If $B_X(x,\varepsilon) \cap B_X(y,\varepsilon)$ were nonempty then there would exist $z \in X$ satisfying $d(x,z) < \varepsilon$ and $d(z,y) < \varepsilon$. But this is impossible, since it would then follow from the Triangle Inequality that $d(x,y) < 2\varepsilon$, contrary to the choice of ε . Thus $x \in B_X(x,\varepsilon)$, $y \in B_X(y,\varepsilon)$, $B_X(x,\varepsilon) \cap B_X(y,\varepsilon) = \emptyset$. This shows that the metric space X is a Hausdorff space.

We now give an example of a topological space which is not a Hausdorff space.

Example.

The Zariski topology on the set \mathbb{R} of real numbers is defined as follows: a subset U of \mathbb{R} is open (with respect to the Zariski topology) if and only if either $U = \emptyset$ or else $\mathbb{R} \setminus U$ is finite. It is a straightforward exercise to verify that the topological space axioms are satisfied, so that the set $\mathbb R$ of real numbers is a topological space with respect to this Zariski topology. Now the intersection of any two non-empty open sets in this topology is always non-empty. (Indeed if U and V are non-empty open sets then $U = \mathbb{R} \setminus F_1$ and $V = \mathbb{R} \setminus F_2$, where F_1 and F_2 are finite sets of real numbers. But then $U \cap V = \mathbb{R} \setminus (F_1 \cup F_2)$, which is non-empty, since $F_1 \cup F_2$ is finite and \mathbb{R} is infinite.) It follows immediately from this that \mathbb{R} , with the Zariski topology, is not a Hausdorff space.

Subspace Topologies.

Let X be a topological space with topology τ , and let A be a subset of X. Let τ_A be the collection of all subsets of A that are of the form $V \cap A$ for $V \in \tau$. Then τ_A is a topology on the set A. (It is a straightforward exercise to verify that the topological space axioms are satisfied.) The topology τ_A on A is referred to as the subspace topology on A.

Any subset of a Hausdorff space is itself a Hausdorff space (with respect to the subspace topology).

Lemma 1.5 Let X be a metric space with distance function d, and let A be a subset of X. A subset W of A is open with respect to the subspace topology on A if and only if, given any point w of W, there exists some $\delta > 0$ such that

 $\{a \in A : d(a, w) < \delta\} \subset W.$

Thus the subspace topology on A coincides with the topology on A obtained on regarding A as a metric space (with respect to the distance function d).

Proof.

Suppose that W is open with respect to the subspace topology on A. Then there exists some open set U in X such that $W = U \cap A$. Let w be a point of W. Then there exists some $\delta > 0$ such that

$$\{x \in X : d(x, w) < \delta\} \subset U.$$

But then

$$\{a \in A : d(a, w) < \delta\} \subset U \cap A = W.$$

Conversely, suppose that W is a subset of A with the property that, for any $w \in W$, there exists some $\delta_w > 0$ such that

$$\{a \in A : d(a, w) < \delta_w\} \subset W.$$

Define U to be the union of the open balls $B_X(w, \delta_w)$ as w ranges over all points of W, where

$$B_X(w,\delta_w) = \{x \in X : d(x,w) < \delta_w\}.$$

The set U is an open set in X, since each open ball $B_X(w, \delta_w)$ is an open set in X (see Lemma 1.1), and any union of open sets is itself an open set. Moreover

 $B_X(w,\delta_w) \cap A = \{a \in A : d(a,w) < \delta_w\} \subset W$

for any $w \in W$. Therefore $U \cap A \subset W$. However $W \subset U \cap A$, since, $W \subset A$ and $\{w\} \subset B_X(w, \delta_w) \subset U$ for any $w \in W$. Thus $W = U \cap A$, where U is an open set in X. We deduce that W is open with respect to the subspace topology on A.

Example.

Let X be any subset of n-dimensional Euclidean space \mathbb{R}^n . Then the subspace topology on X coincides with the topology on X generated by the Euclidean distance function on X. We refer to this topology as the usual topology on X.

Let X be a topological space, and let A be a subset of X. One can readily verify the following:—

- a subset B of A is closed in A (relative to the subspace topology on A) if and only if B = A ∩ F for some closed subset F of X;
- if A is itself open in X then a subset B of A is open in A if and only if it is open in X;
- if A is itself closed in X then a subset B of A is closed in A if and only if it is closed in X.

Continuous Functions between Topological Spaces.

Definition.

A function $f: X \to Y$ from a topological space X to a topological space Y is said to be *continuous* if $f^{-1}(V)$ is an open set in X for every open set V in Y, where

 $f^{-1}(V) \equiv \{x \in X : f(x) \in V\}.$

A continuous function from X to Y is often referred to as a *map* from X to Y.

Lemma 1.6 Let X, Y and Z be topological spaces, and let $f: X \to Y$ and $g: Y \to Z$ be continuous functions. Then the composition $g \circ f: X \to Z$ of the functions f and g is continuous.

Proof.

Let V be an open set in Z. Then $g^{-1}(V)$ is open in Y (since g is continuous), and hence $f^{-1}(g^{-1}(V))$ is open in X (since f is continuous). But $f^{-1}(g^{-1}(V)) = (g \circ f)^{-1}(V)$. Thus the composition function $g \circ f$ is continuous.

Lemma 1.7 Let X and Y be topological spaces, and let $f: X \to Y$ be a function from X to Y. The function f is continuous if and only if $f^{-1}(G)$ is closed in X for every closed subset G of Y.

Proof.

If G is any subset of Y then $X \setminus f^{-1}(G) = f^{-1}(Y \setminus G)$ (i.e., the complement of the preimage of G is the preimage of the complement of G). The result therefore follows immediately from the definitions of continuity and closed sets.

Continuous Functions between Metric Spaces.

The following definition of continuity for functions between metric spaces generalizes that for functions of a real or complex variable.

Definition.

Let X and Y be metric spaces with distance functions d_X and d_Y respectively. A function $f: X \to Y$ from X to Y is said to be *continuous* at a point x of X if and only if the following criterion is satisfied:—

• given any real number ε satisfying $\varepsilon > 0$ there exists some $\delta > 0$ such that $d_Y(f(x), f(x'))$ ε for all points x' of X satisfying $d_X(x, x') < \delta$.

The function $f: X \to Y$ is said to be continuous on X if and only if it is continuous at x for every point x of X.

This definition can be rephrased in terms of open balls: a function $f: X \to Y$ from a metric space X to a metric space Y is continuous at a point x of X if and only if, given any $\varepsilon > 0$, there exists some $\delta > 0$ such that f maps $B_X(x, \delta)$ into $B_Y(f(x), \varepsilon)$ (where $B_X(x, \delta)$ and $B_Y(f(x), \varepsilon)$ denote the open balls of radius δ and ε about x and f(x)respectively).

Let $f: X \to Y$ be a function from a set X to a set Y. Given any subset V of Y, we denote by $f^{-1}(V)$ the *preimage* of V under the map f, defined by

$$f^{-1}(V) = \{x \in X : f(x) \in V\}.$$

The following result shows that the definition of continuity given above for functions between metric spaces is consistent with the more general definition of continuity for functions between topological spaces. **Proposition 1.8** Let X and Y be metric spaces, and let $f: X \to Y$ be a function from X to Y. The function f is continuous if and only if $f^{-1}(V)$ is an open set in X for every open set V of Y.

Proof.

Suppose that $f: X \to Y$ is continuous. Let V be an open set in Y. We must show that $f^{-1}(V)$ is open in X. Let x be a point belonging to $f^{-1}(V)$. We must show that there exists some $\delta > 0$ with the property that $B_X(x,\delta) \subset f^{-1}(V)$. Now f(x) belongs to V. But V is open, hence there exists some $\varepsilon > 0$ with the property that $B_Y(f(x),\varepsilon) \subset V$. But f is continuous at x. Therefore there exists some $\delta > 0$ such that f maps the open ball $B_X(x,\delta)$ into $B_Y(f(x),\varepsilon)$ (see the remarks above). Thus $f(x') \in V$ for all $x' \in B_X(x, \delta)$, showing that $B_X(x,\delta) \subset f^{-1}(V)$. We have thus shown that if $f: X \to Y$ is continuous then $f^{-1}(V)$ is open in X for every open set V in Y.

Conversely suppose that $f: X \to Y$ has the property that $f^{-1}(V)$ is open in X for every open set V in Y. Let x be any point of X. We must show that f is continuous at x. Let $\varepsilon > 0$ be given. The open ball $B_Y(f(x), \varepsilon)$ is an open set in Y, by Lemma 1.1, hence $f^{-1}(B_Y(f(x), \varepsilon))$ is an open set in X which contains x. It follows that there exists some $\delta > 0$ such that $B_X(x, \delta) \subset f^{-1}(B_Y(f(x), \varepsilon))$. We have thus shown that, given any $\varepsilon > 0$, there exists some $\delta > 0$ such that f maps the open ball $B_X(x, \delta)$ into $B_Y(f(x), \varepsilon)$. We conclude that f is continuous at x, as required.

A Criterion for Continuity.

We now show that, if a topological space X is the union of a finite collection of closed sets, and if a function from X to some topological space is continuous on each of these closed sets, then that function is continuous on X. **Lemma 1.9** Let X and Y be topological spaces, let $f: X \to Y$ be a function from X to Y, and let $X = A_1 \cup A_2 \cup \cdots \cup A_k$, where A_1, A_2, \ldots, A_k are closed sets in X. Suppose that the restriction of f to the closed set A_i is continuous for $i = 1, 2, \ldots, k$. Then $f: X \to Y$ is continuous.

Proof.

Let V be an open set in Y. We must show that $f^{-1}(V)$ is open in X. Now the preimage of the open set V under the restriction $f|A_i$ of f to A_i is $f^{-1}(V) \cap A_i$. It follows from the continuity of $f|A_i$ that $f^{-1}(V) \cap A_i$ is relatively open in A_i for each i, and hence there exist open sets U_1, U_2, \ldots, U_k in X such that $f^{-1}(V) \cap A_i = U_i \cap A_i$ for $i = 1, 2, \ldots, k$. Let $W_i = U_i \cup (X \setminus A_i)$ for $i = 1, 2, \ldots, k$. Then W_i is an open set in X (as it is the union of the open sets U_i and $X \setminus A_i$), and $W_i \cap A_i = U_i \cap A_i = f^{-1}(V) \cap A_i$ for each i. We claim that

$$f^{-1}(V) = W_1 \cap W_2 \cap \cdots \cap W_k.$$

Let $W = W_1 \cap W_2 \cap \cdots \cap W_k$. Then $f^{-1}(V) \subset W_i$, since $f^{-1}(V) \subset W_i$ for each *i*. Also

$$W = \bigcup_{i=1}^{k} (W \cap A_i) \subset \bigcup_{i=1}^{k} (W_i \cap A_i)$$
$$= \bigcup_{i=1}^{k} (f^{-1}(V) \cap A_i) \subset f^{-1}(V),$$

since $X = A_1 \cup A_2 \cup \cdots \cup A_k$ and $W_i \cap A_i = f^{-1}(V) \cap A_i$ for each *i*. Therefore $f^{-1}(V) = W$. But *W* is open in *X*, since it is the intersection of a finite collection of open sets. We have thus shown that $f^{-1}(V)$ is open in *X* for any open set *V* in *Y*. Thus $f: X \to Y$ is continuous, as required.

Proof.

[Alternative Proof] A function $f: X \to Y$ is continuous if and only if $f^{-1}(G)$ is closed in Xfor every closed set G in Y (Lemma 1.7). Let G be an closed set in Y. Then $f^{-1}(G) \cap A_i$ is relatively closed in A_i for i = 1, 2, ..., k, since the restriction of f to A_i is continuous for each i. But A_i is closed in X, and therefore a subset of A_i is relatively closed in A_i if and only if it is closed in X. Therefore $f^{-1}(G) \cap$ A_i is closed in X for i = 1, 2, ..., k. Now $f^{-1}(G)$ is the union of the sets $f^{-1}(G) \cap A_i$ for i = 1, 2, ..., k. It follows that $f^{-1}(G)$, being a finite union of closed sets, is itself closed in X. It now follows from Lemma 1.7 that $f: X \to Y$ is continuous.

Example.

Let Y be a topological space, and let α : $[0, 1] \rightarrow Y$ and β : $[0, 1] \rightarrow Y$ be continuous functions defined on the interval [0, 1], where $\alpha(1) = \beta(0)$. Let γ : $[0, 1] \rightarrow Y$ be defined by

$$\gamma(t) = \begin{cases} \alpha(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \beta(2t-1) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Now $\gamma|[0,\frac{1}{2}] = \alpha \circ \rho$ where $\rho: [0,\frac{1}{2}] \rightarrow [0,1]$ is the continuous function defined by $\rho(t) = 2t$ for all $t \in [0,\frac{1}{2}]$. Thus $\gamma|[0,\frac{1}{2}]$ is continuous, being a composition of two continuous functions. Similarly $\gamma|[\frac{1}{2},1]$ is continuous. The subintervals $[0,\frac{1}{2}]$ and $[\frac{1}{2},1]$ are closed in [0,1], and [0,1] is the union of these two subintervals. It follows from Lemma 1.9 that $\gamma: [0,1] \rightarrow Y$ is continuous.

Homeomorphisms.

Definition.

Let X and Y be topological spaces. A function $h: X \to Y$ is said to be a *homeomorphism* if and only if the following conditions are satisfied:

- the function $h: X \to Y$ is both injective and surjective (so that the function $h: X \to Y$ has a well-defined inverse $h^{-1}: Y \to X$),
- the function $h: X \to Y$ and its inverse $h^{-1}: Y \to X$ are both continuous.

Two topological spaces X and Y are said to be *homeomorphic* if there exists a homeomorphism $h: X \to Y$ from X to Y. If $h: X \to Y$ is a homeomorphism between topological spaces X and Y then h induces a one-to-one correspondence between the open sets of X and the open sets of Y. Thus the topological spaces X and Y can be regarded as being identical as topological spaces.

Neighbourhoods, Closures and Interiors.

Definition.

Let X be a topological space, and let x be a point of X. Let N be a subset of X which contains the point x. Then N is said to be a *neighbourhood* of the point x if and only if there exists an open set U for which $x \in U$ and $U \subset N$.

One can readily verify that this definition of neighbourhoods in topological spaces is consistent with that for neighbourhoods in metric spaces. **Lemma 1.10** Let X be a topological space. A subset V of X is open in X if and only if V is a neighbourhood of each point belonging to V.

Proof.

It follows directly from the definition of neighbourhoods that an open set V is a neighbourhood of any point belonging to V. Conversely, suppose that V is a subset of X which is a neighbourhood of each $v \in V$. Then, given any point v of V, there exists an open set U_v such that $v \in U_v$ and $U_v \subset V$. Thus V is an open set, since it is the union of the open sets U_v as v ranges over all points of V.
Definition.

Let X be a topological space and let A be a subset of X. The closure \overline{A} of A in X is defined to be the intersection of all of the closed subsets of X that contain A. The interior A^0 of A in X is defined to be the union of all of the open subsets of X that are contained in A.

Let X be a topological space and let A be a subset of X. It follows directly from the definition of \overline{A} that the closure \overline{A} of A is uniquely characterized by the following two properties:

- (i) the closure \overline{A} of A is a closed set containing A,
- (ii) if F is any closed set containing A then F contains \overline{A} .

Similarly the interior A^0 of A is uniquely characterized by the following two properties:

- (i) the interior A^0 of A is an open set contained in A,
- (ii) if U is any open set contained in A then U is contained in A^0 .

Moreover a point x of A belongs to the interior A^0 of A if and only if A is a neighbourhood of x.

Connected Topological Spaces.

Definition.

A topological space X is said to be *connected* if the empty set \emptyset and the whole space X are the only subsets of X that are both open and closed.

Lemma 1.11 A topological space X is connected if and only if it has the following property: if U and V are non-empty open sets in X such that $X = U \cup V$, then $U \cap V$ is non-empty.

Proof.

If U is a subset of X that is both open and closed, and if $V = X \setminus U$, then U and V are both open, $U \cup V = X$ and $U \cap V = \emptyset$. Conversely if U and V are open subsets of X satisfying $U \cup V = X$ and $U \cap V = \emptyset$, then $U = X \setminus V$, and hence U is both open and closed. Thus a topological space X is connected if and only if there do not exist nonempty open sets U and V such that $U \cup V = X$ and $U \cap V = \emptyset$. The result follows.

Let \mathbb{Z} be the set of integers with the usual topology (i.e., the subspace topology on \mathbb{Z} induced by the usual topology on \mathbb{R}). Then $\{n\}$ is open for all $n \in \mathbb{Z}$, since

 $\{n\} = \mathbb{Z} \cap \{t \in \mathbb{R} : |t - n| < \frac{1}{2}\}.$

It follows that every subset of \mathbb{Z} is open (since it is a union of sets consisting of a single element, and any union of open sets is open). It follows that a function $f: X \to \mathbb{Z}$ on a topological space X is continuous if and only if $f^{-1}(V)$ is open in X for any subset V of \mathbb{Z} . We use this fact in the proof of the next theorem. **Proposition 1.12** A topological space X is connected if and only if every continuous function $f: X \to \mathbb{Z}$ from X to the set \mathbb{Z} of integers is constant.

Proof.

Suppose that X is connected. Let $f: X \to \mathbb{Z}$ be a continuous function. Choose $n \in f(X)$, and let

$$U = \{x \in X : f(x) = n\},\$$
$$V = \{x \in X : f(x) \neq n\}.$$

Then U and V are the preimages of the open subsets $\{n\}$ and $\mathbb{Z} \setminus \{n\}$ of \mathbb{Z} , and therefore both U and V are open in X. Moreover $U \cap$ $V = \emptyset$, and $X = U \cup V$. It follows that V = $X \setminus U$, and thus U is both open and closed. Moreover U is non-empty, since $n \in f(X)$. It follows from the connectedness of X that U = X, so that $f: X \to \mathbb{Z}$ is constant, with value n. Conversely suppose that every continuous function $f: X \to \mathbb{Z}$ is constant. Let S be a subset of X which is both open and closed. Let $f: X \to \mathbb{Z}$ be defined by

$$f(x) = \begin{cases} 1 & \text{if } x \in S; \\ 0 & \text{if } x \notin S. \end{cases}$$

Now the preimage of any subset of \mathbb{Z} under f is one of the open sets \emptyset , S, $X \setminus S$ and X. Therefore the function f is continuous. But then the function f is constant, so that either $S = \emptyset$ or S = X. This shows that X is connected. **Lemma 1.13** The closed interval [a, b] is connected, for all real numbers a and b satisfying $a \leq b$.

Proof.

Let $f:[a,b] \to \mathbb{Z}$ be a continuous integervalued function on [a,b]. We show that fis constant on [a,b]. Indeed suppose that f were not constant. Then $f(\tau) \neq f(a)$ for some $\tau \in [a,b]$. But the Intermediate Value Theorem would then ensure that, given any real number c between f(a) and $f(\tau)$, there would exist some $t \in [a,\tau]$ for which f(t) = c, and this is clearly impossible, since f is integer-valued. Thus f must be constant on [a,b]. We now deduce from Proposition 1.12 that [a,b] is connected.

Example.

Let $X = \{(x, y) \in \mathbb{R}^2 : x \neq 0\}$. The topological space X is not connected. Indeed if $f: X \to \mathbb{Z}$ is defined by

$$f(x,y) = \begin{cases} 1 & \text{if } x > 0, \\ -1 & \text{if } x < 0, \end{cases}$$

then f is continuous on X but is not constant.

A concept closely related to that of connectedness is *path-connectedness*. Let x_0 and x_1 be points in a topological space X. A *path* in X from x_0 to x_1 is defined to be a continuous function $\gamma: [0,1] \to X$ such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$. A topological space X is said to be *path-connected* if and only if, given any two points x_0 and x_1 of X, there exists a path in X from x_0 to x_1 .

Proposition 1.14 Every path-connected topological space is connected.

Proof.

Let *X* be a path-connected topological space, and let $f: X \to \mathbb{Z}$ be a continuous integervalued function on *X*. If x_0 and x_1 are any two points of *X* then there exists a path $\gamma: [0,1] \to X$ such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$. But then $f \circ \gamma: [0,1] \to \mathbb{Z}$ is a continuous integer-valued function on [0,1]. But [0,1] is connected (Lemma 1.13), therefore $f \circ \gamma$ is constant (Proposition 1.12). It follows that $f(x_0) = f(x_1)$. Thus every continuous integer-valued function on *X* is constant. Therefore *X* is connected, by Proposition 1.12. The topological spaces \mathbb{R} , \mathbb{C} and \mathbb{R}^n are all path-connected. Indeed, given any two points of one of these spaces, the straight line segment joining these two points is a continuous path from one point to the other. Also the *n*-sphere S^n is path-connected for all n > 0. We conclude that these topological spaces are connected.

Let A be a subset of a topological space X. Using Lemma 1.11 and the definition of the subspace topology, we see that A is connected if and only if the following condition is satisfied:

• if U and V are open sets in X such that $A \cap U$ and $A \cap V$ are non-empty and $A \subset U \cup V$ then $A \cap U \cap V$ is also non-empty.

Lemma 1.15 Let X be a topological space and let A be a connected subset of X. Then the closure \overline{A} of A is connected.

Proof.

It follows from the definition of the closure of A that $\overline{A} \subset F$ for any closed subset F of Xfor which $A \subset F$. On taking F to be the complement of some open set U, we deduce that $\overline{A} \cap U = \emptyset$ for any open set U for which $A \cap U = \emptyset$. Thus if U is an open set in X and if $\overline{A} \cap U$ is non-empty then $A \cap U$ must also be non-empty.

Now let U and V be open sets in X such that $\overline{A} \cap U$ and $\overline{A} \cap V$ are non-empty and $\overline{A} \subset U \cup V$. Then $A \cap U$ and $A \cap V$ are non-empty, and $A \subset U \cup V$. But A is connected. Therefore $A \cap U \cap V$ is non-empty, and thus $\overline{A} \cap U \cap V$ is non-empty. This shows that \overline{A} is connected. **Lemma 1.16** Let $f: X \to Y$ be a continuous function between topological spaces X and Y, and let A be a connected subset of X. Then f(A) is connected.

Proof.

Let $g: f(A) \to \mathbb{Z}$ be any continuous integervalued function on f(A). Then $g \circ f: A \to \mathbb{Z}$ is a continuous integer-valued function on A. It follows from Proposition 1.12 that $g \circ f$ is constant on A. Therefore g is constant on f(A). We deduce from Proposition 1.12 that f(A) is connected. **Proposition 1.17** Let X be a topological space. For each $x \in X$, let S_x be the union of all connected subsets of X that contain x. Then

(i) S_x is connected,

(ii) S_x is closed,

(iii) if $x, y \in X$, then either $S_x = S_y$, or else $S_x \cap S_y = \emptyset$.

Proof.

Let $f: S_x \to \mathbb{Z}$ be a continuous integer-valued function on S_x , for some $x \in X$. Let y be any point of S_x . Then, by definition of S_x , there exists some connected set A containing both x and y. But then f is constant on A, and thus f(x) = f(y). This shows that the function f is constant on S_x . We deduce that S_x is connected. This proves (i). Moreover the closure $\overline{S_x}$ is connected, by Lemma 1.15. Therefore $\overline{S_x} \subset S_x$. This shows that S_x is closed, proving (ii).

49

Finally, suppose that x and y are points of Xfor which $S_x \cap S_y \neq \emptyset$. Let $f: S_x \cup S_y \to \mathbb{Z}$ be any continuous integer-valued function on $S_x \cup S_y$. Then f is constant on both S_x and S_y . Moreover the value of f on S_x must agree with that on S_y , since $S_x \cap S_y$ is non-empty. We deduce that f is constant on $S_x \cup S_y$. Thus $S_x \cup S_y$ is a connected set containing both x and y, and thus $S_x \cup S_y \subset S_x$ and $S_x \cup S_y \subset S_y$, by definition of S_x and S_y . We conclude that $S_x = S_y$. This proves (iii).

Given any topological space X, the connected subsets S_x of X defined as in the statement of Proposition 1.17 are referred to as the *connected components* of X. We see from Proposition 1.17, part (iii) that the topological space X is the disjoint union of its connected components.

Example.

The connected components of $\{(x,y) \in \mathbb{R}^2 : x \neq 0\}$ are $\{(x,y) \in \mathbb{R}^2 : x > 0\}$

$$\{(x,y)\in\mathbb{R}^2:x<0\}.$$

Example.

The connected components of

 $\{t \in \mathbb{R} : |t - n| < \frac{1}{2} \text{ for some integer } n\}.$ are the sets J_n for all $n \in \mathbb{Z}$, where

$$J_n = (n - \frac{1}{2}, n + \frac{1}{2}).$$

Compact Topological Spaces.

Compact Topological Spaces.

Let X be a topological space, and let A be a subset of X. A collection of subsets of Xin X is said to *cover* A if and only if every point of A belongs to at least one of these subsets. In particular, an *open cover* of X is collection of open sets in X that covers X.

If \mathcal{U} and \mathcal{V} are open covers of some topological space X then \mathcal{V} is said to be a *subcover* of \mathcal{U} if and only if every open set belonging to \mathcal{V} also belongs to \mathcal{U} .

Definition.

A topological space X is said to be *compact* if and only if every open cover of X possesses a finite subcover.

Lemma 2.1 Let X be a topological space. A subset A of X is compact (with respect to the subspace topology on A) if and only if, given any collection \mathcal{U} of open sets in X covering A, there exists a finite collection V_1, V_2, \ldots, V_r of open sets belonging to \mathcal{U} such that $A \subset V_1 \cup V_2 \cup \cdots \cup V_r$.

Proof.

A subset *B* of *A* is open in *A* (with respect to the subspace topology on *A*) if and only if $B = A \cap V$ for some open set *V* in *X*. The desired result therefore follows directly from the definition of compactness.

We now show that any closed bounded interval in the real line is compact. This result is known as the *Heine-Borel Theorem*. The proof of this theorem uses the *least upper bound principle* which states that, given any non-empty set S of real numbers which is bounded above, there exists a *least upper bound* (or *supremum*) sup S for the set S. **Theorem 2.2** (Heine-Borel) Let a and b be real numbers satisfying a < b. Then the closed bounded interval [a,b] is a compact subset of \mathbb{R} .

Proof.

Let \mathcal{U} be a collection of open sets in \mathbb{R} with the property that each point of the interval [a,b] belongs to at least one of these open sets. We must show that [a,b] is covered by finitely many of these open sets.

Let *S* be the set of all $\tau \in [a, b]$ with the property that $[a, \tau]$ is covered by some finite collection of open sets belonging to \mathcal{U} , and let $s = \sup S$. Now $s \in W$ for some open set *W* belonging to \mathcal{U} . Moreover *W* is open in \mathbb{R} , and therefore there exists some $\delta > 0$ such that $(s - \delta, s + \delta) \subset W$. Moreover $s - \delta$ is not an upper bound for the set *S*, hence there exists some $\tau \in S$ satisfying $\tau > s - \delta$. It follows from the definition of *S* that $[a, \tau]$ is covered by some finite collection V_1, V_2, \ldots, V_r of open sets belonging to \mathcal{U} . Let $t \in [a, b]$ satisfy $\tau \leq t < s + \delta$. Then $[a, t] \subset [a, \tau] \cup (s - \delta, s + \delta) \subset V_1 \cup V_2 \cup \cdots \cup V_r \cup W$, and thus $t \in S$. In particular $s \in S$, and moreover s = b, since otherwise s would not be an upper bound of the set S. Thus $b \in S$, and therefore [a, b] is covered by a finite collection of open sets belonging to \mathcal{U} , as required.

Lemma 2.3 Let A be a closed subset of some compact topological space X. Then A is compact.

Proof.

Let \mathcal{U} be any collection of open sets in X covering A. On adjoining the open set $X \setminus A$ to \mathcal{U} , we obtain an open cover of X. This open cover of X possesses a finite subcover, since X is compact. Moreover A is covered by the open sets in the collection \mathcal{U} that belong to this finite subcover. It follows from Lemma 2.1 that A is compact, as required.

Lemma 2.4 Let $f: X \to Y$ be a continuous function between topological spaces X and Y, and let A be a compact subset of X. Then f(A) is a compact subset of Y.

Proof.

Let \mathcal{V} be a collection of open sets in Y which covers f(A). Then A is covered by the collection of all open sets of the form $f^{-1}(V)$ for some $V \in \mathcal{V}$. It follows from the compactness of A that there exists a finite collection V_1, V_2, \ldots, V_k of open sets belonging to \mathcal{V} such that

$$A \subset f^{-1}(V_1) \cup f^{-1}(V_2) \cup \cdots \cup f^{-1}(V_k).$$

But then $f(A) \subset V_1 \cup V_2 \cup \cdots \cup V_k$. This shows that f(A) is compact.

Lemma 2.5 Let $f: X \to \mathbb{R}$ be a continuous real-valued function on a compact topological space X. Then f is bounded above and below on X.

Proof.

The range f(X) of the function f is covered by some finite collection I_1, I_2, \ldots, I_k of open intervals of the form (-m, m), where $m \in \mathbb{N}$, since f(X) is compact (Lemma 2.4) and \mathbb{R} is covered by the collection of all intervals of this form. It follows that $f(X) \subset (-M, M)$, where (-M, M) is the largest of the intervals I_1, I_2, \ldots, I_k . Thus the function f is bounded above and below on X, as required. **Proposition 2.6** Let $f: X \to \mathbb{R}$ be a continuous real-valued function on a compact topological space X. Then there exist points u and v of X such that

 $f(u) \le f(x) \le f(v)$

for all $x \in X$.

Proof.

Let $m = \inf\{f(x) : x \in X\}$ and $M = \sup\{f(x) : x \in X\}$. There must exist $v \in X$ satisfying f(v) = M, for if f(x) < M for all $x \in X$ then the function

$$x \mapsto 1/(M - f(x))$$

would be a continuous real-valued function on X that was not bounded above, contradicting Lemma 2.5. Similarly there must exist $u \in X$ satisfying f(u) = m, since otherwise the function $x \mapsto 1/(f(x) - m)$ would be a continuous function on X that was not bounded above, again contradicting Lemma 2.5. But then $f(u) \leq f(x) \leq f(v)$ for all $x \in X$, as required.

Proposition 2.7 Let A be a compact subset of a metric space X. Then A is closed in X.

Proof.

Let p be a point of X that does not belong to A, and let f(x) = d(x, p), where d is the distance function on X. It follows from Proposition 2.6 that there is a point q of A such that $f(a) \ge f(q)$ for all $a \in A$, since A is compact. Now f(q) > 0, since $q \ne p$. Let δ satisfy $0 < \delta \le f(q)$. Then the open ball of radius δ about the point p is contained in the complement of A, since f(x) < f(q) for all points x of this open ball. It follows that the complement of A is an open set in X, and thus A itself is closed in X.

Proposition 2.8 Let X be a Hausdorff topological space, and let K be a compact subset of X. Let x be a point of $X \setminus K$. Then there exist open sets V and W in X such that $x \in V$, $K \subset W$ and $V \cap W = \emptyset$.

Proof.

For each point $y \in K$ there exist open sets $V_{x,y}$ and $W_{x,y}$ such that $x \in V_{x,y}$, $y \in W_{x,y}$ and $V_{x,y} \cap W_{x,y} = \emptyset$ (since X is a Hausdorff space). But then there exists a finite set $\{y_1, y_2, \ldots, y_r\}$ of points of K such that K is contained in $W_{x,y_1} \cup W_{x,y_2} \cup \cdots \cup W_{x,y_r}$, since K is compact. Define

$$V = V_{x,y_1} \cap V_{x,y_2} \cap \cdots \cap V_{x,y_r},$$

$$W = W_{x,y_1} \cup W_{x,y_2} \cup \cdots \cup W_{x,y_r}.$$

Then V and W are open sets, $x \in V$, $K \subset W$ and $V \cap W = \emptyset$, as required. **Corollary 2.9** A compact subset of a Hausdorff topological space is closed.

Proof.

Let K be a compact subset of a Hausdorff topological space X. It follows immediately from Proposition 2.8 that, for each $x \in X \setminus K$, there exists an open set V_x such that $x \in V_x$ and $V_x \cap K = \emptyset$. But then $X \setminus K$ is equal to the union of the open sets V_x as x ranges over all points of $X \setminus K$, and any set that is a union of open sets is itself an open set. We conclude that $X \setminus K$ is open, and thus K is closed.

Proposition 2.10 Let X be a Hausdorff topological space, and let K_1 and K_2 be compact subsets of X, where $K_1 \cap K_2 = \emptyset$. Then there exist open sets U_1 and U_2 such that $K_1 \subset U_1$, $K_2 \subset U_2$ and $U_1 \cap U_2 = \emptyset$.

Proof.

It follows from Proposition 2.8 that, for each point x of K_1 , there exist open sets V_x and W_x such that $x \in V_x$, $K_2 \subset W_x$ and $V_x \cap$ $W_x = \emptyset$. But then there exists a finite set $\{x_1, x_2, \ldots, x_r\}$ of points of K_1 such that

$$K_1 \subset V_{x_1} \cup V_{x_2} \cup \cdots \cup V_{x_r},$$

since K_1 is compact. Define

$$U_1 = V_{x_1} \cup V_{x_2} \cup \cdots \cup V_{x_r},$$

$$U_2 = W_{x_1} \cap W_{x_2} \cap \cdots \cap W_{x_r}.$$

Then U_1 and U_2 are open sets, $K_1 \subset U_1$, $K_2 \subset U_2$ and $U_1 \cap U_2 = \emptyset$, as required.

Lemma 2.11 Let $f: X \to Y$ be a continuous function from a compact topological space X to a Hausdorff space Y. Then f(K) is closed in Y for every closed set K in X.

Proof.

If K is a closed set in X, then K is compact (Lemma 2.3), and therefore f(K) is compact (Lemma 2.4). But any compact subset of a Hausdorff space is closed (Corollary 2.9). Thus f(K) is closed in Y, as required.

Remark.

If the Hausdorff space Y in Lemma 2.11 is a metric space, then Proposition 2.7 may be used in place of Corollary 2.9 in the proof of the lemma. **Theorem 2.12** A continuous bijection $f: X \rightarrow Y$ from a compact topological space X to a Hausdorff space Y is a homeomorphism.

Proof.

Let $g: Y \to X$ be the inverse of the bijection $f: X \to Y$. If U is open in X then $X \setminus U$ is closed in X, and hence $f(X \setminus U)$ is closed in Y, by Lemma 2.11. But

 $f(X \setminus U) = g^{-1}(X \setminus U) = Y \setminus g^{-1}(U).$

It follows that $g^{-1}(U)$ is open in Y for every open set U in X. Therefore $g: Y \to X$ is continuous, and thus $f: X \to Y$ is a homeomorphism.

We recall that a function $f: X \to Y$ from a topological space X to a topological space Y is said to be an *identification map* if it is surjective and satisfies the following condition: a subset U of Y is open in Y if and only if $f^{-1}(U)$ is open in X. **Proposition 2.13** A continuous surjection $f: X \rightarrow Y$ from a compact topological space X to a Hausdorff space Y is an identification map.

Proof.

Let U be a subset of Y. We claim that

 $Y \setminus U = f(K)$, where $K = X \setminus f^{-1}(U)$. Clearly $f(K) \subset Y \setminus U$. Also, given any $y \in Y \setminus U$, there exists $x \in X$ satisfying y = f(x), since $f: X \to Y$ is surjective. Moreover $x \in K$, since $f(x) \notin U$. Thus $Y \setminus U \subset f(K)$, and hence $Y \setminus U = f(K)$, as claimed.

We must show that the set U is open in Y if and only if $f^{-1}(U)$ is open in X. First suppose that $f^{-1}(U)$ is open in X. Then K is closed in X, and hence f(K) is closed in Y, by Lemma 2.11. It follows that U is open in Y. Conversely if U is open in Y then $f^{-1}(U)$ is open in X, since $f: X \to Y$ is continuous. Thus the surjection $f: X \to Y$ is an identification map.

Example.

Let S^1 be the unit circle in \mathbb{R}^2 , defined by

$$S^{1} = \{ (x, y) \in \mathbb{R}^{2} : x^{2} + y^{2} = 1 \},\$$

and let $q: [0,1] \to S^1$ be defined by

$$q(t) = (\cos 2\pi t, \sin 2\pi t)$$

for all $t \in [0, 1]$. It has been shown that the map q is an identification map. This also follows directly from the fact that $q: [0, 1] \rightarrow S^1$ is a continuous surjection from the compact space [0, 1] to the Hausdorff space S^1 .

The Lebesgue Lemma and Uniform Continuity.

Definition.

Let X be a metric space with distance function d. A subset A of X is said to be bounded if there exists a non-negative real number K such that $d(x,y) \leq K$ for all $x, y \in A$. The smallest real number K with this property is referred to as the diameter of A, and is denoted by diam A. (Note that diam A is the supremum of the values of d(x,y) as x and y range over all points of A.)

Lemma 2.14 (Lebesgue Lemma) Let (X, d)be a compact metric space. Let \mathcal{U} be an open cover of X. Then there exists a positive real number δ such that every subset of X whose diameter is less than δ is contained wholly within one of the open sets belonging to the open cover \mathcal{U} .

67

Proof.

Every point of X is contained in at least one of the open sets belonging to the open cover \mathcal{U} . It follows from this that, for each point x of X, there exists some $\delta_x > 0$ such that the open ball $B(x, 2\delta_x)$ of radius $2\delta_x$ about the point x is contained wholly within one of the open sets belonging to the open cover \mathcal{U} . But then the collection consisting of the open balls $B(x, \delta_x)$ of radius δ_x about the points x of X forms an open cover of the compact space X. Therefore there exists a finite set x_1, x_2, \ldots, x_r of points of X such that

 $B(x_1, \delta_1) \cup B(x_2, \delta_2) \cup \cdots \cup B(x_r, \delta_r) = X,$ where $\delta_i = \delta_{x_i}$ for $i = 1, 2, \ldots, r$. Let $\delta > 0$ be given by

$$\delta = \min(\delta_1, \delta_2, \dots, \delta_r).$$

Suppose that A is a subset of X whose diameter is less than δ . Let u be a point of A. Then u belongs to $B(x_i, \delta_i)$ for some integer i between 1 and r. But then it follows that $A \subset B(x_i, 2\delta_i)$, since, for each point v of A,

 $d(v, x_i) \leq d(v, u) + d(u, x_i) < \delta + \delta_i \leq 2\delta_i$. But $B(x_i, 2\delta_i)$ is contained wholly within one of the open sets belonging to the open cover \mathcal{U} . Thus A is contained wholly within one of the open sets belonging to \mathcal{U} , as required.

Let \mathcal{U} be an open cover of a compact metric space X. A Lebesgue number for the open cover \mathcal{U} is a positive real number δ such that every subset of X whose diameter is less than δ is contained wholly within one of the open sets belonging to the open cover \mathcal{U} . The Lebesgue Lemma thus states that there exists a Lebesgue number for every open cover of a compact metric space. Let X and Y be metric spaces with distance functions d_X and d_Y respectively, and let

$$f: X \to Y$$

be a function from X to Y. The function f is said to be *uniformly continuous* on X if and only if, given $\varepsilon > 0$, there exists some $\delta > 0$ such that

 $d_Y(f(x), f(x')) < \varepsilon$

for all points x and x' of X satisfying

$$d_X(x,x') < \delta.$$

(The value of δ should be independent of both x and x'.)

Theorem 2.15 Let X and Y be metric spaces. Suppose that X is compact. Then every continuous function from X to Y is uniformly continuous.

Proof.

Let d_X and d_Y denote the distance functions for the metric spaces X and Y respectively. Let $f: X \to Y$ be a continuous function from X to Y. We must show that f is uniformly continuous.

Let $\varepsilon > 0$ be given. For each $y \in Y$, define

$$V_y = \{ x \in X : d_Y(f(x), y) < \frac{1}{2}\varepsilon \}.$$

Note that $V_y = f^{-1} \left(B_Y(y, \frac{1}{2}\varepsilon) \right)$, where $B_Y(y, \frac{1}{2}\varepsilon)$ denotes the open ball of radius $\frac{1}{2}\varepsilon$ about yin Y. Now the open ball $B_Y(y, \frac{1}{2}\varepsilon)$ is an open set in Y, and f is continuous. Therefore V_y is open in X for all $y \in Y$. Note that $x \in V_{f(x)}$ for all $x \in X$. Now $\{V_y : y \in Y\}$ is an open cover of the compact metric space X. It follows from the Lebesgue Lemma (Lemma 2.14) that there exists some $\delta > 0$ such that every subset of X whose diameter is less than δ is a subset of some set V_y . Let x and x' be points of X satisfying $d_X(x, x') < \delta$. The diameter of the set $\{x, x'\}$ is $d_X(x, x')$, which is less than δ . Therefore there exists some $y \in Y$ such that $x \in V_y$ and $x' \in V_y$. But then

$$d_Y(f(x),y) < \frac{1}{2}\varepsilon$$

and

$$d_Y(f(x'),y) < \frac{1}{2}\varepsilon,$$

and hence

$$d_Y(f(x), f(x')) \leq d_Y(f(x), y) + d_Y(y, f(x')) < \varepsilon.$$

This shows that $f: X \to Y$ is uniformly continuous, as required.
Let K be a closed bounded subset of \mathbb{R}^n . It follows from Theorem 3.15 and Theorem 2.15 that any continuous function $f: K \to \mathbb{R}^k$ is uniformly continuous.

Product and Quotient Topologies.

Bases for Topologies.

Proposition 3.1 Let X be a set, let β be a collection of subsets of X, and let τ be the collection consisting of the empty set, together with all subsets of X that are unions of sets belonging to the collection β . Then τ is a topology on X if and only if the following conditions are satisfied:—

- (i) the set X is the union of the subsets belonging to the collection β ;
- (ii) given subsets $B_1, B_2 \in \beta$, and given any point p of $B_1 \cap B_2$, there exists some $B \in \beta$ such that $p \in B$ and $B \subset B_1 \cap B_2$.

Proof.

First suppose that τ is a topology on X. Then $X \in \tau$. But any subset of X that belongs to τ is a union of sets belonging to β . Therefore X is a union of subsets belonging to the collection β , and thus condition (i) is satisfied.

Moreover the intersection of any two open subsets of a topological space is required to be open. Thus if τ is a topology on X, and if $B_1, B_2 \in \beta$, then $B_1, B_2 \in \tau$ and therefore $B_1 \cap B_2 \in \tau$. It follows that $B_1 \cap B_2$ is a union of subsets of X that belong to β , and therefore, given any $p \in B_1 \cap B_2$, there exists $B \in \beta$ such that $p \in B$ and $B \subset B_1 \cap B_2$. Thus condition (ii) is satisfied. Conversely we must prove that if the collection β of subsets of a set X satisfies conditions (i) and (ii) then the collection τ of unions of sets belonging to β is a topology on X.

The empty set belongs to τ . Condition (i) ensures that the whole set X belongs to τ . It follows directly from the definition of τ that any union of sets belonging to τ is a union of sets belonging to β , and therefore itself belongs to τ .

It therefore only remains to show that the intersection of any finite collection of sets belonging to τ belongs to τ . It suffices to prove that the intersection of two sets belonging to τ belongs to τ .

Let $V_1, V_2 \in \tau$, and let $p \in V_1 \cap V_2$. Then V_1 and V_2 are union of sets belonging to β , and therefore there exist $B_1, B_2 \in \beta$ such that $p \in B_1$, $p \in B_2$, $B_1 \subset V_1$, and $B_2 \subset V_2$. Now condition (ii) ensures the existence of $B_p \in \beta$ such that $p \in B_p$ and $B_p \subset B_1 \cap B_2$. Then $B_p \subset B_1 \subset V_1$ and $B_p \subset B_2 \subset V_2$. We have thus shown that, given any point p of $V_1 \cap V_2$, there exists some member B_p of the collection β such that $p \in B_p$ and $B_p \subset V_1 \cap V_2$. It follows that $V_1 \cap V_2$ is the union of the sets B_p as p ranges over all points of the intersection $V_1 \cap V_2$, and therefore $V_1 \cap V_2 \in \tau$. We have thus proved that the intersection of two sets belonging to τ belongs to τ . It follows directly by induction on the number of sets involved that the intersection of any finite collection of sets belonging to τ must belong to τ . Thus τ is a topology on the set X, as required.

Definition.

Let X be a set. A collection β of subsets of X is said to be a *basis* for a topology on X if the following conditions are satisfied:—

- (i) the set X is the union of the subsets belonging to the collection β ;
- (ii) given subsets $B_1, B_2 \in \beta$, and given any point p of $B_1 \cap B_2$, there exists some $B \in \beta$ such that $p \in B$ and $B \subset B_1 \cap B_2$.

If β is a basis for a topology on X then the topology generated by β is the topology whose open sets are those subsets of X that are unions of sets belonging to the basis β . **Lemma 3.2** Let X be a set, and let β be a basis for a topology on X. A non-empty subset V is open in X with respect to the topology generated by β if and only if, given any point v of V, there exists $B \in \beta$ such that $v \in B$ and $B \subset V$.

Proof.

This result follows directly from the fact that the non-empty open sets in X are those subsets of X that are unions of sets belonging to the basis β .

Example.

Let X be a metric space. Then the collection of open balls in X (determined by the distance function d on X) is a basis for a topology on X. Indeed let x and y be points of X that are the centres of open balls $B_X(x,r)$ and $B_X(y,s)$ of radii r and s respectively, and let $z \in B_X(x,r) \cap B_X(y,s)$. Then d(z,x) < rand d(z,y) < s. Let δ be the minimum of r-d(z,x) and s-d(z,y). Then $\delta > 0$ and $z \in B_X(z, \delta)$, and it follows from the Triangle Inequality that $B_X(z,\delta) \subset B_X(x,r) \cap B_X(y,s)$. This ensures that the collection of open balls in this metric space is a basis for a topology on the metric space. The topology generated by this basis is the standard topology carried by the metric space when we consider metric spaces to be topological spaces.

The following result is an immediate consequence of the requirements for a collection of subsets of a given set to be a basis for a topology of that set. **Lemma 3.3** Let X be a topological space, and let β be a collection of subsets of X. Suppose that $X \in \beta$ and that the intersection of any finite collection of sets belonging to β itself belongs to β . Then β is a basis for a topology on X.

Subbases for Topologies.

Let X be a set, and let σ be a collection of subsets of X. Let β be the collection of subsets of X consisting of the set X itself together with all subsets of X that are finite intersections of sets belonging to the collection σ . It follows from an immediate application of Lemma 3.3 that β is a basis for a topology τ on X. We refer to τ as the topology on X generated by the collection σ of subsets of X, and we refer to the collection σ as a subbasis for the topology τ which it generates. **Lemma 3.4** Let σ be a collection of subsets of a set X which is a subbasis for a topology τ on X, and let $\tilde{\tau}$ be a topology on X. Suppose that $\sigma \subset \tilde{\tau}$. Then $\tau \subset \tilde{\tau}$. Thus the topology τ generated by the subbasis σ is the smallest topology on X that contains σ .

Proof.

The definition of topologies ensures that $X \in$ $\tilde{\tau}$. It also ensures that any finite intersection of subsets of X that belong to $\tilde{\tau}$ must itself belong to $\tilde{\tau}$. It follows that if $\sigma \subset \tilde{\tau}$ then $\beta \subset \tilde{\tau}$, where β is the collection of subsets of X consisting of the set X itself together with all subsets of X that are finite intersections of sets belonging to the collection σ . Now every non-empty subset of X that belongs to the topology τ is a union of subsets of X that belong to β . These subsets belong to $\tilde{\tau}$, and any union of subsets of X belonging to $\tilde{\tau}$ must itself belong to $\tilde{\tau}$. Therefore $\tau \subset \tilde{\tau}$. This shows that the topology τ generated by a subbasis σ is the smallest topology on X that contains the members of the subbasis σ .

It follows immediately from Lemma 3.4 that the topology τ on a set X generated by a subbasis σ is the intersection of all topologies $\tilde{\tau}$ on X for which $\sigma \subset \tilde{\tau}$.

Product Topologies.

The Cartesian product $X_1 \times X_2 \times \cdots \times X_n$ of sets X_1, X_2, \ldots, X_n is defined to be the set of all ordered *n*-tuples (x_1, x_2, \ldots, x_n) , where $x_i \in X_i$ for $i = 1, 2, \ldots, n$.

The sets \mathbb{R}^2 and \mathbb{R}^3 are the Cartesian products $\mathbb{R} \times \mathbb{R}$ and $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$ respectively. **Lemma 3.5** Let X_1, X_2, \ldots, X_n be topological spaces, let X be the Cartesian product $X_1 \times X_2 \times \cdots \times X_n$ of the sets X_1, X_2, \ldots, X_n , and let β be the collection consisting of all subsets of the Cartesian product X that are of the form $V_1 \times V_2 \times \cdots \times V_n$, where V_j is an open set in X_j for $j = 1, 2, \ldots, n$. Then β is a basis for a topology on X.

Proof.

The definition of topological spaces ensures that each of the sets X_1, X_2, \ldots, X_n is an open set in itself. It follows that $X \in \beta$.

Let U_1, U_2, \ldots, U_m be subsets of X, where $U_i \in \beta$ for $i = 1, 2, \ldots, m$. Then there exist subsets $V_{i,j}$ of X_j , where $V_{i,j}$ is an open set in X_j for $j = 1, 2, \ldots, n$ and $i = 1, 2, \ldots, n$, such that

 $U_i = V_{i,1} \times V_{i,2} \times \cdots \times V_{i,n} \quad \text{for } i = 1, 2, \dots, m.$ Then

$$\bigcap_{i=1}^{m} U_i = V_1 \times V_2 \times \cdots \times V_n$$

where

$$V_j = \bigcap_{i=1}^m V_{i,j}$$
 for $j = 1, 2, ..., n$.

Indeed let $(x_1, x_2, \ldots, x_n) \in X$, where $x_j \in X_j$ for $j = 1, 2, \ldots, n$. Then

$$(x_1, x_2, \dots, x_n) \in \bigcap_{i=1}^m U_i$$

$$\iff (x_1, x_2, \dots, x_n) \in U_i$$
for $i = 1, 2, \dots, m$

$$\iff x_j \in V_{i,j} \text{ for } i = 1, 2, \dots, m$$
and $j = 1, 2, \dots, n$

$$\iff x_j \in \bigcap_{i=1}^m V_{i,j} \text{ for } j = 1, 2, \dots, n$$

$$\iff x_j \in V_j \text{ for } j = 1, 2, \dots, n$$

$$\iff (x_1, x_2, \dots, x_n) \in V_1 \times V_2 \times \dots \times V_n.$$

But V_j is an open set in X_j for j = 1, 2, ..., n, since any finite intersection of open sets in a topological space must itself be an open set in that space, and therefore $\bigcap_{i=1}^{m} U_i \in \beta$. We have now shown that $X \in \beta$, and also that any finite intersection of subsets of X that belong to β must itself belong to β . It now follows from Lemma 3.3 that β is a basis for a topology on X.

Definition.

Let X_1, X_2, \ldots, X_n be topological spaces. The *product topology* on the Cartesian product $X_1 \times X_2 \times \cdots \times X_n$ is the topology on this Cartesian product of sets that is generated by the collection consisting of all subsets of the product set that are of the form

 $V_1 \times V_2 \times \cdots \times V_n$,

where V_j is an open set in X_j for j = 1, 2, ..., n.

Lemma 3.6 Let $X_1, X_2, ..., X_n$ be topological spaces. A non-empty subset U of the Cartesian product $X_1 \times X_2 \times \cdots \times X_n$ is open with respect to the product topology if and only if, given any point p of U, there exist open sets V_i in X_i for i = 1, 2, ..., n such that $p \in V_1 \times V_2 \times \cdots \times V_n$ and $V_1 \times V_2 \times \cdots \times V_n \subset U$.

Proof.

The product topology is the topology generated by the collection β consisting of those subsets of the Cartesian product that are of the form $V_1 \times V_2 \times \cdots \times V_n$, where V_j is an open set in X_j for $j = 1, 2, \ldots, n$. This collection of subsets is a basis for the product topology (Lemma 3.5). It follows that a nonempty subset of X is open with respect to the product topology if and only if it is a union of sets that belong to the basis β . The result follows directly (see Lemma 3.2). **Lemma 3.7** Let X_1, X_2, \ldots, X_n and Z be topological spaces. Then a function

 $f: X_1 \times X_2 \times \cdots \times X_n \to Z$

is continuous if and only if, given any point p of

 $X_1 \times X_2 \times \cdots \times X_n$,

and given any open set U in Z containing f(p), there exist open sets V_i in X_i for i = 1, 2, ..., n such that

$$p \in V_1 \times V_2 \cdots \times V_n$$

and

$$f(V_1 \times V_2 \times \cdots \times V_n) \subset U$$

Proof.

Let V_i be an open set in X_i for i = 1, 2, ..., n, and let U be an open set in Z. Then

$$V_1 \times V_2 \times \cdots \times V_n \subset f^{-1}(U)$$

if and only if

$$f(V_1 \times V_2 \times \cdots \times V_n) \subset U.$$

It follows that $f^{-1}(U)$ is open in the product topology on

$$X_1 \times X_2 \times \cdots \times X_n$$

if and only if, given any point p of

 $X_1 \times X_2 \times \cdots \times X_n$

satisfying $f(p) \in U$, there exist open sets V_i in X_i for i = 1, 2, ..., n such that

$$f(V_1 \times V_2 \times \cdots \times V_n) \subset U.$$

The required result now follows from the definition of continuity. Let X_1, X_2, \ldots, X_n be topological spaces, and let V_i be an open set in X_i for $i = 1, 2, \ldots, n$. It follows directly from the definition of the product topology that

 $V_1 \times V_2 \times \cdots \times V_n$

is open in $X_1 \times X_2 \times \cdots \times X_n$.

Theorem 3.8 Let $X = X_1 \times X_2 \times \cdots \times X_n$, where X_1, X_2, \ldots, X_n are topological spaces and X is given the product topology, and for each i, let $p_i: X \to X_i$ denote the projection function which sends $(x_1, x_2, \ldots, x_n) \in X$ to x_i . Then the functions p_1, p_2, \ldots, p_n are continuous. Moreover a function $f: Z \to X$ mapping a topological space Z into X is continuous if and only if

 $p_i \circ f: Z \to X_i$

is continuous for $i = 1, 2, \ldots, n$.

Proof.

Let V be an open set in X_i . Then $p_i^{-1}(V) = X_1 \times \cdots \times X_{i-1} \times V \times X_{i+1} \times \cdots \times X_n$, and therefore $p_i^{-1}(V)$ is open in X. Thus $p_i: X \to X_i$ is continuous for all i.

Let $f: Z \to X$ be continuous. Then, for each i, $p_i \circ f: Z \to X_i$ is a composition of continuous functions, and is thus itself continuous.

Conversely suppose that $f: Z \to X$ is a function with the property that $p_i \circ f$ is continuous for all *i*. Let *U* be an open set in *X*. We must show that $f^{-1}(U)$ is open in *Z*.

Let z be a point of $f^{-1}(U)$, and let

$$f(z) = (u_1, u_2, \ldots, u_n).$$

Now U is open in X, and therefore there exist open sets V_1, V_2, \ldots, V_n in X_1, X_2, \ldots, X_n respectively such that $u_i \in V_i$ for all i and

$$V_1 \times V_2 \times \cdots \times V_n \subset U.$$

Let

$$N_z = f_1^{-1}(V_1) \cap f_2^{-1}(V_2) \cap \dots \cap f_n^{-1}(V_n),$$

where $f_i = p_i \circ f$ for i = 1, 2, ..., n. Now $f_i^{-1}(V_i)$ is an open subset of Z for i = 1, 2, ..., n, since V_i is open in X_i and $f_i: Z \to X_i$ is continuous. Thus N_z , being a finite intersection of open sets, is itself open in Z. Moreover

$$f(N_z) \subset V_1 \times V_2 \times \cdots \times V_n \subset U,$$

so that $N_z \subset f^{-1}(U)$. It follows that $f^{-1}(U)$ is the union of the open sets N_z as z ranges over all points of $f^{-1}(U)$. Therefore $f^{-1}(U)$ is open in Z. This shows that $f: Z \to X$ is continuous, as required. **Proposition 3.9** The usual topology on \mathbb{R}^n coincides with the product topology on \mathbb{R}^n obtained on regarding \mathbb{R}^n as the Cartesian product $\mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ of n copies of the real line \mathbb{R} .

Proof.

We must show that a subset U of \mathbb{R}^n is open with respect to the usual topology if and only if it is open with respect to the product topology.

Let U be a subset of \mathbb{R}^n that is open with respect to the usual topology, and let $\mathbf{u} \in U$. Then there exists some $\delta > 0$ such that

$$B(\mathbf{u},\delta) \subset U,$$

where

$$B(\mathbf{u},\delta) = {\mathbf{x} \in \mathbb{R}^n : |\mathbf{x} - \mathbf{u}| < \delta}$$

Let I_1, I_2, \ldots, I_n be the open intervals in \mathbb{R} defined by

$$I_i = \{t \in \mathbb{R} : u_i - \frac{\delta}{\sqrt{n}} < t < u_i + \frac{\delta}{\sqrt{n}}\}$$

for i = 1, 2, ..., n. Then $I_1, I_2, ..., I_n$ are open sets in \mathbb{R} . Moreover

 $\{\mathbf{u}\} \subset I_1 \times I_2 \times \cdots \times I_n \subset B(\mathbf{u}, \delta) \subset U,$

since

$$|\mathbf{x} - \mathbf{u}|^2 = \sum_{i=1}^n (x_i - u_i)^2 < n \left(\frac{\delta}{\sqrt{n}}\right)^2 = \delta^2$$

for all

$$\mathbf{x} \in I_1 \times I_2 \times \cdots \times I_n.$$

This shows that any subset U of \mathbb{R}^n that is open with respect to the usual topology on \mathbb{R}^n is also open with respect to the product topology on \mathbb{R}^n . Conversely suppose that U is a subset of \mathbb{R}^n that is open with respect to the product topology on \mathbb{R}^n , and let $\mathbf{u} \in U$. Then there exist open sets V_1, V_2, \ldots, V_n in \mathbb{R} containing u_1, u_2, \ldots, u_n respectively such that

 $V_1 \times V_2 \times \cdots \times V_n \subset U.$

Now we can find $\delta_1, \delta_2, \ldots, \delta_n$ such that $\delta_i > 0$ and

$$(u_i - \delta_i, u_i + \delta_i) \subset V_i$$

for all *i*. Let $\delta > 0$ be the minimum of $\delta_1, \delta_2, \ldots, \delta_n$. Then

$$B(\mathbf{u}, \delta) \subset V_1 \times V_2 \times \cdots \vee V_n \subset U,$$

for if $\mathbf{x} \in B(\mathbf{u}, \delta)$ then

$$|x_i - u_i| < \delta_i$$
 for $i = 1, 2, ..., n$.

This shows that any subset U of \mathbb{R}^n that is open with respect to the product topology on \mathbb{R}^n is also open with respect to the usual topology on \mathbb{R}^n . The following result is now an immediate corollary of Proposition 3.9 and Theorem 3.8.

Corollary 3.10 Let X be a topological space and let $f: X \to \mathbb{R}^n$ be a function from X to \mathbb{R}^n . Let us write

 $f(x) = (f_1(x), f_2(x), \dots, f_n(x))$

for all $x \in X$, where the components

 f_1, f_2, \ldots, f_n

of f are functions from X to \mathbb{R} . The function f is continuous if and only if its components f_1, f_2, \ldots, f_n are all continuous.

Let $f: X \to \mathbb{R}$ and $g: X \to \mathbb{R}$ be continuous real-valued functions on some topological space X. We claim that f+g, f-g and f.gare continuous. Now it is a straightforward exercise to verify that the sum and product functions $s: \mathbb{R}^2 \to \mathbb{R}$ and $p: \mathbb{R}^2 \to \mathbb{R}$ defined by s(x,y) = x + y and p(x,y) = xy are continuous, and $f + g = s \circ h$ and $f.g = p \circ h$, where $h: X \to \mathbb{R}^2$ is defined by h(x) = (f(x), g(x)). Moreover it follows from Corollary 3.10 that the function h is continuous, and compositions of continuous functions are continuous. Therefore f + g and f.g are continuous, as claimed. Also -g is continuous, and f - g = f + (-g), and therefore f - g is continuous. If in addition the continuous function g is non-zero everywhere on X then 1/gis continuous (since 1/g is the composition of g with the reciprocal function $t \mapsto 1/t$), and therefore f/g is continuous. **Lemma 3.11** The Cartesian product $X_1 \times X_2 \times \ldots \times X_n$ of Hausdorff spaces X_1, X_2, \ldots, X_n is Hausdorff.

Proof.

Let $X = X_1 \times X_2 \times \ldots, X_n$, and let u and v be distinct points of X, where

$$u = (x_1, x_2, \dots, x_n)$$

and

$$v = (y_1, y_2, \ldots, y_n).$$

Then $x_i \neq y_i$ for some integer *i* between 1 and *n*. But then there exist open sets *U* and *V* in X_i such that $x_i \in U$, $y_i \in V$ and $U \cap V = \emptyset$ (since X_i is a Hausdorff space). Let $p_i: X \to X_i$ denote the projection function. Then $p_i^{-1}(U)$ and $p_i^{-1}(V)$ are open sets in *X*, since p_i is continuous. Moreover $u \in p_i^{-1}(U)$, $v \in p_i^{-1}(V)$, and $p_i^{-1}(U) \cap p_i^{-1}(V) = \emptyset$. Thus *X* is Hausdorff, as required.

Cartesian Products of Connected Topological Spaces.

Lemma 3.12 The Cartesian product $X \times Y$ of connected topological spaces X and Y is itself connected.

Proof.

Let $f: X \times Y \to \mathbb{Z}$ be a continuous integervalued function from $X \times Y$ to Z. Choose $x_0 \in X$ and $y_0 \in Y$. The function $x \mapsto f(x, y_0)$ is continuous on X, and is thus constant. Therefore $f(x, y_0) = f(x_0, y_0)$ for all $x \in X$. Now fix x. The function $y \mapsto f(x, y)$ is continuous on Y, and is thus constant. Therefore

$$f(x,y) = f(x,y_0) = f(x_0,y_0)$$

for all $x \in X$ and $y \in Y$. We deduce from Proposition 1.12 that $X \times Y$ is connected.

We deduce immediately that a finite Cartesian product of connected topological spaces is connected.

Finite Cartesian Products of Compact Topological Spaces.

We shall show that a finite Cartesian product of compact spaces is compact. To prove this, we apply the following result, known as the *Tube Lemma*. **Lemma 3.13** Let X and Y be topological spaces, let K be a compact subset of Y, and U be an open set in $X \times Y$. Let

$$V = \{x \in X : \{x\} \times K \subset U\}.$$

Then V is an open set in X.

Proof.

Let $x \in V$. For each $y \in K$ there exist open subsets D_y and E_y of X and Y respectively such that $(x, y) \in D_y \times E_y$ and $D_y \times E_y \subset U$. Now there exists a finite set $\{y_1, y_2, \ldots, y_k\}$ of points of K such that

$$K \subset E_{y_1} \cup E_{y_2} \cup \cdots \cup E_{y_k},$$

since K is compact. Set

$$N_x = D_{y_1} \cap D_{y_2} \cap \cdots \cap D_{y_k}.$$

Then N_x is an open set in X. Moreover

$$N_x \times K \subset \bigcup_{i=1}^k (N_x \times E_{y_i}) \subset \bigcup_{i=1}^k (D_{y_i} \times E_{y_i}) \subset U,$$

so that $N_x \subset V$. It follows that V is the union of the open sets N_x for all $x \in V$. Thus V is itself an open set in X, as required.

Theorem 3.14 A Cartesian product of a finite number of compact spaces is itself compact.

Proof.

It suffices to prove that the product of two compact topological spaces X and Y is compact, since the general result then follows easily by induction on the number of compact spaces in the product.

Let \mathcal{U} be an open cover of $X \times Y$. We must show that this open cover possesses a finite subcover. Let x be a point of X. The set $\{x\} \times Y$ is a compact subset of $X \times Y$, since it is the image of the compact space Y under the continuous map from Y to $X \times Y$ which sends $y \in Y$ to (x, y), and the image of any compact set under a continuous map is itself compact (Lemma 2.4). Therefore there exists a finite collection U_1, U_2, \ldots, U_r of open sets belonging to the open cover \mathcal{U} such that $\{x\} \times Y$ is contained in

 $U_1 \cup U_2 \cup \cdots \cup U_r.$

Let V_x denote the set of all points x' of X for which $\{x'\} \times Y$ is contained in

 $U_1 \cup U_2 \cup \cdots \cup U_r.$

Then $x \in V_x$, and Lemma 3.13 ensures That V_x is an open set in X. Note that $V_x \times Y$ is covered by finitely many of the open sets belonging to the open cover \mathcal{U} .

Now $\{V_x : x \in X\}$ is an open cover of the space X. It follows from the compactness of X that there exists a finite set $\{x_1, x_2, \ldots, x_r\}$ of points of X such that

$$X = V_{x_1} \cup V_{x_2} \cup \cdots \cup V_{x_r}$$

. Now $X \times Y$ is the union of the sets $V_{x_j} \times Y$ for j = 1, 2, ..., r, and each of these sets can be covered by a finite collection of open sets belonging to the open cover \mathcal{U} . On combining these finite collections, we obtain a finite collection of open sets belonging to \mathcal{U} which covers $X \times Y$. This shows that $X \times Y$ is compact. **Theorem 3.15** Let K be a subset of \mathbb{R}^n . Then K is compact if and only if K is both closed and bounded.

Proof.

Suppose that K is compact. Then K is closed, since \mathbb{R}^n is Hausdorff, and a compact subset of a Hausdorff space is closed (by Corollary 2.9). For each natural number m, let B_m be the open ball of radius m about the origin, given by

 $B_m = \{ \mathbf{x} \in \mathbb{R}^n : |\mathbf{x}| < m \}.$

Then $\{B_m : m \in \mathbb{N}\}$ is an open cover of \mathbb{R}^n . It follows from the compactness of K that there exist natural numbers m_1, m_2, \ldots, m_k such that $K \subset B_{m_1} \cup B_{m_2} \cup \cdots \cup B_{m_k}$. But then $K \subset B_M$, where M is the maximum of m_1, m_2, \ldots, m_k , and thus K is bounded. Conversely suppose that K is both closed and bounded. Then there exists some real number L such that K is contained within the closed cube C given by

$$C = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : \\ -L \le x_j \le L \text{ for } j = 1, 2, \dots, n\}.$$

Now the closed interval [-L, L] is compact by the Heine-Borel Theorem (Theorem 2.2), and *C* is the Cartesian product of *n* copies of the compact set [-L, L]. It follows from Theorem 3.14 that *C* is compact. But *K* is a closed subset of *C*, and a closed subset of a compact topological space is itself compact, by Lemma 2.3. Thus *K* is compact, as required.

Identification Maps and Quotient Topologies.

Definition.

Let X and Y be topological spaces and let $q: X \to Y$ be a function from X to Y. The function q is said to be an *identification map* if and only if the following conditions are satisfied:

- the function $q: X \to Y$ is surjective,
- a subset U of Y is open in Y if and only if $q^{-1}(U)$ is open in X.

It follows directly from the definition that any identification map is continuous. Moreover, in order to show that a continuous surjection $q: X \to Y$ is an identification map, it suffices to prove that if V is a subset of Y with the property that $q^{-1}(V)$ is open in X then V is open in Y.

Lemma 3.16 Let X be a topological space, let Y be a set, and let $q: X \to Y$ be a surjection. Then there is a unique topology on Yfor which the function $q: X \to Y$ is an identification map.
Proof.

Let τ be the collection consisting of all subsets U of Y for which $q^{-1}(U)$ is open in X. Now $q^{-1}(\emptyset) = \emptyset$, and $q^{-1}(Y) = X$, so that $\emptyset \in \tau$ and $Y \in \tau$. If $\{V_{\alpha} : \alpha \in A\}$ is any collection of subsets of Y indexed by a set A, then it is a straightforward exercise to verify that

$$\bigcup_{\alpha \in A} q^{-1}(V_{\alpha}) = q^{-1} \left(\bigcup_{\alpha \in A} V_{\alpha} \right),$$

$$\bigcap_{\alpha \in A} q^{-1}(V_{\alpha}) = q^{-1} \left(\bigcap_{\alpha \in A} V_{\alpha}\right)$$

(i.e., given any collection of subsets of Y, the union of the preimages of the sets is the preimage of the union of those sets, and the intersection of the preimages of the sets is the preimage of the intersection of those sets). It follows easily from this that unions and finite intersections of sets belonging to τ must themselves belong to τ . Thus τ is a topology on Y, and the function $q: X \to Y$ is an identification map with respect to the topology τ . Clearly τ is the unique topology on Y for which the function $q: X \to Y$ is an identification map.

109

Let X be a topological space, let Y be a set, and let $q: X \to Y$ be a surjection. The unique topology on Y for which the function q is an identification map is referred to as the quotient topology (or identification topology) on Y.

Lemma 3.17 Let X and Y be topological spaces and let $q: X \to Y$ be an identification map. Let Z be a topological space, and let $f: Y \to Z$ be a function from Y to Z. Then the function f is continuous if and only if the composition function $f \circ q: X \to Z$ is continuous.

Proof.

Suppose that f is continuous. Then the composition function $f \circ q$ is a composition of continuous functions and hence is itself continuous.

Conversely suppose that $f \circ q$ is continuous. Let U be an open set in Z. Then $q^{-1}(f^{-1}(U))$ is open in X (since $f \circ q$ is continuous), and hence $f^{-1}(U)$ is open in Y (since the function q is an identification map). Therefore the function f is continuous, as required.

Example.

Let S^1 be the unit circle in \mathbb{R}^2 , and let $q\colon [0,1]\to S^1$ be the map that sends $t\in [0,1]$ to

 $(\cos 2\pi t, \sin 2\pi t).$

Then $q: [0, 1] \to S^1$ is an identification map, and therefore a function $f: S^1 \to Z$ from S^1 to some topological space Z is continuous if and only if $f \circ q: [0, 1] \to Z$ is continuous.

Example.

Let S^n be the *n*-sphere, consisting of all points x in \mathbb{R}^{n+1} satisfying $|\mathbf{x}| = 1$. Let $\mathbb{R}P^n$ be the set of all lines in \mathbb{R}^{n+1} passing through the origin (i.e., $\mathbb{R}P^n$ is the set of all onedimensional vector subspaces of \mathbb{R}^{n+1}). Let $q: S^n \to \mathbb{R}P^n$ denote the function which sends a point x of S^n to the element of $\mathbb{R}P^n$ represented by the line in \mathbb{R}^{n+1} that passes through both \mathbf{x} and the origin. Note that each element of $\mathbb{R}P^n$ is the image (under q) of exactly two antipodal points x and -x of S^n . The function q induces a corresponding quotient topology on $\mathbb{R}P^n$ such that $q: S^n \to \mathbb{R}P^n$ is an identification map. The set $\mathbb{R}P^n$, with this topology, is referred to as real projective *n*-dimensional space. In particular $\mathbb{R}P^2$ is referred to as the real projective plane. It follows from Lemma 3.17 that a function $f: \mathbb{R}P^n \to Z$ from $\mathbb{R}P^n$ to any topological space Z is continuous if and only if the composition function $f \circ q: S^n \to Z$ is continuous.

Covering Maps and the Monodromy Theorem.

The Helicoidal Covering of the Punctured Plane.

Let the topological space \boldsymbol{X} be the punctured plane

$$X = \{ (x, y) \in \mathbb{R}^2 : (x, y) \neq (0, 0) \}$$

with the usual topology, and let

$$\tilde{X} = \{(x, y, z) \in \mathbb{R}^3 : x \sin 2\pi z = y \cos 2\pi z \\ \text{and } x \cos 2\pi z + y \sin 2\pi z > 0\}.$$

Lemma 4.1 Let

$$\tilde{X} = \{(x, y, z) \in \mathbb{R}^3 : x \sin 2\pi z = y \cos 2\pi z \\ and \ x \cos 2\pi z + y \sin 2\pi z > 0\}.$$

Then a point (x, y, z) of \mathbb{R}^3 belongs to the surface \tilde{X} if and only if there exists some positive real number ρ such that

 $x = \rho \cos 2\pi z$ and $y = \rho \sin 2\pi z$.

Proof.

Suppose that $(x, y, z) \in \tilde{X}$. Let

$$\rho = x \cos 2\pi z + y \sin 2\pi z.$$

Then $\rho > 0$. Now

 $x\sin 2\pi z = y\cos 2\pi z.$

It follows that

$$\rho \cos 2\pi z = x \cos^2 2\pi z + y \sin 2\pi z \cos 2\pi z$$

= $x \cos^2 2\pi z + x \sin^2 2\pi z$
= $x;$
$$\rho \sin 2\pi z = x \sin 2\pi z \cos 2\pi z + y \sin^2 2\pi z$$

= $y \cos^2 2\pi z + y \sin^2 2\pi z$
= $y.$

Thus if (x, y, z) is a point of the surface \tilde{X} then there exists some positive real number ρ such that

 $x = \rho \cos 2\pi z$ and $y = \rho \sin 2\pi z$.

Conversely suppose that (x, y, z) is a point of \mathbf{R}^3 for which there exists some $\rho > 0$ such that

 $x = \rho \cos 2\pi z$ and $y = \rho \sin 2\pi z$.

Then

$$x\sin 2\pi z = y\cos 2\pi z$$

and

 $x\cos 2\pi z + y\sin 2\pi z = \rho > 0,$

and therefore $(x,y,z)\in \widetilde{X}$, as required

Now, for each real number θ , the intersection of the surface \tilde{X} with the plane $z = \frac{\theta}{2\pi}$ is the half line consisting of all points of \mathbb{R}^3 that are of the form

$$\left(\rho\cos\theta,\rho\sin\theta,\frac{\theta}{2\pi}\right)$$

for some $\rho > 0$. Moreover both the height $\frac{\theta}{2\pi}$ of this half line and the angle which the halfline makes with the *x*-direction increase at a constant rate as θ increases. The surface \tilde{X} thus takes the form of a spiral ramp about the *z*-axis in \mathbb{R}^3 . The surface \tilde{X} is an example of a *helicoid*.

We let $p: \tilde{X} \to X$ be the continuous map defined such that

$$p(x, y, z) = (x, y)$$

for all $(x, y, z) \in \tilde{X}$. The map p is a surjective map from the helicoid \tilde{X} to the punctured plane X.

Given any real number θ , let

$$\widetilde{U}_{\theta} = \left\{ (x, y, z) \in \widetilde{X} : \left| z - \frac{\theta}{2\pi} \right| < \frac{1}{2} \right\},$$

and let $U_{\theta} = p(\tilde{U}_{\theta})$. Then U_{θ} is the sector of the punctured plane consisting all all halflines starting at the origin that make an angle of less than π with the half-line in the direction of the vector ($\cos \theta$, $\sin \theta$). It follows that $U_{\theta} = X \setminus L_{\theta}$, where L_{θ} is the half-line from the origin in the direction of the vector $(-\cos \theta, -\sin \theta)$, defined so that

 $L_{\theta} = \{(-t\cos\theta, -t\sin\cos\theta) : t \in \mathbb{R} \text{ and } t > 0\}.$

Then the preimage $p^{-1}(U_{\theta})$ of U_{θ} is the disjoint union $\bigcup_{n \in \mathbb{Z}} V_n$ of the open subsets V_n of \tilde{X} for all integers n, where

$$V_n = \{(x, y, z) \in \tilde{X} : (x, y, z - n) \in \tilde{U}_{\theta}\}$$

= $\{(x, y, z) \in \tilde{X} : \frac{\theta}{2\pi} + n - \frac{1}{2} < z$
 $< \frac{\theta}{2\pi} + n + \frac{1}{2}\}.$

Each of these open set V_n is mapped homeomorphically onto U_{θ} by the map $p: \tilde{X} \to X$. Indeed let $s_n: \tilde{U}_{\theta} \to V_n$ be defined such that

$$s_n(\rho\cos(\theta+\varphi),\rho\sin(\theta+\varphi)) = \left(\rho\cos(\theta+\varphi),\rho\sin(\theta+\varphi),\frac{\theta+\varphi}{2\pi}\right)$$

for all angles φ satisfying $-\pi < \varphi < \pi$. Then $s_n: U_\theta \to V_n$ is a continuous map, and this map is the inverse of the restriction of the map $p: \tilde{X} \to X$ to V_n .

It follows that the preimage $p^{-1}(U_{\theta})$ of the open subset U_{θ} of X is a disjoint union of open sets, each of which is mapped homeomorphically onto U_{θ} by the map $p: \tilde{X} \to X$. We say that the open set U_{θ} is *evenly covered* by the continuous map $p: \tilde{X} \to X$.

A continuous map between topological spaces is said to be a *covering map* if it is surjective, and if in addition every point of the codomain is contained in some evenly covered open set. The map $p: \tilde{X} \to X$ from the helicoid \tilde{X} to the punctured plane X is an example of a covering map.

A path $\gamma: [0, 1] \to X$ defined on the unit interval [0, 1] is by definition a continuous map from [0,1] into the topological space X. It can be shown that, given any path $\gamma: [0, 1] \rightarrow$ X, and given any point w of \tilde{X} satisfying $p(w) = \gamma(0)$, there exists a unique path $\tilde{\gamma}: [0, 1] \rightarrow 0$ \tilde{X} such that $\tilde{\gamma}(0) = w$ and $p \circ \tilde{\gamma} = \gamma$. Such a path $\tilde{\gamma}: [0,1] \to \tilde{X}$ is said to be a *lift* of the path $\gamma: [0, 1] \to X$. This result is a special case of the Path Lifting Theorem for covering maps. It can also be shown that, given any continuous map $H: [0, 1] \times [0, 1] \rightarrow X$ defined on the unit square $[0,1] \times [0,1]$, and given any point w of \tilde{X} satisfying p(w) =H(0,0), there exists a unique continuous map $\tilde{H}\colon [0,1]\times [0,1]\to \tilde{X}$ such that $\tilde{H}(0,0)=w$ and $p \circ \tilde{H} = H$. This result is a special case of the Monodromy Theorem, a theorem applicable to any covering map.

Covering Maps.

Definition.

Let X and \tilde{X} be topological spaces and let $p: \tilde{X} \to X$ be a continuous map. An open subset U of X is said to be *evenly covered* by the map p if and only if $p^{-1}(U)$ is a disjoint union of open sets of \tilde{X} each of which is mapped homeomorphically onto U by p. The map $p: \tilde{X} \to X$ is said to be a *covering map* if $p: \tilde{X} \to X$ is surjective and in addition every point of X is contained in some open set that is evenly covered by the map p.

If $p: \tilde{X} \to X$ is a covering map, then we say that \tilde{X} is a *covering space* of X.

Example.

Let S^1 be the unit circle in \mathbb{R}^2 . Then the map $p: \mathbb{R} \to S^1$ defined by

 $p(t) = (\cos 2\pi t, \sin 2\pi t)$

is a covering map. Indeed let n be a point of S^1 . Consider the open set U in S^1 containing n defined by $U = S^1 \setminus \{-n\}$. Now n = $(\cos 2\pi t_0, \sin 2\pi t_0)$ for some $t_0 \in \mathbb{R}$. Then $p^{-1}(U)$ is the union of the disjoint open sets J_n for all integers n, where

$$J_n = \{ t \in \mathbb{R} : t_0 + n - \frac{1}{2} < t < t_0 + n + \frac{1}{2} \}.$$

Each of the open sets J_n is mapped homeomorphically onto U by the map p. This shows that $p: \mathbb{R} \to S^1$ is a covering map.

Example.

The map

$$p_{\mathsf{exp}}: \mathbb{C} \to \mathbb{C} \setminus \{\mathbf{0}\}$$

defined by $p_{\exp}(z) = \exp(z)$ is a covering map, where

$$\exp(x+iy) = e^x \cos y + ie^x \sin y$$

for all real numbers x and y. This covering map is closely related to the helicoidal covering $p_{\text{vert}}: \tilde{X} \to X$ of the punctured plane X, where

$$X = \{(x, y) \in \mathbb{R}^2 : (x, y) \neq (0, 0)\},\$$

$$\tilde{X} = \{(x, y, z) \in \mathbb{R}^3 : x \sin 2\pi z = y \cos 2\pi z \\$$

and $x \cos 2\pi z + y \sin 2\pi z > 0\}.$

and $p_{\text{vert}}(x, y, z) = (x, y)$ for all $(x, y) \in \tilde{X}$.

Indeed let $h: \mathbb{C} \to \tilde{X}$ be defined such that

$$h(u+iv) = \left(e^u \cos v, e^u \sin v, \frac{v}{2\pi}\right)$$

for all $u, v \in \mathbb{R}$. Then the function h is a homeomorphism from the complex plane \mathbb{C} to the helicoid \tilde{X} whose inverse $h^{-1}\tilde{X} \to \mathbb{C}$ satisfies

$$h^{-1}(x, y, z) = \left(\frac{1}{2}\log(x^2 + y^2), 2\pi z\right)$$

for all $(x, y, z) \in \tilde{X}$. Also let $\nu: X \to \mathbb{C} \setminus \{0\}$ be the natural homeomorphism between the punctured Euclidean plane $\mathbb{R}^2 \setminus \{(0,0)\}$ and the punctured complex plane $\mathbb{C} \setminus \{0\}$, defined such that $\nu(x, y) = x + iy$ for all $(x, y) \in X$. Then $p_{exp}(z) = \nu(p_{vert}(h(z)))$ for all $z \in \mathbb{C}$, and thus $p_{exp} = \nu \circ p_{vert} \circ h$. Moreover $h: \mathbb{C} \to \tilde{X}$ and $\nu: X \to \mathbb{C} \setminus \{0\}$ are homeomorphisms. It follows that $p_{exp}: \mathbb{C} \to \mathbb{C} \setminus \{0\}$ is a covering map that corresponds naturally to the helicoidal covering of the punctured plane. Given any $\theta \in [-\pi,\pi]$ let us define

 $W_{\theta} = \{z \in \mathbb{C} \setminus \{0\} : \arg(-z) \neq \theta\}.$

Then $p_{\exp}^{-1}(W_{\theta})$ is the disjoint union of the open sets

$$\left\{z\in\mathbb{C}:\left|\operatorname{Im}z-\theta-2\pi n\right|<\pi\right\},$$

for all integers n, and p_{exp} maps each of these open sets homeomorphically onto W_{θ} . Thus W_{θ} is evenly covered by the exponential map p_{exp} .

Example.

Consider the map $\alpha: (-2,2) \rightarrow S^1$, where $\alpha(t) = (\cos 2\pi t, \sin 2\pi t)$ for all $t \in (-2,2)$. It can easily be shown that there is no open set U containing the point (1,0) that is evenly covered by the map α . Indeed suppose that there were to exist such an open set U. Then there would exist some δ satisfying $0 < \delta < \frac{1}{2}$ such that $U_{\delta} \subset U$, where

 $U_{\delta} = \{(\cos 2\pi t, \sin 2\pi t) : -\delta < t < \delta\}.$

The open set U_{δ} would then be evenly covered by the map α . However the connected components of $\alpha^{-1}(U_{\delta})$ are

$$(-2, -2 + \delta), (-1 - \delta, -1 + \delta), (-\delta, \delta),$$

 $(1-\delta,1+\delta)$ and $(2-\delta,2)$,

and neither $(-2, -2+\delta)$ nor $(2-\delta, 2)$ is mapped homeomorphically onto U_{δ} by α .

Example.

Let $Z = \mathbb{C} \setminus \{1, -1\}$, let

 $\tilde{Z} = \{(z, w) \in \mathbb{C}^2 : w \neq 0 \text{ and } w^2 = z^2 - 1\},\$ and let $p: \tilde{Z} \to Z$ be defined such that

$$p(z,w) = z$$

for all $(z, w) \in \tilde{Z}$. Let $(z_0, w_0) \in \tilde{Z}$, let $z = z_0 + \zeta$, and let $(z, w) \in \mathbb{C}$ be a complex number for which $(z, w) \in \tilde{Z}$. Then

$$w^{2} = z^{2} - 1 = z_{0}^{2} - 1 + 2z_{0}\zeta + \zeta^{2}$$

= $w_{0}^{2} + 2z_{0}\zeta + \zeta^{2}$
= $w_{0}^{2} \left(1 + \frac{2z_{0}\zeta + \zeta^{2}}{w_{0}^{2}}\right).$

Now there exists $\delta > 0$ such that

$$\left|\frac{2z_0\zeta+\zeta^2}{w_0^2}\right|<1$$

whenever $|\zeta| < \delta$.

127

It follows from Taylor's Theorem (for holomorphic functions of a complex variable) that if $|z - z_0| < \delta$ and $(z, w) \in \tilde{W}$, then

$$w = \pm w_0 \sum_{n=0}^{+\infty} a_n \left(\frac{2z_0 \zeta + \zeta^2}{w_0^2} \right)^n,$$

where $a_0 = 1$ and

$$a_n = \frac{(\frac{3}{2} - n)a_{n-1}}{n}$$

for all natural numbers n satisfying n > 1. This ensures that the open disk $\{z \in Z : |z - z_0| < \delta\}$ is evenly covered by the map $p: \tilde{Z} \rightarrow Z$. We conclude from this that every point z_0 of Z is contained in some open set that is evenly covered by the map $p: \tilde{Z} \rightarrow Z$. This map is thus a covering map. Let $\tilde{f}(z,w) = w$ for all $(z,w) \in \tilde{Z}$. Then

$$\tilde{f}(\tilde{z})^2 = p(\tilde{z})^2 - 1$$

for all $\tilde{z} \in \tilde{Z}$. It follows that the function $\tilde{f}: \tilde{Z} \to \mathbb{C}$ represents in some sense the manyvalued 'function' $\sqrt{z^2 - 1}$. However this function \tilde{z} is not defined on the open subset Z of the complex plane, but is instead defined over a covering space \tilde{Z} of this open set. This covering space is the *Riemann surface* for the 'function' $\sqrt{z^2 - 1}$. This method of representing many-valued 'functions' of a complex variable using single-valued functions defined over a covering space was initiated and extensively developed by Bernhard Riemann (1826–1866) in his doctoral thesis. **Lemma 4.2** Let $p: \tilde{X} \to X$ be a covering map. Then p(V) is open in X for every open set V in \tilde{X} . In particular, a covering map $p: \tilde{X} \to X$ is a homeomorphism if and only if it is a bijection.

Proof.

Let V be open in \tilde{X} , and let $x \in p(V)$. Then x = p(v) for some $v \in V$. Now there exists an open set U containing the point x which is evenly covered by the covering map p. Then $p^{-1}(U)$ is a disjoint union of open sets, each of which is mapped homeomorphically onto U by the covering map p. One of these open sets contains v; let \tilde{U} be this open set, and let $N_x = p(V \cap \tilde{U})$. Now N_x is open in X, since $V \cap \tilde{U}$ is open in \tilde{U} and $p|\tilde{U}$ is a homeomorphism from \tilde{U} to U. Also $x \in N_x$ and $N_x \subset p(V)$.

It follows that p(V) is the union of the open sets N_x as x ranges over all points of p(V), and thus p(V) is itself an open set, as required. The result that a bijective covering map is a homeomorphism then follows directly from the fact that a continuous bijection is a homeomorphism if and only if it maps open sets to open sets.

Path Lifting and the Monodromy Theorem.

Let $p: \tilde{X} \to X$ be a covering map over a topological space X. Let Z be a topological space, and let $f: Z \to X$ be a continuous map from Z to X. A continuous map $\tilde{f}: Z \to \tilde{X}$ is said to be a *lift* of the map $f: Z \to X$ if and only if $p \circ \tilde{f} = f$. We shall prove various results concerning the existence and uniqueness of such lifts.

Proposition 4.3 Let $p: \tilde{X} \to X$ be a covering map, let Z be a connected topological space, and let $g: Z \to \tilde{X}$ and $h: Z \to \tilde{X}$ be continuous maps. Suppose that $p \circ g = p \circ h$ and that g(z) = h(z) for some $z \in Z$. Then g = h.

Proof.

Let $Z_0 = \{z \in Z : g(z) = h(z)\}$. Note that Z_0 is non-empty, by hypothesis. We show that Z_0 is both open and closed in Z.

Let z be a point of Z. There exists an open set U in X containing the point p(g(z)) which is evenly covered by the covering map p. Then $p^{-1}(U)$ is a disjoint union of open sets, each of which is mapped homeomorphically onto U by the covering map p. One of these open sets contains g(z); let this set be denoted by \tilde{U} . Also one of these open sets contains h(z); let this open set be denoted by \tilde{V} . Let $N_z = g^{-1}(\tilde{U}) \cap h^{-1}(\tilde{V})$. Then N_z is an open set in Z containing z.

Consider the case when $z \in Z_0$. Then g(z) = h(z), and therefore $\tilde{V} = \tilde{U}$. It follows from this that both g and h map the open set N_z into \tilde{U} . But $p \circ g = p \circ h$, and $p|\tilde{U}:\tilde{U} \to U$ is a homeomorphism. Therefore $g|N_z = h|N_z$, and thus $N_z \subset Z_0$. We have thus shown that, for each $z \in Z_0$, there exists an open set N_z such that $z \in N_z$ and $N_z \subset Z_0$. We conclude that Z_0 is open.

Next consider the case when $z \in Z \setminus Z_0$. In this case $\tilde{U} \cap \tilde{V} = \emptyset$, since $g(z) \neq h(z)$. But $g(N_z) \subset \tilde{U}$ and $h(N_z) \subset \tilde{V}$. Therefore $g(z') \neq$ h(z') for all $z' \in N_z$, and thus $N_z \subset Z \setminus Z_0$. We have thus shown that, for each $z \in Z \setminus Z_0$, there exists an open set N_z such that $z \in N_z$ and $N_z \subset Z \setminus Z_0$. We conclude that $Z \setminus Z_0$ is open.

The subset Z_0 of Z is therefore both open and closed. Also Z_0 is non-empty by hypothesis. We deduce that $Z_0 = Z$, since Z is connected. Thus g = h, as required.

Lemma 4.4 Let $p: \tilde{X} \to X$ be a covering map, let Z be a topological space, let A be a connected subset of Z, and let $f: Z \to X$ and $g: A \to \tilde{X}$ be continuous maps with the property that $p \circ g = f | A$. Suppose that $f(Z) \subset U$, where U is an open subset of X that is evenly covered by the covering map p. Then there exists a continuous map $\tilde{f}: Z \to \tilde{X}$ such that $\tilde{f} | A = g$ and $p \circ \tilde{f} = f$.

134

Proof.

The open set U is evenly covered by the covering map p, and therefore $p^{-1}(U)$ is a disjoint union of open sets, each of which is mapped homeomorphically onto U by the covering map p. One of these open sets contains g(a) for some $a \in A$; let this set be denoted by \tilde{U} . Let $\sigma: U \to \tilde{U}$ be the inverse of the homeomorphism $p|\tilde{U}:\tilde{U} \to U$, and let $\tilde{f} = \sigma \circ f$. Then $p \circ \tilde{f} = f$. Also $p \circ \tilde{f}|A = p \circ g$ and $\tilde{f}(a) = g(a)$. It follows from Proposition 4.3 that $\tilde{f}|A = g$, since A is connected. Thus $\tilde{f}: Z \to \tilde{X}$ is the required map.

Theorem 4.5 (Path Lifting Theorem) Let $p: \tilde{X} \to X$ be a covering map, let $\gamma: [0, 1] \to X$ be a continuous path in X, and let w be a point of \tilde{X} satisfying $p(w) = \gamma(0)$. Then there exists a unique continuous path

$$\tilde{\gamma}$$
: [0, 1] $\rightarrow \tilde{X}$

such that $\tilde{\gamma}(0) = w$ and $p \circ \tilde{\gamma} = \gamma$.

Proof.

The map $p: \tilde{X} \to X$ is a covering map; therefore there exists an open cover \mathcal{U} of X such that each open set U belonging to X is evenly covered by the map p. Now the collection consisting of the preimages $\gamma^{-1}(U)$ of the open sets U belonging to \mathcal{U} is an open cover of the interval [0, 1]. But [0, 1] is compact, by the Heine-Borel Theorem. It follows from the Lebesgue Lemma that there exists some $\delta > 0$ such that every subinterval of length less than δ is mapped by γ into one of the open sets belonging to \mathcal{U} . Partition the interval [0, 1] into subintervals $[t_{i-1}, t_i]$, where

$$0 = t_0 < t_1 < \dots < t_{n-1} < t_n = 1,$$

and where the length of each subinterval is less than δ . Then each subinterval $[t_{i-1}, t_i]$ is mapped by γ into some open set in X that is evenly covered by the map p.

It follows from Lemma 4.4 that once $\tilde{\gamma}(t_{i-1})$ has been determined, we can extend $\tilde{\gamma}$ continuously over the *i*th subinterval $[t_{i-1}, t_i]$. Thus by extending $\tilde{\gamma}$ successively over

$$[t_0, t_1], [t_1, t_2], \ldots, [t_{n-1}, t_n],$$

we can lift the path $\gamma: [0,1] \to X$ to a path $\tilde{\gamma}: [0,1] \to \tilde{X}$ starting at w. The uniqueness of $\tilde{\gamma}$ follows from Proposition 4.3.

Theorem 4.6 (The Monodromy Theorem) Let $p: \tilde{X} \to X$ be a covering map, let

 $H: [0,1] \times [0,1] \to X$

be a continuous map, and let w be a point of \tilde{X} satisfying p(w) = H(0,0). Then there exists a unique continuous map

 $\tilde{H} \colon [0,1] \times [0,1] \to \tilde{X}$

such that $\tilde{H}(0,0) = w$ and $p \circ \tilde{H} = H$.

Proof.

The unit square $[0,1] \times [0,1]$ is compact. By applying the Lebesgue Lemma to an open cover of the square by preimages of evenly covered open sets in X (as in the proof of Theorem 4.5), we see that there exists some $\delta > 0$ with the property that any square contained in $[0,1] \times [0,1]$ whose sides have length less than δ is mapped by H into some open set in X which is evenly covered by the covering map p. It follows from Lemma 4.4 that if the lift \tilde{H} of H has already been determined over a corner, or along one side, or along two adjacent sides of a square whose sides have length less than δ , then \tilde{H} can be extended over the whole of that square. Thus if we subdivide $[0, 1] \times [0, 1]$ into squares $S_{i,k}$, where

$$\begin{split} S_{j,k} &= \{(s,t) \in [0,1] \times [0,1] : \frac{j-1}{n} \le s \le \frac{j}{n} \\ & \text{and} \quad \frac{k-1}{n} \le t \le \frac{k}{n} \}, \end{split}$$

and $1/n < \delta$, then we can construct a lift \tilde{H} of H by defining $\tilde{H}(0,0) = w$, and then successively extending \tilde{H} in turn over each of these smaller squares. (Indeed the map \tilde{H} can be extended successively over the squares

$$S_{1,1}, S_{1,2}, \ldots, S_{1,n}, S_{2,1}, S_{2,2}, \ldots, S_{2,n},$$

$$S_{3,1},\ldots,S_{n-1,n},\ldots,S_{n,1},S_{n,2},\ldots,S_{n,n}$$
)

The uniqueness of \tilde{H} follows from Proposition 4.3.

Homotopies and the Fundamental Group.

Homotopies.

Definition.

Let $f: X \to Y$ and $g: X \to Y$ be continuous maps between topological spaces X and Y. The maps f and g are said to be *homotopic* if there exists a continuous map $H: X \times [0, 1] \to$ Y such that H(x, 0) = f(x) and H(x, 1) =g(x) for all $x \in X$. If the maps f and g are homotopic then we denote this fact by writing $f \simeq g$. The map H with the properties stated above is referred to as a *homotopy* between f and g.

Continuous maps f and g from X to Y are homotopic if and only if it is possible to 'continuously deform' the map f into the map g. **Lemma 5.1** Let X and Y be topological spaces. The homotopy relation \simeq is an equivalence relation on the set of all continuous maps from X to Y.

Proof.

Clearly $f \simeq f$, since $(x,t) \mapsto f(x)$ is a homotopy between f and itself. Thus the relation is reflexive. If $f \simeq g$ then there exists a homotopy

$$H: X \times [0, 1] \to Y$$

between f and g (so that H(x,0) = f(x) and H(x,1) = g(x) for all $x \in X$). But then $(x,t) \mapsto H(x,1-t)$ is a homotopy between g and f. Therefore $f \simeq g$ if and only if $g \simeq f$. Thus the relation is symmetric. Finally, suppose that $f \simeq g$ and $g \simeq h$. Then there exist homotopies $H_1: X \times [0, 1] \to Y$ and $H_2: X \times [0, 1] \to Y$ such that

$$H_1(x,0) = f(x), H_1(x,1) = g(x) = H_2(x,0), H_2(x,1) = h(x)$$

for all $x \in X$. Define $H: X \times [0, 1] \to Y$ by

$$H(x,t) = \begin{cases} H_1(x,2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ H_2(x,2t-1) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Now $H|X \times [0, \frac{1}{2}]$ and $H|X \times [\frac{1}{2}, 1]$ are continuous. It follows from elementary point set topology that H is continuous on $X \times [0, 1]$. Moreover H(x, 0) = f(x) and H(x, 1) = h(x) for all $x \in X$. Thus $f \simeq h$. Thus the relation is transitive. The relation \simeq is therefore an equivalence relation.

Definition.

Let X and Y be topological spaces, and let A be a subset of X. Let $f: X \to Y$ and $g: X \to Y$ be continuous maps from X to some topological space Y, where f|A = g|A (i.e., f(a) = g(a) for all $a \in A$). We say that f and g are homotopic relative to A (denoted by $f \simeq g$ rel A) if and only if there exists a (continuous) homotopy $H: X \times [0,1] \to Y$ such that H(x,0) = f(x) and H(x,1) = g(x) for all $x \in X$ and H(a,t) = f(a) = g(a) for all $a \in A$.

Homotopy relative to a chosen subset of X is also an equivalence relation on the set of all continuous maps between topological spaces X and Y.

The Fundamental Group of a Topological Space.

Definition.

Let X be a topological space, and let x_0 and x_1 be points of X. A path in X from x_0 to x_1 is defined to be a continuous map $\gamma: [0, 1] \rightarrow X$ for which $\gamma(0) = x_0$ and $\gamma(1) = x_1$. A loop in X based at x_0 is defined to be a continuous map $\gamma: [0, 1] \rightarrow X$ for which

$$\gamma(0) = \gamma(1) = x_0.$$
We can concatenate paths. Let $\gamma_1: [0,1] \rightarrow X$ and $\gamma_2: [0,1] \rightarrow X$ be paths in some topological space X. Suppose that $\gamma_1(1) = \gamma_2(0)$. We define the *product path* $\gamma_1.\gamma_2: [0,1] \rightarrow X$ by

$$(\gamma_1.\gamma_2)(t) = \begin{cases} \gamma_1(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \gamma_2(2t-1) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

(The continuity of $\gamma_1.\gamma_2$ may be deduced from Lemma 5.1.)

If $\gamma: [0, 1] \to X$ is a path in X then we define the *inverse path* $\gamma^{-1}: [0, 1] \to X$ by

$$\gamma^{-1}(t) = \gamma(1-t).$$

(Thus if γ is a path from the point x_0 to the point x_1 then γ^{-1} is the path from x_1 to x_0 obtained by traversing γ in the reverse direction.) Let X be a topological space, and let $x_0 \in X$ be some chosen point of X. We define an equivalence relation on the set of all (continuous) loops based at the basepoint x_0 of X, where two such loops γ_0 and γ_1 are equivalent if and only if $\gamma_0 \simeq \gamma_1$ rel $\{0, 1\}$. We denote the equivalence class of a loop $\gamma: [0, 1] \rightarrow X$ based at x_0 by $[\gamma]$. This equivalence class is referred to as the *based homotopy class* of the loop γ . The set of equivalence classes of loops based at x_0 is denoted by $\pi_1(X, x_0)$. Thus two loops γ_0 and γ_1 represent the same element of $\pi_1(X, x_0)$ if and only if $\gamma_0 \simeq \gamma_1$ rel $\{0, 1\}$ (i.e., there exists a homotopy

$F\colon [\mathsf{0},\mathsf{1}]\times [\mathsf{0},\mathsf{1}]\to X$

between γ_0 and γ_1 which maps $(0, \tau)$ and $(1, \tau)$ to x_0 for all $\tau \in [0, 1]$).

Theorem 5.2 Let *X* be a topological space, let x_0 be some chosen point of *X*, and let $\pi_1(X, x_0)$ be the set of all based homotopy classes of loops based at the point x_0 . Then $\pi_1(X, x_0)$ is a group, the group multiplication on $\pi_1(X, x_0)$ being defined according to the rule $[\gamma_1][\gamma_2] = [\gamma_1.\gamma_2]$ for all loops γ_1 and γ_2 based at x_0 .

Proof.

First we show that the group operation on $\pi_1(X, x_0)$ is well-defined. Let γ_1 , γ'_1 , γ_2 and γ'_2 be loops in X based at the point x_0 . Suppose that $[\gamma_1] = [\gamma'_1]$ and $[\gamma_2] = [\gamma'_2]$. Let the map $F: [0, 1] \times [0, 1] \to X$ be defined by

$$F(t,\tau) = \begin{cases} F_1(2t,\tau) & \text{if } 0 \le t \le \frac{1}{2}, \\ F_2(2t-1,\tau) & \text{if } \frac{1}{2} \le t \le 1, \end{cases}$$

where $F_1: [0,1] \times [0,1] \to X$ is a homotopy between γ_1 and γ'_1 , $F_2: [0,1] \times [0,1] \to X$ is a homotopy between γ_2 and γ'_2 , and where the homotopies F_1 and F_2 map $(0,\tau)$ and $(1,\tau)$ to x_0 for all $\tau \in [0,1]$. Then F is itself a homotopy from $\gamma_1.\gamma_2$ to $\gamma'_1.\gamma'_2$, and maps $(0,\tau)$ and $(1,\tau)$ to x_0 for all $\tau \in [0,1]$. Thus $[\gamma_1.\gamma_2] = [\gamma'_1.\gamma'_2]$, showing that the group operation on $\pi_1(X,x_0)$ is well-defined. Next we show that the group operation on $\pi_1(X, x_0)$ is associative. Let γ_1 , γ_2 and γ_3 be loops based at x_0 , and let $\alpha = (\gamma_1.\gamma_2).\gamma_3$. Then $\gamma_1.(\gamma_2.\gamma_3) = \alpha \circ \theta$, where

$$\theta(t) = \begin{cases} \frac{1}{2}t & \text{if } 0 \le t \le \frac{1}{2}; \\ t - \frac{1}{4} & \text{if } \frac{1}{2} \le t \le \frac{3}{4}; \\ 2t - 1 & \text{if } \frac{3}{4} \le t \le 1. \end{cases}$$

Thus the map $G: [0, 1] \times [0, 1] \to X$ defined by $G(t, \tau) = \alpha((1 - \tau)t + \tau\theta(t))$ is a homotopy between $(\gamma_1.\gamma_2).\gamma_3$ and $\gamma_1.(\gamma_2.\gamma_3)$, and moreover this homotopy maps $(0, \tau)$ and $(1, \tau)$ to x_0 for all $\tau \in [0, 1]$. It follows that

$$(\gamma_1.\gamma_2).\gamma_3 \simeq \gamma_1.(\gamma_2.\gamma_3)$$
 rel $\{0,1\}$

and hence

$$([\gamma_1][\gamma_2])[\gamma_3] = [\gamma_1]([\gamma_2][\gamma_3]).$$

This shows that the group operation on $\pi_1(X, x_0)$ is associative.

Let ε : $[0,1] \to X$ denote the constant loop at x_0 , defined by $\varepsilon(t) = x_0$ for all $t \in [0,1]$. Then $\varepsilon \cdot \gamma = \gamma \circ \theta_0$ and $\gamma \cdot \varepsilon = \gamma \circ \theta_1$ for any loop γ based at x_0 , where

$$\theta_0(t) = \begin{cases} 0 & \text{if } 0 \le t \le \frac{1}{2}, \\ 2t - 1 & \text{if } \frac{1}{2} \le t \le 1, \end{cases} \\ \theta_1(t) = \begin{cases} 2t & \text{if } 0 \le t \le \frac{1}{2}, \\ 1 & \text{if } \frac{1}{2} \le t \le 1, \end{cases}$$

for all $t \in [0, 1]$. But the continuous map

$$(t,\tau)\mapsto\gamma((1-\tau)t+\tau\theta_j(t))$$

is a homotopy between γ and $\gamma \circ \theta_j$ for j = 0, 1 which sends $(0, \tau)$ and $(1, \tau)$ to x_0 for all $\tau \in [0, 1]$. Therefore

$$\varepsilon.\gamma\simeq\gamma\simeq\gamma.\varepsilon$$
 rel $\{0,1\},$

and hence

$$[\varepsilon][\gamma] = [\gamma] = [\gamma][\varepsilon].$$

We conclude that $[\varepsilon]$ represents the identity element of $\pi_1(X, x_0)$.

It only remains to verify the existence of inverses. Now the map $K: [0,1] \times [0,1] \rightarrow X$ defined by

$$K(t,\tau) = \begin{cases} \gamma(2\tau t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \gamma(2\tau(1-t)) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

is a homotopy between the loops $\gamma . \gamma^{-1}$ and ε , and moreover this homotopy sends $(0, \tau)$ and $(1, \tau)$ to x_0 for all $\tau \in [0, 1]$. Therefore

$$\gamma . \gamma^{-1} \simeq \varepsilon \operatorname{rel}\{0, 1\},$$

and thus

$$[\gamma][\gamma^{-1}] = [\gamma \cdot \gamma^{-1}] = [\varepsilon].$$

On replacing γ by $\gamma^{-1},$ we see also that

$$[\gamma^{-1}][\gamma] = [\varepsilon],$$

and thus $[\gamma^{-1}] = [\gamma]^{-1}$, as required.

Let x_0 be a point of some topological space X. The group $\pi_1(X, x_0)$ is referred to as the fundamental group of X based at the point x_0 .

Let $f: X \to Y$ be a continuous map between topological spaces X and Y, and let x_0 be a point of X. Then f induces a homomorphism

$$f_{\#}: \pi_1(X, x_0) \to \pi_1(Y, f(x_0)),$$

where $f_{\#}([\gamma]) = [f \circ \gamma]$ for all loops $\gamma: [0, 1] \rightarrow X$ based at x_0 . If x_0 , y_0 and z_0 are points belonging to topological spaces X, Y and Z, and if $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are continuous maps satisfying $f(x_0) = y_0$ and $g(y_0) = z_0$, then the induced homomorphisms

$$f_{\#}: \pi_1(X, x_0) \to \pi_1(Y, y_0)$$

and

$$g_{\#}: \pi_1(Y, x_0) \to \pi_1(Z, z_0)$$

satisfy $g_{\#} \circ f_{\#} = (g \circ f)_{\#}$. It follows easily from this that any homeomorphism of topological spaces induces a corresponding isomorphism of fundamental groups, and thus the fundamental group is a topological invariant.

Simply-Connected Topological Spaces.

Definition.

A topological space X is said to be *simply-connected* if it is path-connected, and any continuous map

$$f:\partial D\to X$$

mapping the boundary circle ∂D of a closed disc D into X can be extended continuously over the whole of the disk.

Example.

 \mathbb{R}^n is simply-connected for all n. Indeed any continuous map $f: \partial D \to \mathbb{R}^n$ defined over the boundary ∂D of the closed unit disk D can be extended to a continuous map $F: D \to \mathbb{R}^n$ over the whole disk by setting $F(r\mathbf{x}) = rf(\mathbf{x})$ for all $\mathbf{x} \in \partial D$ and $r \in [0, 1]$. Let E be a topological space that is homeomorphic to the closed disk D, and let $\partial E = h(\partial D)$, where ∂D is the boundary circle of the disk D and $h: D \to E$ is a homeomorphism from D to E. Then any continuous map $g: \partial E \to X$ mapping ∂E into a simplyconnected space X extends continuously to the whole of E. Indeed there exists a continuous map $F: D \to X$ which extends $g \circ h: \partial D \to$ X, and the map $F \circ h^{-1}: E \to X$ then extends the map g. **Theorem 5.3** A path-connected topological space X is simply-connected if and only if $\pi_1(X, x)$ is trivial for all $x \in X$.

Proof.

Suppose that X is simply-connected. Let $\gamma: [0,1] \to X$ be a loop based at some point x of X. Now the unit square is homeomorphic to the unit disk, and therefore any continuous map defined over the boundary of the square can be continuously extended over the whole of the square. It follows that there exists a continuous map $F: [0,1] \times [0,1] \to X$ such that $F(t,0) = \gamma(t)$ and F(t,1) = x for all $t \in [0,1]$, and $F(0,\tau) = F(1,\tau) = x$ for all $\tau \in [0,1]$. Thus $\gamma \simeq \varepsilon_x \operatorname{rel}\{0,1\}$, where ε_x is the constant loop at x, and hence $[\gamma] = [\varepsilon_x]$ in $\pi_1(X,x)$. This shows that $\pi_1(X,x)$ is trivial.

Conversely suppose that X is path-connected and $\pi_1(X, x)$ is trivial for all $x \in X$. Let $f: \partial D \to X$ be a continuous function defined on the boundary circle ∂D of the closed unit disk D in \mathbb{R}^2 . We must show that f can be extended continuously over the whole of D. Let x = f(1,0). There exists a continuous map $G: [0,1] \times [0,1] \to X$ such that

$$G(t,0) = f(\cos(2\pi t), \sin(2\pi t))$$

and G(t, 1) = x for all $t \in [0, 1]$ and

 $G(0,\tau) = G(1,\tau) = x$

for all $\tau \in [0, 1]$, since $\pi_1(X, x)$ is trivial. Moreover $G(t_1, \tau_1) = G(t_2, \tau_2)$ whenever $q(t_1, \tau_1) = q(t_2, \tau_2)$, where

$$q(t,\tau) = ((1-\tau)\cos(2\pi t) + \tau, (1-\tau)\sin(2\pi t))$$

for all $t,\tau \in [0,1]$. It follows that there is
a well-defined function $F: D \to X$ such that
 $F \circ q = G$.

However $q: [0, 1] \times [0, 1] \rightarrow D$ is a continuous surjection from a compact space to a Hausdorff space and is therefore an identification map. It follows that $F: D \rightarrow X$ is continuous (since a basic property of identification maps ensures that a function $F: D \rightarrow X$ is continuous if and only if $F \circ q: [0, 1] \times [0, 1] \rightarrow X$ is continuous). Moreover $F: D \rightarrow X$ extends the map f. We conclude that the space X is simply-connected, as required.

One can show that, if two points x_1 and x_2 in a topological space X can be joined by a path in X then $\pi_1(X, x_1)$ and $\pi_1(X, x_2)$ are isomorphic. On combining this result with Theorem 5.3, we see that a path-connected topological space X is simply-connected if and only if $\pi_1(X, x)$ is trivial for some $x \in X$.

Theorem 5.4 Let X be a topological space, and let U and V be open subsets of X, with $U \cup V = X$. Suppose that U and V are simplyconnected, and that $U \cap V$ is non-empty and path-connected. Then X is itself simplyconnected.

Proof.

We must show that any continuous function $f: \partial D \to X$ defined on the unit circle ∂D can be extended continuously over the closed unit disk D. Now the preimages $f^{-1}(U)$ and $f^{-1}(V)$ of U and V are open in ∂D (since f is continuous), and $\partial D = f^{-1}(U) \cup f^{-1}(V)$. It follows from the Lebesgue Lemma that there exists some $\delta > 0$ such that any arc in ∂D whose length is less than δ is entirely contained in one or other of the sets $f^{-1}(U)$ and $f^{-1}(V)$. Choose points z_1, z_2, \ldots, z_n around ∂D such that the distance from z_i to z_{i+1} is less than δ for $i = 1, 2, \ldots, n-1$ and the distance from z_n to z_1 is also less than δ . Then, for each i, the short arc joining z_{i-1} to z_i is mapped by f into one or other of the open sets U and V.

Let x_0 be some point of $U \cap V$. Now the sets U, V and $U \cap V$ are all path-connected. Therefore we can choose paths $\alpha_i:[0,1] \rightarrow X$ for i = 1, 2, ..., n such that $\alpha_i(0) = x_0$, $\alpha_i(1) = f(z_i), \alpha_i([0,1]) \subset U$ whenever $f(z_i) \in U$, and $\alpha_i([0,1]) \subset V$ whenever $f(z_i) \in V$. For convenience let $\alpha_0 = \alpha_n$.

Now, for each i, consider the sector T_i of the closed unit disk bounded by the line segments joining the centre of the disk to the points z_{i-1} and z_i and by the short arc joining z_{i-1} to z_i . Now this sector is homeomorphic to the closed unit disk, and therefore any continuous function mapping the boundary ∂T_i of T_i into a simply-connected space can be extended continuously over the whole of T_i .

In particular, let F_i be the function on ∂T_i defined by

$$F_i(z) = \begin{cases} f(z) & \text{if } z \in T_i \cap \partial D, \\ \alpha_{i-1}(t) & \text{if } z = tz_{i-1} \\ & \text{for some } t \in [0, 1], \\ \alpha_i(t) & \text{if } z = tz_i \\ & \text{for some } t \in [0, 1], \end{cases}$$

Note that $F_i(\partial T_i) \subset U$ whenever the short arc joining z_{i-1} to z_i is mapped by f into U, and $F_i(\partial T_i) \subset V$ whenever this short arc is mapped into V. But U and V are both simplyconnected. It follows that each of the functions F_i can be extended continuously over the whole of the sector T_i . Moreover the functions defined in this fashion on each of the sectors T_i agree with one another wherever the sectors intersect, and can therefore be pieced together to yield a continuous map defined over the the whole of the closed disk D which extends the map f, as required.

Example.

Let S^n be the $n\mbox{-dimensional sphere, defined}$ by

$$S^n = \{ \mathbf{x} \in \mathbb{R}^{n+1} : |\mathbf{x}| = 1 \}.$$

Then S^n is simply-connected for all n > 1. Indeed let $U = \{x \in S^n : x_{n+1} > -\frac{1}{2}\}$ and $V = \{x \in S^n : x_{n+1} < \frac{1}{2}\}$. Then U and Vare homeomorphic to an n-dimensional ball, and are therefore simply-connected. Moreover $U \cap V$ is path-connected, provided that n > 1. It follows that S^n is simply-connected for all n > 1.

Monodromy.

The Fundamental Group of the Punctured Plane.

Let \boldsymbol{X} be the punctured plane, defined such that

$$X = \{ (x, y) \in \mathbb{R}^2 : (x, y) \neq (0, 0) \},\$$

let \tilde{X} be the helicoidal surface in \mathbb{R}^3 , defined such that

$$\tilde{X} = \{(x, y, z) \in \mathbb{R}^3 : x \sin 2\pi z = y \cos 2\pi z \\ \text{and } x \cos 2\pi z + y \sin 2\pi z > 0\}.$$

and let $p_{\text{vert}}: \tilde{X} \to X$ be the covering map with covering space \tilde{X} and base space X defined by vertical projection, so that

$$p_{\mathsf{vert}}(x,y,z) = (x,y)$$

for all $(x,y) \in \tilde{X}$.

A point (x, y, z) of \mathbb{R}^3 lies on the helicoidal surface \tilde{X} if and only if there exists some positive real number ρ such that

 $x = \rho \cos 2\pi z \quad \text{and} \quad y = \rho \sin 2\pi z,$ in which case $\rho = \sqrt{x^2 + y^2}$ (see Lemma 4.1).

Let $\gamma: [0,1] \to X$ be a path in X, and let (x_0, y_0, z_0) be a point of the helicoidal surface for which

$$p_{\text{vert}}(x_0, y_0, z_0) = (x_0, y_0) = \gamma(0).$$

It follows from the Path Lifting Theorem (Theorem 4.5) that there exists a lift

$$\tilde{\gamma}$$
: [0, 1] $\rightarrow \tilde{X}$

of the path γ to the helicoidal surface \tilde{X} starting at the point (x_0, y_0, z_0) . Then

$$\tilde{\gamma}(0) = (x_0, y_0, z_0)$$

and

$$p_{\text{vert}}(\tilde{\gamma}(t)) = \gamma(t)$$

for all $t \in [0, 1]$.

164

Let

$$\tilde{\gamma}(t) = (x(t), y(t), z(t))$$

for all $t \in [0, 1]$, where $x: [0, 1] \to \mathbb{R}$, $y: [0, 1] \to \mathbb{R}$ and $z: [0, 1] \to \mathbb{R}$ are maps expressing the Cartesian coordinates of the point $\tilde{\gamma}(t)$ as functions of the parameter t. Then

$$\begin{aligned} \gamma(t) &= (x(t), y(t)), \\ x(t) &= \rho(t) \cos 2\pi z(t), \\ y(t) &= \rho(t) \sin 2\pi z(t) \end{aligned}$$

for all $t \in [0, 1]$, where

$$\rho(t) = \sqrt{x(t)^2 + y(t)^2}$$

for all $t \in [0, 1]$.

Now suppose that the path γ is a loop in the punctured plane X, starting and ending at a point (x_0, y_0) . Then $\tilde{\gamma}(0) = (x_0, y_0, z_0)$ for some $z_0 \in [0, 1]$. Also

$$p_{\text{vert}}(\tilde{\gamma}(1)) = \gamma(1) = \gamma(0) = (x_0, y_0).$$

It follows that

$$\tilde{\gamma}(1) = (x_0, y_0, z_0 + n(\gamma))$$

for some integer $n(\gamma)$. This integer $n(\gamma)$ is determined by the loop γ , but is independent of the choice of the lift $\tilde{\gamma}$ of that loop. Indeed suppose that $\eta: [0,1] \to \tilde{X}$ is also a lift of the loop γ to the helicoidal surface \tilde{X} . Let $\eta(0) = (x_0, y_0, z'_0)$. Then $z'_0 = z_0 + m$ for some integer m. Now the map that sends $t \in [0,1]$ to $\tilde{\gamma}(t) + m$ is a lift of the loop γ starting at the point $\eta(0)$. But the Path Lifting Theorem (Theorem 4.5) guarantees that any lift of the loop γ is uniquely determined by its starting point. Therefore $\eta(t) = \tilde{\gamma}(t) + m$ for all $t \in [0,1]$, and therefore

 $\eta(1) = \tilde{\gamma}(1) + m = z_0 + n(\gamma) + m = z'_0 + n(\gamma).$ We refer to the integer $n(\gamma)$ as the *winding number* of the loop γ about the point (0,0)of $\mathbb{R}^2 \setminus X$. Now let α : $[0,1] \to X$ and β : $[0,1] \to X$ be loops in X based at the point (x_0, y_0) , so that

$$\alpha(0) = \alpha(1) = \beta(0) = \beta(1) = (x_0, y_0).$$

Suppose that $\alpha \simeq \beta$ rel $\{0,1\}$. Then there exists a continuous map $F: [0,1] \times [0,1] \rightarrow X$ from the unit square $[0,1] \times [0,1]$ to the punctured plane X such that

 $F(t,0) = \alpha(t)$ and $F(t,1) = \beta(t)$

for all $t \in [0, 1]$ and

$$F(0,\tau) = F(1,\tau) = (x_0, y_0)$$

for all $\tau \in [0,1]$. It then follows from the Monodromy Theorem (Theorem 4.6) that there exists a continuous map

$$G: [0, 1] \times [0, 1] \rightarrow \tilde{X}$$

from the unit square $[0,1]\times[0,1]$ to the helicoidal surface \tilde{X} such that

$$p_{\mathsf{vert}}(G(t,\tau)) = F(t,\tau))$$

for all $t, \tau \in [0, 1]$.

Let

$$G(t,\tau) = (u(t,\tau), v(t,\tau), w(t,\tau))$$

for all $t, \tau \in [0, 1]$, where $u: [0, 1] \times [0, 1] \rightarrow \mathbb{R}$, $v: [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ and $w: [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ are maps expressing the Cartesian coordinates of the point $G(t, \tau)$ as functions of the parameters t and τ . Then

$$F(t,\tau) = (u(t,\tau), v(t,\tau)),$$

$$u(t,\tau) = s(t,\tau) \cos 2\pi w(t,\tau),$$

$$v(t,\tau) = s(t,\tau) \sin 2\pi w(t,\tau)$$

for all $t, \tau \in [0, 1]$, where

$$s(t,\tau) = \sqrt{x(t,\tau)^2 + y(t,\tau)^2}$$

for all $t, \tau \in [0, 1]$.

Let $w_0 = w(0,0)$. Now the map that sends $t \in [0,1]$ to G(t,0) is a lift of the loop α . It follows from the definition of the winding number $n(\alpha)$ that $w(1,0) = w_0 + n(\alpha)$. Also the maps that send $\tau \in [0,1]$ to $G(0,\tau)$ and $G(1,\tau)$ are lifts of constant loops based the the point (0,0), and must therefore themselves be constant paths. It follows that $G(0,1) = w_0$ and $G(1,1) = w_0 + n(\alpha)$. But the map that sends $t \in [0,1]$ to G(t,1) is a lift of the loop β starting at the point w_0 , and therefore $G(1,1) = w_0 + n(\beta)$. This shows that $n(\alpha) = n(\beta)$. We have now shown that each loop γ in the punctured plane X based at the point (x_0, y_0) determines an integer $n(\gamma)$ that represents the winding number of the loop about the point (0,0) of $\mathbb{R}^2 \setminus X$. Moreover if two loops based at (x_0, y_0) represent the same element of the fundamental group $\pi_1(X, (x_0, y_0))$ then they have the same winding number about (0,0). It follows that there is a well-defined function

 $\lambda:\pi_1(X,(x_0,y_0))\to\mathbb{Z}$

defined such that $\lambda([\gamma]) = n(\gamma)$ for all loops γ based at the point (x_0, y_0) .

The function

$$\lambda$$
: $\pi_1(X, (x_0, y_0)) \to \mathbb{Z}$

is a homomorphism. Indeed let $\alpha: [0,1] \rightarrow X$ and $\beta: [0,1] \rightarrow X$ be loops in X based at the point (x_0, y_0) , and let $\alpha.\beta: [0,1] \rightarrow X$ be the concatenation of the two loops α and β , defined such that

$$(\alpha.\beta)(t) = \begin{cases} \alpha(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \beta(2t-1) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Let z_0 be a real number for which $(x_0, y_0, z_0) \in \tilde{X}$. Then there exists a lift of the loop α that is a path in the helicoidal surface \tilde{X} starting at (x_0, y_0, z_0) and finishing at (x_0, y_0, z_1) , where $z_1 = z_0 + n(\alpha)$. Also there exists a lift of the loop β that is a path in the helicoidal surface \tilde{X} starting at (x_0, y_0, z_1) and finishing at (x_0, y_0, z_2) , where $z_2 = z_1 + n(\beta)$. But then the concatenation of these two paths in the helicoidal surface \tilde{X} is a path in \tilde{X} that is a lift of the loop $\alpha.\beta$ starting at (x_0, y_0, z_0) and finishing at (x_0, y_0, z_2) . It follows that

$$n(\alpha.\beta) = z_2 - z_0 = n(\alpha) + n(\beta).$$

This shows that

$$\lambda$$
: $\pi_1(X, (x_0, y_0)) \to \mathbb{Z}$

is indeed a homomorphism.

Now let α and β be loops in the punctured plane X based at the point (x_0, y_0) , and let $\tilde{\alpha}$ and $\tilde{\beta}$ be paths in the helicoidal surface \tilde{X} starting at some point (x_0, y_0, z_0) of that surface, where $\tilde{\alpha}$ and $\tilde{\beta}$ are lifts of the loops α and β respectively. Suppose that

$$\lambda([\alpha]) = \lambda([\beta]).$$

Then

$$\tilde{\alpha}(1) = z_0 + n(\alpha) = z_0 + \lambda([\alpha])$$
$$= z_0 + \lambda([\beta]) = z_0 + n(\beta) = \tilde{\beta}(1).$$

Now there exist continuous maps

$$\rho_{\alpha}: [0, 1] \to \mathbb{R}, \quad \rho_{\beta}: [0, 1] \to \mathbb{R},$$

 $z_{\alpha}: [0,1] \to \mathbb{R}$ and $z_{\beta}: [0,1] \to \mathbb{R}$

such that $ho_{lpha}(t)>0$, $ho_{eta}(t)>0$,

 $\tilde{\alpha}(t) = (\rho_{\alpha}(t) \cos 2\pi z_{\alpha}(t), \ \rho_{\alpha}(t) \sin 2\pi z_{\alpha}(t), \ z_{\alpha}(t))$ and

$$\tilde{\beta}(t) = (\rho_{\beta}(t) \cos 2\pi z_{\beta}(t), \, \rho_{\beta}(t) \sin 2\pi z_{\beta}(t), \, z_{\beta}(t)).$$

Let

$$G(t,\tau) = (\rho(t,\tau)\cos 2\pi z(t,\tau), \rho(t,\tau)\sin 2\pi z(t,\tau), z(t,\tau)),$$

where

$$\rho(t,\tau) = (1-\tau)\rho_{\alpha}(t) + \tau \rho_{\beta}(t),$$

and

$$z(t,\tau) = (1-\tau)z_{\alpha}(t) + \tau z_{\beta}(t).$$

Then $\rho(t,\tau) > 0$ for all $t,\tau \in [0,1]$, and therefore the function G is a homotopy in \tilde{X} between the paths $\tilde{\alpha}$ and $\tilde{\beta}$. Let

$$F(t,\tau) = p_{\mathsf{vert}}(G(t,\tau))$$

for all $t, \tau \in [0, 1]$. Then $F(t, 0) = \alpha(t)$ and $F(t, 1) = \beta(t)$ for all $t \in [0, 1]$, and thus the map $F: [0, 1] \times [0, 1] \rightarrow X$ is a homotopy between the loops α and β . Also

$$F(0,\tau) = F(1,\tau) = (x_0, y_0)$$

for all $\tau \in [0,1]$, because $\tilde{\alpha}(0) = \tilde{\beta}(0)$ and $\tilde{\alpha}(1) = \tilde{\beta}(1)$. It follows that the loops α and β represent the same element of the fundamental group $\pi_1(X, (x_0, y_0))$.

We have accordingly shown that if α and β are loops in the punctured plane X based at the point (x_0, y_0) , and if $\lambda([\alpha]) = \lambda([\beta])$, then $\alpha \simeq \beta$ rel {0, 1}, and therefore $[\alpha] = [\beta]$ in $\pi_1(X, (x_0, y_0))$. Thus the homomorphism

$$\lambda:\pi_1(X,(x_0,y_0))\to\mathbb{Z}$$

is injective.

Given any integer m, let $\eta: [0,1] \to \tilde{Z}$ be a path in helicoidal surface \tilde{X} which starts at the point (x_0, y_0, z_0) and ends at $(x_0, y_0, z_0 + m)$, and let $\gamma = p_{\text{vert}} \circ \eta$. Then $\gamma: [0,1] \to X$ is a loop in the punctured plane X based at the point (x_0, y_0) , and moreover

$$\lambda[\gamma]) = n(\gamma) = m.$$

This shows that the homomorphism

$$\lambda$$
: $\pi_1(X, (x_0, y_0)) \to \mathbb{Z}$

is surjective.

We have thus constructed an isomorphism $\lambda: \pi_1(X, (x_0, y_0)) \to \mathbb{Z}$ between the fundamental group $\pi_1(X, (x_0, y_0))$ of the punctured plane X and the group \mathbb{Z} of integers under addition.

Remark.

We showed that the fundamental group of the punctured plane X is isomorphic to the group of integers under addition by exploiting properties of the covering map $p_{\text{vert}}: \tilde{X} \to X$. The methods used can be generalized so as to find the fundamental group $\pi_1(Z, z_0)$ of the base space Z of a covering map $p: \tilde{Z} \to$ Z for which the covering space \tilde{Z} is simplyconnected. In such cases, the elements of the fundamental group $\pi_1(Z, z_0)$ are in one-toone correspondence with the elements of the preimage $p^{-1}(\{z_0\})$ of the base point under the covering map.

The Fundamental Group of the Circle.

Theorem 6.1 $\pi_1(S^1, b) \cong \mathbb{Z}$ for any $b \in S^1$.

Proof.

We regard S^1 as the unit circle in \mathbb{R}^2 . Without loss of generality, we can take b = (1,0). Now the map $p: \mathbb{R} \to S^1$ which sends $t \in \mathbb{R}$ to $(\cos 2\pi t, \sin 2\pi t)$ is a covering map, and b = p(0). Moreover $p(t_1) = p(t_2)$ if and only if $t_1 - t_2$ is an integer; in particular p(t) = bif and only if t is an integer.

Let $\alpha: [0,1] \to S^1$ be a loop in S^1 . It follows from the Path Lifting Theorem (Theorem 4.5) that there exists a continuous function $\tilde{\alpha}: [0,1] \to \mathbb{R}$ such that $p \circ \tilde{\alpha} = \alpha$. Then $p(\tilde{\alpha}(1)) = b = p(\tilde{\alpha}(0))$, and therefore $\tilde{\alpha}(1) - \tilde{\alpha}(0)$ is an integer. We show that the value of this integer does not depend on the choice of the function $\tilde{\alpha}$.

177

Let $\eta: [0,1] \to \mathbb{R}$ be a map from $[0,1] \to \mathbb{R}$ that satisfies $p \circ \eta = \alpha$, and let

$$m = \eta(0) - \tilde{\alpha}(0).$$

Then the function that sends $t \in [0,1]$ to $\tilde{\alpha}(t) + m$ is a continuous function on the interval [0,1], and

$$p(\eta(t)) = \alpha(t) = p(\tilde{\alpha}(t) + m)$$

for all $t \in [0, 1]$. It follows from this that

$$\eta(t) = \tilde{\alpha}(t) + m$$

for all $t \in [0, 1]$. This result is a consequence of the fact that lifts of paths from a base space to a covering space are uniquely determined by their starting point (see Theorem 4.5). Note also that the function that maps $t \in [0, 1]$ to $\eta(t) - \tilde{\alpha}(t) - m$ is a continuous integer-valued function on the interval [0, 1], and the connectedness of the interval [0, 1] ensures that every continuous integervalued function [0, 1] must be constant. Thus

$$\eta(1) - \eta(0) = \tilde{\alpha}(1) - \tilde{\alpha}(0).$$

Given any loop α in S^1 based at the point b, we define the *winding number* $n(\alpha)$ of the loop α to be the unique integer with the property that

$$n(\alpha) = \tilde{\alpha}(1) - \tilde{\alpha}(0)$$

for all paths $\tilde{\alpha}$: $[0,1] \to \mathbb{R}$ in \mathbb{R} satisfying

$$p \circ \tilde{\alpha} = \alpha.$$

Now let α and β be loops in S^1 based at b that represent the same element of $\pi_1(S^1, b)$. Then there exists a homotopy

$$F: [0,1] \times [0,1] \rightarrow S^1$$

such that $F(t,0) = \alpha(t)$ and $F(t,1) = \beta(t)$ for all $t \in [0,1]$, and $F(0,\tau) = F(1,\tau) = b$ for all $\tau \in [0,1]$. It follows from the Monodromy Theorem (Theorem 4.6) that this homotopy lifts to a continuous map

$$G: [0,1] \times [0,1] \rightarrow \mathbb{R}$$

satisfying $p \circ G = F$.

Now the functions that send $\tau \in [0,1]$ to $G(0,\tau)$ and $G(1,\tau)$ are constant functions on the interval [0,1], because they are continuous functions of τ that satisfy

$$p(G(0,\tau)) = b = p(G(1,\tau))$$

for all $\tau \in [0, 1]$. Thus G(0, 0) = G(0, 1) and G(1, 0) = G(1, 1). Let

$$\tilde{\alpha}(t) = G(t, 0)$$

and

$$\tilde{\beta}(t) = G(t, 1)$$

for all $t \in [0, 1]$, and let

$$\tilde{b}_0 = G(0,0) = G(0,1)$$

and

$$\tilde{b}_1 = G(1,0) = G(1,1).$$

Then $p \circ \tilde{\alpha} = \alpha$ and $p \circ \tilde{\beta} = \beta$, and thus the paths $\tilde{\alpha}$ and $\tilde{\beta}$ are lifts of the loops α and β respectively that start at \tilde{b}_0 and end at \tilde{b}_1 .
It follows that

$$n(\alpha) = \tilde{b}_1 - \tilde{b}_0 = n(\beta).$$

Thus there is a well-defined function

$$\lambda: \pi_1(S^1, b) \to \mathbb{Z}$$

with the property that $\lambda([\alpha]) = n(\alpha)$ for all loops α based at the point b.

Next we show that λ is a homomorphism. Let α and β be any loops based at b, and let $\tilde{\alpha}$ and $\tilde{\beta}$ be lifts of α and β . The element $[\alpha][\beta]$ of $\pi_1(S^1, b)$ is represented by the product path $\alpha.\beta$, where

$$(\alpha.\beta)(t) = \begin{cases} \alpha(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \beta(2t-1) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Define a continuous path $\sigma: [0,1] \to \mathbb{R}$ by

$$\sigma(t) = \begin{cases} \tilde{\alpha}(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \tilde{\beta}(2t-1) + \tilde{\alpha}(1) - \tilde{\beta}(0) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$
(Note that $\sigma(t)$ is well-defined when $t = \frac{1}{2}$.)

Then $p \circ \sigma = \alpha . \beta$ and thus

$$\lambda([\alpha][\beta]) = \lambda([\alpha.\beta]) = \sigma(1) - \sigma(0)$$

= $\tilde{\alpha}(1) - \tilde{\alpha}(0) + \tilde{\beta}(1) - \tilde{\beta}(0)$
= $\lambda([\alpha]) + \lambda([\beta]).$

Thus $\lambda: \pi_1(S^1, b) \to \mathbb{Z}$ is a homomorphism.

Now suppose that $\lambda([\alpha]) = \lambda([\beta])$. Let

 $F: [0,1] \times [0,1] \rightarrow S^1$

be the homotopy between α and β defined by

$$F(t,\tau) = p\left((1-\tau)\tilde{\alpha}(t) + \tau\tilde{\beta}(t)\right),\,$$

where $\tilde{\alpha}$ and $\tilde{\beta}$ are the lifts of α and β respectively starting at 0. Now

$$\tilde{\beta}(1) = \lambda([\beta]) = \lambda([\alpha]) = \tilde{\alpha}(1),$$

and $\tilde{\beta}(0) = \tilde{\alpha}(0) = 0$. Therefore

$$F(0,\tau) = b = p(\tilde{\alpha}(1)) = F(1,\tau)$$

for all $\tau \in [0,1]$. Thus $\alpha \simeq \beta$ rel $\{0,1\}$, and therefore $[\alpha] = [\beta]$. This shows that $\lambda: \pi_1(S^1, b) \to \mathbb{Z}$ is injective.

182

The homomorphism λ is surjective, since

$$n = \lambda([\gamma_n])$$

for all $n \in \mathbb{Z}$, where the loop $\gamma_n: [0, 1] \to S^1$ is given by

$$\gamma_n(t) = p(nt) = (\cos 2\pi nt, \sin 2\pi nt)$$

for all $t \in [0, 1]$. We conclude that

$$\lambda$$
: $\pi_1(S^1, b) \to \mathbb{Z}$

is an isomorphism.

Remark.

Let S^1 be the unit circle in the punctured plane X, let $i: S^1 \to X$ be the inclusion map, and let $r: X \to S^1$ be the radial projection map, defined such that

$$r(x,y) = \left(\frac{x}{\sqrt{x^2 + y^2}}, \frac{y}{\sqrt{x^2 + y^2}}\right)$$

for all $(x, y) \in X$. Now the composition map $r \circ i$ is the identity map of S^1 .

183

Let

$$u(x, y, \tau) = \frac{1 - \tau}{\sqrt{x^2 + y^2}} + \tau$$

for all $(x, y) \in X$ and $\tau \in [0, 1]$. Then the function $F: X \times [0, 1] \rightarrow X$ that sends

$$((x,y),\tau) \in X \times [0,1]$$

to

is a homotopy between the composition map $i \circ r$ and the identity map of the punctured plane X. It follows that the homomorphism

$$i_{\#}: \pi_1(S^1, (1, 0)) \to \pi_1(X, (1, 0))$$

of fundamental groups induced by the inclusion map $i: S^1 \to X$ is an isomorphism whose inverse is the homomorphism

$$r_{\#}: \pi_1(X, (1, 0)) \to \pi_1(S^1, (1, 0))$$

of fundamental groups induced by the radial projection map $r: X \to S^1$. Therefore

$$\pi_1(S^1, (1, 0)) \cong \pi_1(X, (1, 0)) \cong \mathbb{Z}.$$

Indeed the function that sends each loop in S^1 based at the point (1,0) to its winding number about the origin (0,0) induces an isomorphism from $\pi_1(S^1, (1,0))$ to \mathbb{Z} . Thus the result of Theorem 6.1 can be deduced as a corollary of the previous discussion concerning the fundamental group of the punctured plane. Indeed the proof of Theorem 6.1 utilizes the same basic methods and results used to find the fundamental group of the punctured plane. We shall develop the theory of the relationship between covering spaces and the fundamental group in order to find the fundamental groups of other topological spaces that are base spaces of covering maps for which the corresponding covering spaces are simply-connected.

Covering Maps and Induced Homomorphisms of the Fundamental Group.

Let $p: \tilde{X} \to X$ be a covering map and let $\alpha: [0,1] \to X$ and $\beta: [0,1] \to X$ be paths in the base space X which both start at some point x_0 of X and finish at some point x_1 of X, so that

 $\alpha(0) = \beta(0) = x_0$ and $\alpha(1) = \beta(1) = x_1$. Let \tilde{x}_0 be some point of the covering space \tilde{X} that projects down to x_0 , so that $p(\tilde{x}_0) = x_0$. It follows from the Path Lifting Theorem (Theorem 4.5) that there exist paths $\tilde{\alpha}: [0,1] \to \tilde{X}$ and $\tilde{\beta}: [0,1] \to \tilde{X}$ in the covering space \tilde{X} that both start at \tilde{x}_0 and that are lifts of the paths α and β respectively. Thus

$$\tilde{\alpha}(0) = \tilde{\beta}(0) = \tilde{x}_0,$$
$$p(\tilde{\alpha}(t)) = \alpha(t)$$

and

$$p(\tilde{\beta}(t)) = \beta(t)$$

for all $t \in [0, 1]$. These lifts $\tilde{\alpha}$ and $\tilde{\beta}$ of the paths α and β are uniquely determined by their starting point \tilde{x}_0 (see Proposition 4.3).

Now, though the lifts $\tilde{\alpha}$ and $\tilde{\beta}$ of the paths α and β have been chosen such that they start at the same point \tilde{x}_0 of the covering space \tilde{X} , they need not in general end at the same point of \tilde{X} . However we shall prove that if $\alpha \simeq \beta$ rel {0,1}, then the lifts $\tilde{\alpha}$ and $\tilde{\beta}$ of α and β respectively that both start at some point \tilde{x}_0 of \tilde{X} will both finish at some point \tilde{x}_1 of \tilde{x} , so that $\tilde{\alpha}(1) = \tilde{\beta}(1) = \tilde{x}_1$. This result is established in Proposition 6.2 below. **Proposition 6.2** Let $p: \tilde{X} \to X$ be a covering map over a topological space X, let $\alpha: [0,1] \to X$ and $\beta: [0,1] \to X$ be paths in X, where $\alpha(0) = \beta(0)$ and $\alpha(1) = \beta(1)$, and let $\tilde{\alpha}: [0,1] \to \tilde{X}$ and $\tilde{\beta}: [0,1] \to \tilde{X}$ be paths in \tilde{X} such that $p \circ \tilde{\alpha} = \alpha$ and $p \circ \tilde{\beta} = \beta$. Suppose that $\tilde{\alpha}(0) = \tilde{\beta}(0)$ and that $\alpha \simeq \beta$ rel $\{0,1\}$. Then $\tilde{\alpha}(1) = \tilde{\beta}(1)$ and $\tilde{\alpha} \simeq \tilde{\beta}$ rel $\{0,1\}$.

Proof.

Let x_0 and x_1 be the points of X given by

 $x_0 = \alpha(0) = \beta(0),$ $x_1 = \alpha(1) = \beta(1).$ Now $\alpha \simeq \beta$ rel {0,1}, and therefore there exists a homotopy $F: [0,1] \times [0,1] \rightarrow X$ such

that

 $F(t,0) = \alpha(t)$ and $F(t,1) = \beta(t)$ for all $t \in [0,1]$, and

 $F(0,\tau) = x_0$ and $F(1,\tau) = x_1$

for all $\tau \in [0, 1]$.

It then follows from the Monodromy Theorem (Theorem 4.6) that there exists a continuous map $G: [0,1] \times [0,1] \rightarrow \tilde{X}$ such that $p \circ G = F$ and $G(0,0) = \tilde{\alpha}(0)$. Then

$$p(G(0,\tau)) = x_0$$

and

$$p(G(1,\tau)) = x_1$$

for all $\tau \in [0,1]$. A straightforward application of Proposition 4.3 shows that any continuous lift of a constant path must itself be a constant path. Therefore $G(0,\tau) = \tilde{x}_0$ and $G(1,\tau) = \tilde{x}_1$ for all $\tau \in [0,1]$, where

$$\tilde{x}_0 = G(0,0) = \tilde{\alpha}(0), \qquad \tilde{x}_1 = G(1,0).$$

However

$$G(0,0) = G(0,1) = \tilde{x}_0 = \tilde{\alpha}(0) = \tilde{\beta}(0),$$

$$p(G(t,0)) = F(t,0) = \alpha(t) = p(\tilde{\alpha}(t))$$

and

$$p(G(t,1)) = F(t,1) = \beta(t) = p(\tilde{\beta}(t))$$

for all $t \in [0, 1]$.

It follows that the map that sends $t \in [0, 1]$ to G(t, 0) is a lift of the path α that starts at \tilde{x}_0 , and the map that sends $t \in [0, 1]$ to G(t, 1) is a lift of the path β that also starts at \tilde{x}_0 . However Proposition 4.3 ensures that the lifts $\tilde{\alpha}$ and $\tilde{\beta}$ of the paths α and β are uniquely determined by their starting points. It follows that $G(t, 0) = \tilde{\alpha}(t)$ and $G(t, 1) = \tilde{\beta}(t)$ for all $t \in [0, 1]$. In particular,

 $\tilde{\alpha}(1) = G(1,0) = \tilde{x}_1 = G(1,1) = \tilde{\beta}(1).$

Moreover the map $G: [0,1] \times [0,1] \rightarrow \tilde{X}$ is a homotopy between the paths $\tilde{\alpha}$ and $\tilde{\beta}$ which satisfies $G(0,\tau) = \tilde{x}_0$ and $G(1,\tau) = \tilde{x}_1$ for all $\tau \in [0,1]$. It follows that $\tilde{\alpha} \simeq \tilde{\beta}$ rel $\{0,1\}$, as required.

Homomorphisms of Fundamental Groups induced by Covering Maps.

Proposition 6.3 Let $p: \tilde{X} \to X$ be a covering map, and let \tilde{x}_0 be a point of the covering space \tilde{X} . Then the homomorphism

 $p_{\#}: \pi_1(\tilde{X}, \tilde{x}_0) \to \pi_1(X, p(\tilde{x}_0))$

of fundamental groups induced by the covering map p is injective.

Proof.

Let σ_0 and σ_1 be loops in \tilde{X} based at the point \tilde{x}_0 , representing elements $[\sigma_0]$ and $[\sigma_1]$ of $\pi_1(\tilde{X}, \tilde{x}_0)$. Suppose that $p_{\#}[\sigma_0] = p_{\#}[\sigma_1]$. Then $p \circ \sigma_0 \simeq p \circ \sigma_1$ rel $\{0, 1\}$. Also $\sigma_0(0) =$ $\tilde{x}_0 = \sigma_1(0)$. Therefore $\sigma_0 \simeq \sigma_1$ rel $\{0, 1\}$, by Proposition 6.2, and thus $[\sigma_0] = [\sigma_1]$. We conclude that the homomorphism

$$p_{\#}: \pi_1(\tilde{X}, \tilde{x}_0) \to \pi_1(X, p(\tilde{x}_0))$$

is injective.

Proposition 6.4 Let $p: \tilde{X} \to X$ be a covering map, let \tilde{x}_0 be a point of the covering space \tilde{X} , and let γ be a loop in X based at $p(\tilde{x}_0)$. Then $[\gamma] \in p_{\#}(\pi_1(\tilde{X}, \tilde{x}_0))$ if and only if there exists a loop $\tilde{\gamma}$ in \tilde{X} , based at the point \tilde{x}_0 , such that $p \circ \tilde{\gamma} = \gamma$.

Proof.

If $\gamma = p \circ \tilde{\gamma}$ for some loop $\tilde{\gamma}$ in \tilde{X} based at \tilde{x}_0 then $[\gamma] = p_{\#}[\tilde{\gamma}]$, and therefore $[\gamma] \in p_{\#}(\pi_1(\tilde{X}, \tilde{x}_0))$.

Conversely suppose that $[\gamma] \in p_{\#}(\pi_1(\tilde{X}, \tilde{x}_0))$. We must show that there exists some loop $\tilde{\gamma}$ in \tilde{X} based at \tilde{x}_0 such that $\gamma = p \circ \tilde{\gamma}$. Now there exists a loop σ in \tilde{X} based at the point \tilde{x}_0 such that $[\gamma] = p_{\#}([\sigma])$ in $\pi_1(X, p(\tilde{x}_0))$. Then $\gamma \simeq p \circ \sigma$ rel $\{0, 1\}$. It follows from the Path Lifting Theorem for covering maps (Theorem 4.5) that there exists a unique path

$$\tilde{\gamma}$$
: [0, 1] $\rightarrow \tilde{X}$

in \tilde{X} for which $\tilde{\gamma}(0) = \tilde{x}_0$ and $p \circ \tilde{\gamma} = \gamma$.

It then follows from Proposition 6.2 that $\tilde{\gamma}(1) = \sigma(1)$ and $\tilde{\gamma} \simeq \sigma$ rel $\{0, 1\}$. But $\sigma(1) = \tilde{x}_0$. Therefore the path $\tilde{\gamma}$ is the required loop in \tilde{X} based the point \tilde{x}_0 which satisfies $p \circ \tilde{\gamma} = \gamma$.

Corollary 6.5 Let $p: \tilde{X} \to X$ be a covering map over a topological space X, let w_0 and w_1 be points of \tilde{X} satisfying $p(w_0) = p(w_1)$, and let $\alpha: [0,1] \to \tilde{X}$ be a path in \tilde{X} from w_0 to w_1 . Suppose that $[p \circ \alpha] \in p_{\#}(\pi_1(\tilde{X}, w_0))$. Then the path α is a loop in \tilde{X} , and thus $w_0 = w_1$.

Proof.

It follows from Proposition 6.4 that there exists a loop β based at w_0 satisfying $p \circ \beta = p \circ \alpha$. Then $\alpha(0) = \beta(0)$. Now Proposition 4.3 ensures that the lift to \tilde{X} of any path in X is uniquely determined by its starting point. It follows that $\alpha = \beta$. But then the path α must be a loop in \tilde{X} , and therefore $w_0 = w_1$, as required. **Corollary 6.6** Let $p: \tilde{X} \to X$ be a covering map over a topological space X. Let $\alpha: [0, 1] \to X$ and $\beta: [0, 1] \to X$ be paths in X such that

 $\alpha(0) = \beta(0)$ and $\alpha(1) = \beta(1)$, and let $\alpha.\beta^{-1}$ be the loop in X defined such that

$$(\alpha.\beta^{-1})(t) = \begin{cases} \alpha(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \beta(2-2t) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Let $\tilde{\alpha}$: $[0,1] \to \tilde{X}$ and $\tilde{\beta}$: $[0,1] \to \tilde{X}$ be paths in \tilde{X} such that $p \circ \tilde{\alpha} = \alpha$, and $p \circ \tilde{\beta} = \beta$. Suppose that $\tilde{\alpha}(0) = \tilde{\beta}(0)$. Then

$$\tilde{\alpha}(1) = \tilde{\beta}(1)$$

if and only if

$$[\alpha.\beta^{-1}] \in p_{\#}(\pi_1(\tilde{X}, \tilde{x}_0)),$$

where $\tilde{x}_0 = \tilde{\alpha}(0) = \tilde{\beta}(0).$

Proof.

Suppose that $\tilde{\alpha}(1) = \tilde{\beta}(1)$. Then the concatenation $\tilde{\alpha}.\tilde{\beta}^{-1}$ is a loop in \tilde{X} based at \tilde{x}_0 , and $[\alpha.\beta^{-1}] = p_{\#}([\tilde{\alpha}.\tilde{\beta}^{-1}])$, and therefore $[\alpha.\beta^{-1}] \in p_{\#}(\pi_1(\tilde{X},\tilde{x}_0)).$ Conversely suppose that $\tilde{\alpha}$ and $\tilde{\beta}$ are paths in \tilde{X} satisfying $p \circ \tilde{\alpha} = \alpha$, $p \circ \tilde{\beta} = \beta$ and

$$\tilde{\alpha}(0) = \tilde{\beta}(0) = \tilde{x}_0,$$

and that

$$[\alpha.\beta^{-1}] \in p_{\#}(\pi_1(\tilde{X},\tilde{x}_0)).$$

We must show that $\tilde{\alpha}(1) = \tilde{\beta}(1)$. Let

$$\gamma: [0, 1] \to X$$

be the loop based at $p(\tilde{x}_0)$ given by $\gamma = \alpha . \beta^{-1}$. Thus

$$\gamma(t) = \begin{cases} \alpha(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \beta(2-2t) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Then $[\gamma] \in p_{\#}(\pi_1(\tilde{X}, \tilde{x}_0))$. It follows from Proposition 6.4 that there exists a loop $\tilde{\gamma}$ in \tilde{X} based at \tilde{x}_0 such that $p \circ \tilde{\gamma} = \gamma$. Let $\hat{\alpha}: [0,1] \to \tilde{X}$ and Let $\hat{\beta}: [0,1] \to \tilde{X}$ be the paths in \tilde{X} defined such that $\hat{\alpha}(t) = \tilde{\gamma}(\frac{1}{2}t)$ and $\hat{\beta}(t) = \tilde{\gamma}(1 - \frac{1}{2}t)$ for all $t \in [0,1]$. Then

$$\tilde{\alpha}(0) = \hat{\alpha}(0) = \tilde{\beta}(0) = \hat{\beta}(0) = \tilde{x}_0,$$

 $p \circ \hat{\alpha} = \alpha = p \circ \tilde{\alpha}$ and $p \circ \hat{\beta} = \beta = p \circ \tilde{\beta}$.

But Proposition 4.3 ensures that the lift to \tilde{X} of any path in X is uniquely determined by its starting point. Therefore $\tilde{\alpha} = \hat{\alpha}$ and $\tilde{\beta} = \hat{\beta}$. It follows that

$$\tilde{\alpha}(1) = \hat{\alpha}(1) = \tilde{\gamma}(\frac{1}{2}) = \hat{\beta}(1) = \tilde{\beta}(1),$$

as required.

Theorem 6.7 Let $p: \tilde{X} \to X$ be a covering map over a topological space X. Suppose that \tilde{X} is path-connected and that X is simplyconnected. Then the covering map $p: \tilde{X} \to X$ is a homeomorphism.

Proof.

We show that the map $p: \tilde{X} \to X$ is a bijection. This map is surjective (since covering maps are by definition surjective). We must show that it is injective.

Let w_0 and w_1 be points of \tilde{X} with the property that $p(w_0) = p(w_1)$. Then there exists a path α : $[0,1] \to \tilde{X}$ with $\alpha(0) = w_0$ and $\alpha(1) = w_1$, since \tilde{X} is path-connected. Then $p \circ \alpha$ is a loop in X based at the point x_0 , where $x_0 = p(w_0)$. However $\pi_1(X, p(w_0))$ is the trivial group, since X is simply-connected. It follows from Corollary 6.5 that the path α is a loop in \tilde{X} based at w_0 , and therefore $w_0 = w_1$. This shows that the the covering map $p: \tilde{X} \to X$ is injective. Thus the map $p: \tilde{X} \to X$ is a bijection, and thus has a well-defined inverse $p^{-1}: X \to \tilde{X}$. It now follows from Lemma 4.2 that $p: \tilde{X} \to X$ is a homeomorphism, as required.

Let $p: \tilde{X} \to X$ be a covering map over some topological space X, and let x_0 be some chosen basepoint of X. We shall investigate the dependence of the subgroup $p_{\#}(\pi_1(\tilde{X}, \tilde{x}))$ of $\pi_1(X, x_0)$ on the choice of the point \tilde{x} in \tilde{X} , where \tilde{x} is chosen such that $p(\tilde{x}) = x_0$. We first introduce some concepts from group theory.

Let G be a group, and let H be a subgroup of G. Given any $g \in G$, let gHg^{-1} denote the subset of G defined by

 $gHg^{-1} = \{g' \in G : g' = ghg^{-1} \text{ for some } h \in H\}.$ It is easy to verify that gHg^{-1} is a subgroup of G.

Definition.

Let G be a group, and let H and H' be subgroups of G. We say that H and H' are conjugate if and only if there exists some $g \in G$ for which $H' = gHg^{-1}$. Note that if $H' = gHg^{-1}$ then $H = g^{-1}H'g$. The relation of conjugacy is an equivalence relation on the set of all subgroups of the group G. Moreover conjugate subgroups of Gare isomorphic, since the homomorphism sending $h \in H$ to ghg^{-1} is an isomorphism from H to gHg^{-1} whose inverse is the homorphism sending $h' \in gHg^{-1}$ to $g^{-1}h'g$.

A subgroup H of a group G is said to be a normal subgroup of G if $ghg^{-1} \in H$ for all $h \in H$ and $g \in G$. If H is a normal subgroup of G then $gHg^{-1} \subset H$ for all $g \in G$. But then $g^{-1}Hg \subset H$ and $H = g(g^{-1}Hg)g^{-1}$ for all $g \in G$, and therefore $H \subset gHg^{-1}$ for all $g \in G$. It follows from this that a subgroup H of G is a normal subgroup if and only if $gHg^{-1} = H$ for all $g \in G$. Thus a subgroup H of G is a normal subgroup if and only if there is no other subgroup of G conjugate to H. **Lemma 6.8** Let $p: \tilde{X} \to X$ be a covering map over a topological space X. Let x_0 be a point of X, and let w_0 and w_1 be points of \tilde{X} for which

 $p(w_0) = x_0 = p(w_1).$

Let H_0 and H_1 be the subgroups of $\pi_1(X, x_0)$ defined by

 $H_0 = p_{\#}(\pi_1(\tilde{X}, w_0)), \quad H_1 = p_{\#}(\pi_1(\tilde{X}, w_1)).$ Suppose that the covering space \tilde{X} is pathconnected. Then the subgroups H_0 and H_1 of $\pi_1(X, x_0)$ are conjugate. Moreover if H is any subgroup of $\pi_1(X, x_0)$ which is conjugate to H_0 then there exists an element w of \tilde{X} for which p(w) = x and $p_{\#}(\pi_1(\tilde{X}, w)) = H.$

Proof.

Let $\alpha: [0,1] \to \tilde{X}$ be a path in \tilde{X} for which $\alpha(0) = w_0$ and $\alpha(1) = w_1$. (Such a path exists since \tilde{X} is path-connected.) Then each loop σ in \tilde{X} based at w_1 determines a corresponding loop $\alpha.\sigma.\alpha^{-1}$ in \tilde{X} based at w_0 , where

$$(\alpha.\sigma.\alpha^{-1})(t) \equiv \begin{cases} \alpha(3t) & \text{if } 0 \le t \le \frac{1}{3}; \\ \sigma(3t-1) & \text{if } \frac{1}{3} \le t \le \frac{2}{3}; \\ \alpha(3-3t) & \text{if } \frac{2}{3} \le t \le 1. \end{cases}$$

(This loop traverses the path α from w_0 to w_1 , then continues round the loop σ , and traverses the path α in the reverse direction in order to return from w_1 to w_0 .) Let $\eta: [0,1] \to X$ be the loop in X based at the point x_0 given by $\eta = p \circ \alpha$, and let

$$\varphi: \pi_1(X, x_0) \to \pi_1(X, x_0)$$

be the automorphism of the group $\pi_1(X, x_0)$ defined such that $\varphi([\gamma]) = [\eta][\gamma][\eta]^{-1}$ for all loops γ in X based at the point x_0 . Then $p \circ (\alpha.\sigma.\alpha^{-1}) = \eta.(p \circ \sigma).\eta^{-1}$, and there-fore

 $p_{\#}([\alpha.\sigma.\alpha^{-1}]) = [\eta]p_{\#}([\sigma])[\eta]^{-1} = \varphi(p_{\#}([\sigma]))$ in $\pi_1(X, x_0)$. It follows that $\varphi(H_1) \subset H_0$. Similarly $\varphi^{-1}(H_0) \subset H_1$, where

$$\varphi^{-1}([\gamma]) = [\eta]^{-1}[\gamma][\eta]$$

for all loops γ in X based at the point x_0 . It follows that $\varphi(H_1) = H_0$, and thus the subgroups H_0 and H_1 are conjugate

Now let H be a subgroup of $\pi_1(X, x_0)$ which is conjugate to H_0 . Then $H_0 = [\eta]H[\eta]^{-1}$ for some loop η in X based at the point x_0 . It follows from the Path Lifting Theorem for covering maps (Theorem 4.5) that there exists a path α : $[0, 1] \rightarrow \tilde{X}$ in \tilde{X} for which $\alpha(0) = w_0$ and $p \circ \alpha = \eta$. Let $w = \alpha(1)$. Then

$$p_{\#}(\pi_1(\tilde{X}, w)) = [\eta]^{-1} H_0[\eta] = H,$$

as required.

The Brouwer Fixed Point Theorem in Two Dimensions.

Theorem 6.9 Let $f: D \to D$ be a continuous map which maps the closed disk D into itself. Then $f(\mathbf{x}_0) = \mathbf{x}_0$ for some $\mathbf{x}_0 \in D$.

Proof.

Let ∂D denote the boundary circle of D. The inclusion map $i: \partial D \hookrightarrow D$ induces a corresponding homomorphism

 $i_{\#}: \pi_1(\partial D, \mathbf{b}) \to \pi_1(D, \mathbf{b})$

of fundamental groups for any $\mathbf{b} \in \partial D$.

Suppose that it were the case that the map f has no fixed point in D. Then one could define a continuous map $r: D \to \partial D$ as follows: for each $\mathbf{x} \in D$, let $r(\mathbf{x})$ be the point on the boundary ∂D of D obtained by continuing the line segment joining $f(\mathbf{x})$ to \mathbf{x} beyond \mathbf{x} until it intersects ∂D at the point $r(\mathbf{x})$. Note that $r|\partial D$ is the identity map of ∂D .

Let $r_{\#}: \pi_1(D, \mathbf{b}) \to \pi_1(\partial D, \mathbf{b})$ be the homomorphism of fundamental groups induced by $r: D \to \partial D$. Now

 $(r \circ i)_{\#}: \pi_1(\partial D, \mathbf{b}) \to \pi_1(\partial D, \mathbf{b})$

is the identity isomorphism of $\pi_1(\partial D, \mathbf{b})$, since $r \circ i: \partial D \to \partial D$ is the identity map. But it follows directly from the definition of induced homomorphisms that $(r \circ i)_{\#} = r_{\#} \circ i_{\#}$. Therefore $i_{\#}: \pi_1(\partial D, \mathbf{b}) \to \pi_1(D, \mathbf{b})$ is injective, and $r_{\#}: \pi_1(D, \mathbf{b}) \to \pi_1(\partial D, \mathbf{b})$ is surjective. But this is impossible, since $\pi_1(\partial D, \mathbf{b}) \cong$ \mathbb{Z} (Theorem 6.1) and $\pi_1(D, \mathbf{b})$ is the trivial group. This contradiction shows that the continuous map $f: D \to D$ must have at least one fixed point.

204

Free Discontinuous Group Actions on Topological Spaces.

Discontinuous Group Actions.

Definition.

Let G be a group, and let X be a set. The group G is said to *act* on the set X (on the left) if each element g of G determines a corresponding function $\theta_g: X \to X$ from the set X to itself, where

(i)
$$\theta_{gh} = \theta_g \circ \theta_h$$
 for all $g, h \in G$;

(ii) the function θ_e determined by the identity element e of G is the identity function of X. Let G be a group acting on a set X. Given any element x of X, the orbit $[x]_G$ of x (under the group action) is defined to be the subset $\{\theta_q(x) : g \in G\}$ of X, and the sta*bilizer* of x is defined to the subgroup $\{g \in G : \theta_q(x) = x\}$ of the group G. Thus the orbit of an element x of X is the set consisting of all points of X to which x gets mapped under the action of elements of the group G. The stabilizer of x is the subgroup of G consisting of all elements of this group that fix the point x. The group G is said to act freely on X if $\theta_q(x) \neq x$ for all $x \in X$ and $g \in G$ satisfying $g \neq e$. Thus the group G acts freely on X if and only if the stabilizer of every element of X is the trivial subgroup of G.

Let *e* be the identity element of *G*. Then $x = \theta_e(x)$ for all $x \in X$, and therefore $x \in [x]_G$ for all $x \in X$, where $[x]_G = \{\theta_g(x) : g \in G\}$. Let x and y be elements of G for which

$[x]_G \cap [y]_G$

is non-empty, and let $z \in [x]_G \cap [y]_G$. Then there exist elements h and k of G such that

$$z = \theta_h(x) = \theta_k(y).$$

Then

$$\theta_g(z) = \theta_{gh}(x) = \theta_{gk}(y),$$

 $\theta_g(x) = \theta_{gh^{-1}}(z)$

and

$$\theta_g(y) = \theta_{gk^{-1}}(z)$$

for all $g \in G$, and therefore

$$[x]_G = [z]_G = [y]_G.$$

It follows from this that the group action partitions the set X into orbits, so that each element of X determines an orbit which is the unique orbit for the action of G on X to which it belongs. We denote by X/G the set of orbits for the action of G on X.

Now suppose that the group G acts on a topological space X. Then there is a surjective function $q: X \to X/G$, where $q(x) = [x]_G$ for all $x \in X$. This surjective function induces a quotient topology on the set of orbits: a subset U of X/G is open in this quotient topology if and only if $q^{-1}(U)$ is an open set in X (see Lemma 3.16). We define the *orbit* space X/G for the action of G on X to be the topological space whose underlying set is the set of orbits for the action of G on X, the topology on X/G being the quotient topology induced by the function $q: X \to X/G$. This function $q: X \to X/G$ is then an identification map: we shall refer to it as the quotient map from X to X/G.

We shall be concerned here with situations in which a group action on a topological space gives rise to a covering map. The relevant group actions are those where the group acts *freely and properly discontinuously* on the topological space.

Definition.

Let G be a group with identity element e, and let X be a topological space. The group G is said to act freely and properly discontinuously on X if each element g of G determines a corresponding continuous map $\theta_g: X \to X$, where the following conditions are satisfied:

(i)
$$\theta_{gh} = \theta_g \circ \theta_h$$
 for all $g, h \in G$;

- (ii) the continuous map θ_e determined by the identity element e of G is the identity map of X;
- (iii) given any point x of X, there exists an open set U in X such that $x \in U$ and $\theta_g(U) \cap U = \emptyset$ for all $g \in G$ satisfying $g \neq e$.

Let G be a group which acts freely and properly discontinuously on a topological space X. Given any element g of G, the corresponding continuous function $\theta_g: X \to X$ determined by X is a homeomorphism. Indeed it follows from conditions (i) and (ii) in the above definition that $\theta_{g^{-1}} \circ \theta_g$ and $\theta_g \circ \theta_{g^{-1}}$ are both equal to the identity map of X, and therefore $\theta_g: X \to X$ is a homeomorphism with inverse $\theta_{g^{-1}}: X \to X$.

Remark.

The terminology 'freely and properly discontinuously' is traditional, but is hardly ideal. The adverb 'freely' refers to the requirement that $\theta_g(x) \neq x$ for all $x \in X$ and for all $g \in G$ satisfying $g \neq e$. The adverb 'discontinuously' refers to the fact that, given any point x of G, the elements of the orbit

$\{\theta_g(x) : g \in G\}$

of x are separated; it does not signify that the functions defining the action are in any way discontinuous or badly-behaved.

The adverb 'properly' refers to the fact that, given any compact subset K of X, the number of elements of g for which $K \cap \theta_q(K) \neq \emptyset$ is finite. Moreover the definitions of properly discontinuous actions in textbooks and in sources of reference are not always in agreement: some say that an action of a group Gon a topological space X (where each group element determines a corresponding homeomorphism of the topological space) is properly discontinuous if, given any $x \in X$, there exists an open set U in X such that the number of elements g of the group for which $g(U) \cap U \neq \emptyset$ is finite; others say that the action is properly discontinuous if it satisfies the conditions given in the definition above for a group acting freely and properly discontinuously on the set. William Fulton, in his textbook Algebraic topology: a first course (Springer, 1995), introduced the term 'evenly' in place of 'freely and properly discontinuously', but this change in terminology does not appear to have been generally adopted.

Proposition 7.1 Let G be a group acting freely and properly discontinuously on a topological space X. Then the quotient map $q: X \rightarrow X/G$ from X to the corresponding orbit space X/G is a covering map.

Proof.

The quotient map $q: X \to X/G$ is surjective. Let V be an open set in X. Then $q^{-1}(q(V))$ is the union $\bigcup_{g \in G} \theta_g(V)$ of the open sets $\theta_g(V)$ as g ranges over the group G, since $q^{-1}(q(V))$ is the subset of X consisting of all elements of X that belong to the orbit of some element of V. But any union of open sets in a topological space is an open set. We conclude therefore that if V is an open set in Xthen q(V) is an open set in X/G. Let x be a point of X. Then there exists an open set U in X such that $x \in U$ and $\theta_g(U) \cap U = \emptyset$ for all $g \in G$ satisfying $g \neq e$. Now

$$q^{-1}(q(U)) = \bigcup_{g \in G} \theta_g(U).$$

We claim that the sets $\theta_g(U)$ are disjoint. Let g and h be elements of G. Suppose that

$$\theta_g(U) \cap \theta_h(U) \neq \emptyset.$$

Then

$$\theta_{h^{-1}}(\theta_g(U) \cap \theta_h(U)) \neq \emptyset.$$

But $\theta_{h-1}: X \to X$ is a bijection, and therefore

$$\begin{aligned} \theta_{h^{-1}}(\theta_g(U) \cap \theta_h(U)) \\ &= \theta_{h^{-1}}(\theta_g(U)) \cap \theta_{h^{-1}}(\theta_h(U)) \\ &= \theta_{h^{-1}g}(U) \cap U, \end{aligned}$$

and therefore $\theta_{h^{-1}g}(U) \cap U \neq \emptyset$. It follows that $h^{-1}g = e$, where e denotes the identity element of G, and therefore g = h. Thus if gand h are elements of g, and if $g \neq h$, then

$$\theta_g(U) \cap \theta_h(U) = \emptyset.$$

We conclude therefore that the preimage

$$q^{-1}(q(U))$$

of q(U) is the disjoint union of the sets $\theta_g(U)$ as g ranges over the group G. Moreover each these sets $\theta_g(U)$ is an open set in X.

Now $U \cap [u]_G = \{u\}$ for all $u \in U$, since

 $[u]_G = \{\theta_g(u) : g \in G\}$

and $U \cap \theta_g(U) = \emptyset$ when $g \neq e$. Thus if u and v are elements of U, and if q(u) = q(v) then $[u]_G = [v]_G$ and therefore u = v. It follows that the restriction $q|U: U \to X/G$ of the quotient map q to U is injective, and therefore q maps U bijectively onto q(U). But q maps open sets onto open sets, and any continuous bijection that maps open sets onto open sets onto open sets is a homeomorphism. We conclude therefore that the restriction of $q: X \to X/G$ to the open set U maps U homeomorphically onto q(U). Moreover, given any element g of G, the quotient map q satisfies $q = q \circ \theta_{g^{-1}}$, and the homeomorphism $\theta_{g^{-1}}$ maps $\theta_g(U)$ homeomorphically onto U.

It follows that the quotient map q maps $\theta_g(U)$ homeomorphically onto q(U) for all $g \in U$. We conclude therefore that q(U) is an evenly covered open set in X/G whose preimage $q^{-1}(q(U))$ is the disjoint union of the open sets $\theta_g(U)$ as g ranges over the group G. It follows that the quotient map $q: X \to X/G$ is a covering map, as required. **Theorem 7.2** Let G be a group acting freely and properly discontinuously on a path-connected topological space X, let $q: X \to X/G$ be the quotient map from X to the orbit space X/G, and let x_0 be a point of X. Then there exists a surjective homomorphism

 λ : $\pi_1(X/G, q(x_0)) \to G$

with the property that

 $\tilde{\gamma}(1) = \theta_{\lambda([\gamma])}(x_0)$

for any loop γ in X/G based at $q(x_0)$, where $\tilde{\gamma}$ denotes the unique path in X for which $\tilde{\gamma}(0) = x_0$ and $q \circ \tilde{\gamma} = \gamma$. The kernel of this homomorphism is the subgroup

$$q_{\#}(\pi_1(X,x_0))$$

of $\pi_1(X/G, q(x_0))$.
Proof.

Let $\gamma: [0,1] \to X/G$ be a loop in the orbit space with $\gamma(0) = \gamma(1) = q(x_0)$. It follows from the Path Lifting Theorem for covering maps (Theorem 4.5) that there exists a unique path $\tilde{\gamma}: [0,1] \to X$ for which $\tilde{\gamma}(0) = x_0$ and $q \circ \tilde{\gamma} = \gamma$. Now $\tilde{\gamma}(0)$ and $\tilde{\gamma}(1)$ must belong to the same orbit, since

$$q(\tilde{\gamma}(0)) = \gamma(0) = \gamma(1) = q(\tilde{\gamma}(1)).$$

Therefore there exists some element g of Gsuch that $\tilde{\gamma}(1) = \theta_g(x_0)$. This element g is uniquely determined, since the group G acts freely on X. Moreover the value of g is determined by the based homotopy class $[\gamma]$ of γ in $\pi_1(X/G, q(x_0))$. Indeed it follows from Proposition 6.2 that if σ is a loop in X/Gbased at $q(x_0)$, if $\tilde{\sigma}$ is the lift of σ starting at x_0 (so that $q \circ \tilde{\sigma} = \sigma$ and $\tilde{\sigma}(0) = x_0$), and if $[\gamma] = [\sigma]$ in $\pi_1(X/G, q(x_0))$ (so that $\gamma \simeq \sigma$ rel $\{0, 1\}$), then $\tilde{\gamma}(1) = \tilde{\sigma}(1)$. We conclude therefore that there exists a well-defined function

$$\lambda: \pi_1(X/G, q(x_0)) \to G,$$

which is characterized by the property that $\tilde{\gamma}(1) = \theta_{\lambda([\gamma])}(x_0)$ for any loop γ in X/G based at $q(x_0)$, where $\tilde{\gamma}$ denotes the unique path in X for which $\tilde{\gamma}(0) = x_0$ and $q \circ \tilde{\gamma} = \gamma$.

Now let $\alpha: [0,1] \to X/G$ and $\beta: [0,1] \to X/G$ be loops in X/G based at x_0 , and let $\tilde{\alpha}: [0,1] \to X$ and $\tilde{\beta}: [0,1] \to X$ be the lifts of α and β respectively starting at x_0 , so that $q \circ \tilde{\alpha} = \alpha$, $q \circ \tilde{\beta} = \beta$ and $\tilde{\alpha}(0) = \tilde{\beta}(0) = x_0$. Then $\tilde{\alpha}(1) = \theta_{\lambda([\alpha])}(x_0)$ and $\tilde{\beta}(1) = \theta_{\lambda([\beta])}(x_0)$. Then the path $\theta_{\lambda([\alpha])} \circ \tilde{\beta}$ is also a lift of the loop β , and is the unique lift of β starting at $\tilde{\alpha}(1)$. Let $\alpha.\beta$ be the concatenation of the loops α and $\beta,$ where

$$(\alpha.\beta)(t) = \begin{cases} \alpha(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \beta(2t-1) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

Then the unique lift of α . β to X starting at x_0 is the path σ : $[0, 1] \rightarrow X$, where

$$\sigma(t) = \begin{cases} \tilde{\alpha}(2t) & \text{if } 0 \le t \le \frac{1}{2}; \\ \theta_{\lambda([\alpha])}(\tilde{\beta}(2t-1)) & \text{if } \frac{1}{2} \le t \le 1. \end{cases}$$

It follows that

$$\begin{aligned} \theta_{\lambda([\alpha][\beta])}(x_{0}) &= \theta_{\lambda([\alpha,\beta])}(x_{0}) \\ &= \sigma(1) \\ &= \theta_{\lambda([\alpha])}(\tilde{\beta}(1)) \\ &= \theta_{\lambda([\alpha])}(\theta_{\lambda([\beta])}(x_{0})) \\ &= \theta_{\lambda([\alpha])\lambda([\beta])}(x_{0}) \end{aligned}$$

and therefore

$$\lambda([\alpha][\beta]) = \lambda([\alpha])\lambda([\beta]).$$

Therefore the function

$$\lambda: \pi_1(X/G, q(x_0)) \to G$$

is a homomorphism.

Let $g \in G$. Then there exists a path α in X from x_0 to $\theta_g(x_0)$, since the space X is pathconnected. Then $q \circ \alpha$ is a loop in X/G based at $q(x_0)$, and $g = \lambda([q \circ \alpha])$. This shows that the homomorphism λ is surjective.

Let $\gamma: [0, 1] \to X/G$ be a loop in X/G based at $q(x_0)$. Suppose that $[\gamma] \in \ker \lambda$. Then

$$\tilde{\gamma}(1) = \theta_e(x_0) = x_0,$$

and therefore $\tilde{\gamma}$ is a loop in X based at x_0 . Moreover $[\gamma] = q_{\#}[\tilde{\gamma}]$, and therefore

$$[\gamma] \in q_{\#}(\pi_1(X, x_0)).$$

On the other hand, if

$$[\gamma] \in q_{\#}(\pi_1(X, x_0))$$

then $\gamma = q \circ \tilde{\gamma}$ for some loop $\tilde{\gamma}$ in X based at x_0 (see Proposition 6.4). But then

$$x_0 = \tilde{\gamma}(1) = \theta_{\lambda([\gamma])}(x_0),$$

and therefore $\lambda([\gamma]) = e$, where e is the identity element of G. Thus

$$\ker \lambda = q_{\#}(\pi_1(X, x_0)),$$

as required.

Corollary 7.3 Let G be a group acting freely and properly discontinuously on a path-connected topological space X, let $q: X \to X/G$ be the quotient map from X to the orbit space X/G, and let x_0 be a point of X. Then $q_{\#}(\pi_1(X, x_0))$ is a normal subgroup of the fundamental group $\pi_1(X/G, q(x_0))$ of the orbit space, and

$$\frac{\pi_1(X/G, q(x_0))}{q_{\#}(\pi_1(X, x_0))} \cong G.$$

Proof.

The subgroup $q_{\#}(\pi_1(X, x_0))$ is the kernel of the homomorphism

$$\lambda: \pi_1(X/G, q(x_0)) \to G$$

described in the statement of Theorem 7.2. It is therefore a normal subgroup of

$$\pi_1(X/G,q(x_0)),$$

since the kernel of any homomorphism is a normal subgroup.

The homomorphism λ is surjective, and the image of any group homomorphism is isomorphism of the quotient of its domain by its kernel. The result follows.

Corollary 7.4 Let G be a group acting freely and properly discontinuously on a simply-connected topological space X, let $q: X \to X/G$ be the quotient map from X to the orbit space X/G, and let x_0 be a point of X. Then

 $\pi_1(X/G, q(x_0)) \cong G.$

Proof.

This is a special case of Corollary 7.3.

Example.

The group \mathbb{Z} of integers under addition acts freely and properly discontinuously on the real line \mathbb{R} . Indeed each integer n determines a corresponding homeomorphism $\theta_n : \mathbb{R} \to \mathbb{R}$, where

$$\theta_n(x) = x + n$$

for all $x \in \mathbb{R}$. Moreover

$$\theta_m \circ \theta_n = \theta_{m+n}$$

for all $m, n \in \mathbb{Z}$, and θ_0 is the identity map of \mathbb{R} . If $U = (-\frac{1}{2}, \frac{1}{2})$ then $\theta_n(U) \cap U = \emptyset$ for all non-zero integers n. The real line \mathbb{R} is simply-connected. It follows from Corollary 7.4 that

 $\pi_1(\mathbb{R}/\mathbb{Z},b)\cong\mathbb{Z}$

for any point b of \mathbb{R}/\mathbb{Z} .

Now the orbit space \mathbb{R}/\mathbb{Z} is homeomorphic to a circle. Indeed let

$$q:\mathbb{R}\to\mathbb{R}/\mathbb{Z}$$

be the quotient map. Then the surjective function $p: \mathbb{R} \to S^1$ which sends $t \in \mathbb{R}$ to

 $(\cos 2\pi t, \sin 2\pi t)$

induces a continuous map

$$h: \mathbb{R}/\mathbb{Z} \to S^1$$

defined on the orbit space which satisfies

$$h \circ q = p,$$

since the quotient map q is an identification map. Moreover real numbers t_1 and t_2 satisfy $p(t_1) = p(t_2)$ if and only if $q(t_1) = q(t_2)$. It follows that the induced map

$$h: \mathbb{R}/\mathbb{Z} \to S^1$$

is a bijection. This map also maps open sets to open sets, for if W is any open set in the orbit space \mathbb{R}/\mathbb{Z} then $q^{-1}(W)$ is an open set in \mathbb{R} , and therefore $p(q^{-1}(W))$ is an open set in S^1 , since the covering map $p: \mathbb{R} \to S^1$ maps open sets to open sets (Lemma 4.2). But

$$p(q^{-1}(W)) = h(W)$$

for all open sets W in \mathbb{R}/\mathbb{Z} . Thus the continuous bijection

$h: \mathbb{R}/\mathbb{Z} \to S^1$

maps open sets to open sets, and is therefore a homeomorphism. It follows from Corollary 7.4 that $\pi_1(S^1, b) \cong \mathbb{Z}$ for any point *b* of the circle S^1 . This shows that Theorem 6.1 concerning the fundamental group of the circle can be obtained as a special case of the more general result Corollary 7.4 concerning fundamental groups of orbit spaces obtained via discontinuous group actions on simply connected topological spaces.

Example.

The group \mathbb{Z}^n of ordered *n*-tuples of integers under addition acts freely and properly discontinuously on \mathbb{R}^n , where

$$\theta_{(m_1,m_2,\dots,m_n)}(x_1,x_2,\dots,x_n) \\ = (x_1 + m_1, x_2 + m_2,\dots,x_n + m_n)$$

for all

$$(m_1, m_2, \ldots, m_n) \in \mathbb{Z}^n$$

and

$$(x_1, x_2, \ldots, x_n) \in \mathbb{R}^n.$$

The orbit space $\mathbb{R}^n/\mathbb{Z}^n$ is an *n*-dimensional torus, homeomorphic to the product of *n* circles. It follows from Corollary 7.4 that the fundamental group of this *n*-dimensional torus is isomorphic to the group \mathbb{Z}^n .

Example.

Let C_2 be the cyclic group of order 2. Then $C_2 = \{e, a\}$ where e is the identity element, $a \neq e$, $a^2 = e$. Then the group C_2 acts freely and properly discontinuously on the ndimensional sphere S^n for each non-negative integer n. We represent S^n as the unit sphere centred on the origin in \mathbb{R}^{n+1} . The homeomorphism θ_e determined by the identity element e of C_2 is the identity map of S^n ; the homeomorphism θ_a determined by the element a of C_2 is the antipodal map that sends each point x of S^n to -x. The orbit space S^n/C_2 is homeomorphic to real projective ndimensional space $\mathbb{R}P^n$. The *n*-dimensional sphere is simply-connected if n > 1. It follows from Corollary 7.4 that the fundamental group of $\mathbb{R}P^n$ is isomorphic to the cyclic group C_2 when n > 1.

Note that S^0 is a pair of points, and $\mathbb{R}P^0$ is a single point. Also S^1 is a circle (which is not simply-connected) and $\mathbb{R}P^1$ is homeomorphic to a circle. Moreover, for any $b \in S^1$, the homomorphism

$$q_{\#}:\pi_1(S^1,b)\to\pi_1(\mathbb{R}P^1,q(b))$$

corresponds to the homomorphism from \mathbb{Z} to \mathbb{Z} that sends each integer n to 2n. This is consistent with the conclusions of Corollary 7.3 in this example.

Example.

Given a pair (m, n) of integers, let

$$\theta_{m,n}: \mathbb{R}^2 \to \mathbb{R}^2$$

be the homeomorphism of the plane \mathbb{R}^2 defined such that

$$\theta_{m,n}(x,y) = (x+m,(-1)^m y + n)$$

for all $(x,y) \in \mathbb{R}^2$. Let (m_1, n_1) and (m_2, n_2) be ordered pairs of integers. Then

$$\theta_{m_1,n_1} \circ \theta_{m_2,n_2} = \theta_{m_1+m_2,n_1+(-1)^{m_1}n_2}.$$

Let Γ be the group whose elements are represented as ordered pairs of integers, where the group operation # on Γ is defined such that

 $(m_1, n_1) \# (m_2, n_2) = (m_1 + m_2, n_1 + (-1)^{m_1} n_2)$ for all $(m_1, n_1), (m_2, n_2) \in \Gamma$. The group Γ is non-Abelian, and its identity element is (0,0). This group acts on the plane \mathbb{R}^2 : given $(m,n) \in \Gamma$ the corresponding symmetry $\theta_{m,n}$ is a translation if m is even, and is a glide reflection if m is odd. Given a pair (m, n) of integers, the corresponding homeomorphism $\theta_{m,n}$ maps an open disk about the point (x, y) onto an open disk of the same radius about the point $\theta_{(m,n)}(x,y)$. It follows that if D is the open disk of radius $\frac{1}{2}$ about the point (x, y), and if $D \cap \theta_{m,n}(D)$ is non-empty, then (m,n) = (0,0). Thus the group Γ maps freely and properly discontinuously on the plane \mathbb{R}^2 .

Now each orbit intersects the closed unit square S, where $S = [0,1] \times [0,1]$. If 0 < x < 1and 0 < y < 1 then the orbit of (x, y) intersects the square S in one point, namely the point (x, y). If 0 < x < 1, then the orbit of (x, 0) intersects the square in two points (x,0) and (x,1). If 0 < y < 1 then the orbit of (0, y) intersects the square S in the two points (0, y) and (1, 1 - y). (Note that $(1, 1 - y) = \theta_{1,1}(0, y)$.) And the orbit of any corner of the square S intersects the square in the four corners of the square. The restriction q|S of the quotient map $q: \mathbb{R}^2 \to \mathbb{R}^2/\Gamma$ to the square S is a continuous surjection defined on the square: one can readily verify that it is an identification map. It follows that the orbit space \mathbb{R}^2/Γ is homeomorphic to the identification space obtained from the closed square S by identifying together the points (x,0) and (x,1) where the real number x satisfies 0 < x < 1, identifying together the points (0, y) and (1, 1-y) where the real number y satisfies 0 < y < 1, and identifying together the four corners of the square.

The identification space obtained in this fashion is a closed non-orientable surface, first described by Felix Klein in 1882, and now known as the *Klein bottle*. Apparently the surface was initially referred to as the *Kleinsche Fläche* (Klein's Surface), but this name was incorrectly translated into English, and, as a result the surface is now referred to as the Klein Bottle (*Kleinsche Flasche*).

The plane \mathbb{R}^2 is simply-connected. It follows from Corollary 7.4 that the fundamental group of the Klein bottle is isomorphic to the group Γ defined above.