

Module MA22200: Advanced Analysis
(Semester 2)
Hilary Term 2020
Part IV (Section 9)

D. R. Wilkins

Copyright © David R. Wilkins 2015–2020

Contents

9	Modes of Convergence on Measure Spaces	89
9.1	Egorov's Theorem	89
9.2	Almost Uniform Convergence	91
9.3	Convergence in Measure	92
9.4	Sequences of Functions that are Cauchy in Measure	93
9.5	Convergence in Mean	97
9.6	Hölder's Inequality	100
9.7	Minkowski's Inequality	102
9.8	Convergence in L^p norm	103
9.9	The L^p Spaces	106

9 Modes of Convergence on Measure Spaces

9.1 Egorov's Theorem

Proposition 9.1 *Let (X, \mathcal{A}, μ) be a measure space, and let E be a measurable subset of X , where $\mu(E) < +\infty$. Let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions that converges on E to a measurable function f . Then, given any strictly positive real numbers ε and δ , there exists a measurable subset F of E and a positive integer N such that $\mu(F) < \delta$ and $|f_j(x) - f(x)| < \varepsilon$ whenever $x \in E \setminus F$ and $j \geq N$.*

Proof Let strictly positive real numbers ε and δ be given and, for each positive integer k , let

$$E_k = \bigcap_{j=k}^{+\infty} \{x \in X : |f_j(x) - f(x)| < \varepsilon\}.$$

Let $x \in E$. Then $\lim_{j \rightarrow +\infty} f_j(x) = f(x)$, and therefore there exists some positive integer k such that $|f_j(x) - f(x)| < \varepsilon$ whenever $j \geq k$. But then $x \in E_k$. It follows from this that $E = \bigcup_{k=1}^{+\infty} E_k$.

Now $E_j \subset E_{j+1}$ for all positive integers j . It follows from the countable additivity of the measure μ that $\mu(E) = \lim_{k \rightarrow +\infty} \mu(E_k)$ (see Lemma 7.25). Now $\mu(E) < +\infty$. It follows that there exists some positive integer N that is large enough to ensure that $\mu(E_N) > \mu(E) - \delta$. Let $F = E \setminus E_N$. Then $\mu(F) < \delta$ and $|f(x) - f_j(x)| < \varepsilon$ whenever $x \in E \setminus F$ and $j \geq N$, as required. ■

Let X be a set, and let E be a subset of X . Let f_1, f_2, f_3, \dots be an infinite sequence of real-valued functions on X , and let f be a real-valued function on X . The sequence f_1, f_2, f_3, \dots is said to converge *uniformly* on E to the limit function f if, given any strictly positive real number ε , there exists some positive integer N (independent of the choice of x in the set E) such that $|f_j(x) - f(x)| < \varepsilon$ whenever $x \in E$ and $j \geq N$.

Sequences of functions that converge pointwise do not necessarily converge uniformly. An infinite sequence f_1, f_2, f_3, \dots of real-valued functions on the set X is said to converge *pointwise* to the real-valued function f on a subset E of X if, given any strictly positive real number ε , and given any point x of E , there exists some positive integer $N(x)$ (in general dependent on the choice of both ε and x , such set E) such that $|f_j(x) - f(x)| < \varepsilon$ whenever $j \geq N(x)$.

Now let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f be a measurable real-valued function on f . If there exists a subset F of X satisfying $\mu(F) = 0$, and if the infinite sequence f_1, f_2, f_3, \dots converges pointwise to f on the complement $X \setminus F$ of F in X , then the sequence f_1, f_2, f_3, \dots is said to converge *pointwise almost everywhere* on X .

Theorem 9.2 (Egorov's Theorem) *Let (X, \mathcal{A}, μ) be a measure space, and let E be a measurable subset of X , where $\mu(E) < +\infty$. Let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions that converges pointwise almost everywhere on E to a measurable function f . Then, given any strictly positive real number δ , there exists a subset F of E such that $\mu(F) < \delta$ and the sequence f_1, f_2, f_3, \dots of real-valued functions converges uniformly to the limit function f on $E \setminus F$.*

Proof If the infinite sequence f_1, f_2, f_3, \dots does not converge to f throughout the set E , then there exists some subset E_0 of E such that $\mu(E \setminus E_0) = 0$ and the sequence f_1, f_2, f_3, \dots converges to the limit function f at all points of E_0 , because the sequence f_1, f_2, f_3, \dots does at least converge to f almost everywhere on E . If then, for a given strictly positive real number δ , we show the existence of a subset F_0 of E_0 such that $\mu(F_0) < \delta$ and the sequence f_1, f_2, f_3, \dots converges uniformly to f on $E_0 \setminus F_0$, and if we take $F = F_0 \cup (E \setminus E_0)$, then $\mu(F) < \delta$ and the sequence f_1, f_2, f_3, \dots of functions converges uniformly to the limit function f on $E \setminus F$. Thus we may assume, without loss of generality, that the sequence of functions f_1, f_2, f_3, \dots converges to the limit function throughout the set E .

Thus suppose that the sequence f_1, f_2, f_3, \dots converges pointwise to the limit function f throughout E , and let some strictly positive real number δ be given. It follows on applying Proposition 9.1, that there exist positive integers N_1, N_2, N_3, \dots and measurable subsets F_1, F_2, F_3, \dots of E such that $\mu(F_k) < 2^{-k}\delta$ and $|f_j(x) - f(x)| < 1/k$ whenever $x \in E \setminus F_k$ and $j \geq N_k$.

Let $F = \bigcup_{k=1}^{+\infty} F_k$. Then

$$\mu(F) \leq \sum_{k=1}^{+\infty} \mu(F_k) < \delta.$$

The required result is thus established once we prove that the infinite sequence f_1, f_2, f_3, \dots of measurable functions converges uniformly to the limit function f on the set $E \setminus F$.

Let some strictly positive real number ε be given. Then some positive integer k can be chosen large enough to ensure that $1/k < \varepsilon$. Let $N = N_k$.

Now $E \setminus F \subset E \setminus F_k$. It follows that $|f_j(x) - f(x)| < \varepsilon$ whenever $x \in E \setminus F$ and $j \geq N$. Thus the functions f_1, f_2, f_3, \dots do indeed converge uniformly to the limit function f on $E \setminus F$. The result follows. ■

9.2 Almost Uniform Convergence

Definition Let (X, \mathcal{A}, μ) be a measure space, let E be a measurable subset of X , let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f be a measurable real-valued function on X . The infinite sequence f_1, f_2, f_3, \dots is said to converge *almost uniformly* on E if, given any positive real number δ , there exists a measurable subset F of E such that $\mu(F) < \delta$ and the infinite sequence f_1, f_2, f_3, \dots converges uniformly to the limit function f on $E \setminus F$.

Egorov's Theorem (Theorem 9.2) ensures that if (X, \mathcal{A}, μ) is a measure space, and if E is a measurable subset of X whose measure $\mu(E)$ is finite, then any infinite sequence of measurable real-valued functions that converges pointwise almost everywhere on E also converges almost uniformly on E .

Example For each positive integer j , let $f_j: \mathbb{R} \rightarrow \mathbb{R}$ be defined so that

$$f_j(x) = \begin{cases} 1 & \text{if } j-1 < x \leq j; \\ 0 & \text{otherwise.} \end{cases}$$

Then the infinite sequence f_1, f_2, f_3, \dots converges pointwise to the zero function throughout the real line \mathbb{R} but does not converge almost uniformly with respect to Lebesgue measure on the real line.

Lemma 9.3 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f be a measurable real-valued function on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions converges almost uniformly to the limit function f . Then the sequence f_1, f_2, f_3, \dots converges to f pointwise almost everywhere.*

Proof Let F be the subset of X consisting of those points x for which the infinite sequence $f_1(x), f_2(x), f_3(x), \dots$ does not converge to $f(x)$. We must show that $\mu(F) = 0$.

Now the infinite sequence f_1, f_2, f_3, \dots converges almost uniformly on X to the limit function f , and therefore, given any strictly positive real number δ , there exists some measurable set F_δ such that $\mu(F_\delta) < \delta$ and $\lim_{j \rightarrow +\infty} f_j(x) = f(x)$ for all $x \in X \setminus F_\delta$. But then $F \subset F_\delta$, and consequently $\mu(F) \leq \mu(F_\delta) < \delta$. We conclude therefore that $\mu(F) < \delta$ for all positive real numbers δ . It follows that $\mu(F) = 0$, as required. ■

9.3 Convergence in Measure

Definition Let (X, \mathcal{A}, μ) be a measure space, and let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f be a measurable real-valued function on X . The sequence f_1, f_2, f_3, \dots is said to *converge in measure* to the function f if, given any strictly positive real numbers ε and δ , there exists some positive integer N such that

$$\mu(\{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}) < \delta$$

whenever $j \geq N$.

The following result follows immediately from the relevant definitions.

Lemma 9.4 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f be a measurable real-valued function on X . Then the infinite sequence f_1, f_2, f_3, \dots converges in measure to the limit function f if and only if, given any strictly positive real number ε ,*

$$\lim_{j \rightarrow +\infty} \mu(\{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}) = 0.$$

Proposition 9.5 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f and \tilde{f} be measurable real-valued functions on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots converges in measure both to the function f and also to the function \tilde{f} . Then the functions f and \tilde{f} are equal almost everywhere.*

Proof For each positive integer j and positive real number ε , let

$$\begin{aligned} E_{j,\varepsilon} &= \{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}, \\ \tilde{E}_{j,\varepsilon} &= \{x \in X : |f_j(x) - \tilde{f}(x)| \geq \varepsilon\}. \end{aligned}$$

Then

$$\lim_{j \rightarrow +\infty} \mu(E_{j,\varepsilon}) \rightarrow 0 \quad \text{and} \quad \lim_{j \rightarrow +\infty} \mu(\tilde{E}_{j,\varepsilon}) \rightarrow 0.$$

Now if x is a point of X , j is a positive integer, ε is a positive real number, and if $|f(x) - \tilde{f}(x)| \geq 2\varepsilon$, then either $|f_j(x) - f(x)| \geq \varepsilon$ or else $|f_j(x) - \tilde{f}(x)| \geq \varepsilon$. It follows that

$$\{x \in X : |f(x) - \tilde{f}(x)| \geq 2\varepsilon\} \subset E_{j,\varepsilon} \cup \tilde{E}_{j,\varepsilon}$$

for all positive integers j , and therefore

$$\mu\left(\{x \in X : |f(x) - \tilde{f}(x)| \geq 2\varepsilon\}\right) \leq \mu(E_{j,\varepsilon}) + \mu(\tilde{E}_{j,\varepsilon})$$

for all positive integers j . Taking limits as $j \rightarrow +\infty$, we conclude that

$$\mu\left(\{x \in X : |f(x) - \tilde{f}(x)| \geq 2\varepsilon\}\right) = 0$$

for all positive real numbers ε . Now

$$\{x \in X : f(x) \neq \tilde{f}(x)\} = \bigcup_{k=1}^{+\infty} \{x \in X : |f(x) - \tilde{f}(x)| \geq 2/k\}.$$

It follows that $\{x \in X : f(x) \neq \tilde{f}(x)\}$ is expressible as a countable union of subsets of X that are each of measure zero, and thus must itself be of measure zero. The result follows. ■

Proposition 9.6 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , and let f be a measurable real-valued function on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots converges almost uniformly on X to the limit function f . Then the sequence f_1, f_2, f_3, \dots converges in measure on X to the limit function f .*

Proof Let strictly positive real numbers ε and δ be given. There then exists a measurable subset F of X such that $\mu(F) < \delta$ and f_1, f_2, f_3, \dots converges uniformly to f on $X \setminus F$. There then exists some positive integer N such that $|f_j(x) - f(x)| < \varepsilon$ whenever $x \in X \setminus F$ and $j \geq N$. But then

$$\{x \in X : |f_j(x) - f(x)| \geq \varepsilon\} \subset F$$

whenever $j \geq N$. But $\mu(F) < \delta$. It follows that

$$\mu(\{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}) < \delta$$

whenever $j \geq N$. Thus f_1, f_2, f_3, \dots converges in measure to the limit function f , as required. ■

9.4 Sequences of Functions that are Cauchy in Measure

Definition Let (X, \mathcal{A}, μ) be a measure space, and let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X . The sequence

f_1, f_2, f_3, \dots is said to be *Cauchy in measure* if, given any positive real numbers ε and δ , there exists some positive integer N such that

$$\mu(\{x \in X : |f_j(x) - f_k(x)| \geq \varepsilon\}) < \delta$$

whenever $j \geq N$ and $k \geq N$.

Lemma 9.7 *Let (X, \mathcal{A}, μ) be a measure space, and let f_1, f_2, f_3, \dots be a sequence of measurable real-valued functions on X that converges in measure to some limit function. Then the infinite sequence f_1, f_2, f_3, \dots of functions is Cauchy in measure.*

Proof For all positive integers j and k and positive real numbers ε , let

$$E_{j,\varepsilon} = \{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}$$

and

$$G_{j,k,\varepsilon} = \{x \in X : |f_j(x) - f_k(x)| \geq \varepsilon\}.$$

Let strictly positive real numbers ε and δ be given. Then there exists some positive integer N such that $\mu(E_{j,\frac{1}{2}\varepsilon}) < \frac{1}{2}\delta$ whenever $j \geq N$, because the sequence f_1, f_2, f_3, \dots converges in measure to the function f . Now if $x \in X$, if j and k are positive integers, and if $|f_j(x) - f_k(x)| \geq \varepsilon$ then either $|f_j(x) - f(x)| \geq \frac{1}{2}\varepsilon$ or else $|f_k(x) - f(x)| \geq \frac{1}{2}\varepsilon$. It follows that

$$G_{j,k,\varepsilon} \subset E_{j,\frac{1}{2}\varepsilon} \cup E_{k,\frac{1}{2}\varepsilon}$$

for all positive integers j and k , and therefore

$$\mu(G_{j,k,\varepsilon}) \leq \mu(E_{j,\frac{1}{2}\varepsilon}) + \mu(E_{k,\frac{1}{2}\varepsilon}) < \frac{1}{2}\delta + \frac{1}{2}\delta = \delta$$

whenever $j \geq N$ and $k \geq N$. The result follows. ■

Proposition 9.8 *Let (X, \mathcal{A}, μ) be a measure space, and let f_1, f_2, f_3, \dots be a sequence of measurable real-valued functions on X that is Cauchy in measure. Then the sequence f_1, f_2, f_3, \dots has a subsequence that converges almost uniformly on X .*

Proof The sequence f_1, f_2, f_3, \dots is Cauchy in measure, and therefore there exists an infinite sequence j_1, j_2, j_3, \dots of positive integers, where

$$j_1 < j_2 < j_3 < \dots,$$

such that, for each positive integer k ,

$$\mu\left(\left\{x \in X : |f_r(x) - f_s(x)| \geq \frac{1}{2^k}\right\}\right) < \frac{1}{2^k}$$

whenever $r \geq j_k$ and $s \geq j_k$. It then follows in particular that $\mu(E_k) < 2^{-k}$ for all positive integers k , where

$$E_k = \left\{x \in X : |f_{j_k}(x) - f_{j_{k+1}}(x)| \geq \frac{1}{2^k}\right\}.$$

Let $F_k = \bigcup_{p=k}^{+\infty} E_p$ for all positive integers k . Then

$$\mu(F_k) \leq \sum_{p=k}^{+\infty} \mu(E_p) \leq \sum_{p=k}^{+\infty} \frac{1}{2^p} = \frac{1}{2^{k-1}}.$$

for all positive integers k .

Let k be a positive integer, and let $x \in X \setminus F_k$. Then $x \notin E_p$ for all integers p satisfying $p \geq k$, and therefore $|f_{j_p}(x) - f_{j_{p+1}}(x)| < 2^{-p}$ for all $p \geq k$. Thus if p and q are integers satisfying $k \leq p < q$ then

$$|f_{j_p}(x) - f_{j_q}(x)| < \sum_{r=p}^{q-1} \frac{1}{2^r} < \frac{1}{2^{p-1}}.$$

It follows that, given any positive real number ε , there exists some positive integer M such that $|f_{j_p}(x) - f_{j_q}(x)| < \varepsilon$ for all integers p and q satisfying $M \leq p < q$. Thus $f_1(x), f_2(x), f_3(x), \dots$ is a Cauchy sequence of real numbers. The completeness of the real number system ensures that every Cauchy sequence of real numbers is convergent. It follows that $f(x)$ is well-defined for all $x \in X \setminus F_k$, where $f(x) = \lim_{p \rightarrow +\infty} f_{j_p}(x)$.

Now we showed that $|f_{j_p}(x) - f_{j_q}(x)| < 2^{1-p}$ whenever $x \notin F_k$ and $k \leq p < q$. Taking limits as $q \rightarrow +\infty$, it follows that $|f_{j_p}(x) - f(x)| \leq 2^{1-p}$ whenever $p \geq k$. Therefore, given any positive real number ε , there exists a positive integer M_k , independent of the choice of x , such that $|f_{j_p}(x) - f(x)| < \varepsilon$ for all $x \in X \setminus F_k$ and for all integers p satisfying $p \geq M_k$. The sequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ of real-valued functions therefore converges uniformly on $X \setminus F_k$ to the limit function f .

Now $f(x)$ is well-defined for all positive integers k and for all points x of $X \setminus F_k$. It is therefore well-defined for all points x of $X \setminus F_\infty$, where $F_\infty = \bigcap_{k=1}^{+\infty} F_k$. Now F_∞ is a measurable set, and $\mu(F_\infty) \leq \mu(F_k) \leq 2^{1-k}$ for

all positive integers. It follows that $\mu(F_\infty) = 0$. Thus the limit function f is defined almost everywhere on X . Also, given any positive real number δ , the positive integer k can be chosen large enough to ensure that $\mu(F_k) < \delta$. The sequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ then converges uniformly to the limit function f on $X \setminus F_k$. The sequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ is therefore almost uniformly convergent on X . Moreover it is a subsequence of the given sequence f_1, f_2, f_3, \dots . The result follows. ■

Proposition 9.9 *Let (X, \mathcal{A}, μ) be a measure space, and let f_1, f_2, f_3, \dots be a sequence of measurable real-valued functions on X that is Cauchy in measure. Then the sequence f_1, f_2, f_3, \dots converges in measure to some measurable real-valued function f on X .*

Proof The sequence f_1, f_2, f_3, \dots of functions is Cauchy in measure and therefore has a subsequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ that converges almost uniformly on X to some measurable real-valued function f (Proposition 9.8). We show that the sequence f_1, f_2, f_3, \dots converges in measure to this limit function f .

Let strictly positive real numbers ε and δ be given. The subsequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ converges in measure on X to the limit function f , because it converges almost uniformly to f on X (Proposition 9.6). Therefore there exists some positive integer M such that

$$\mu(\{x \in X : |f_{j_k}(x) - f(x)| \geq \tfrac{1}{2}\varepsilon\}) < \tfrac{1}{2}\delta$$

whenever $k \geq M$. Also there exists some positive integer N such that

$$\mu(\{x \in X : |f_p(x) - f_q(x)| \geq \tfrac{1}{2}\varepsilon\}) < \tfrac{1}{2}\delta$$

whenever $p \geq N$ and $q \geq N$, because the infinite sequence f_1, f_2, f_3, \dots is Cauchy in measure on X . Choose a positive integer k large enough to ensure that $k \geq M$ and $j_k \geq N$, and let $g = f_{j_k}$. Also let

$$F_j = \{x \in X : |f_j(x) - g(x)| \geq \tfrac{1}{2}\varepsilon\}$$

for all positive integers j , and let

$$G = \{x \in X : |g(x) - f(x)| \geq \tfrac{1}{2}\varepsilon\}.$$

Then $\mu(F_j) < \tfrac{1}{2}\delta$ whenever $j \geq N$. Also $\mu(G) < \tfrac{1}{2}\delta$. Now if $x \in X$, and if $|f_j(x) - g(x)| < \tfrac{1}{2}\varepsilon$ and $|g(x) - f(x)| < \tfrac{1}{2}\varepsilon$ then

$$|f_j(x) - f(x)| \leq |f_j(x) - g(x)| + |g(x) - f(x)| < \varepsilon.$$

Thus

$$\{x \in X : |f_j(x) - f(x)| \geq \varepsilon\} \subset F_j \cup G$$

for all positive integers j . It follows that if $j \geq N$ then

$$\mu(\{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}) \leq \mu(F_j \cup G) \leq \mu(F_j) + \mu(G) < \delta.$$

Thus the infinite sequence f_1, f_2, f_3, \dots of measurable real-valued functions converges in measure to the limit function f , as required. ■

9.5 Convergence in Mean

Definition Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j| d\mu < +\infty$ for all positive integers j , and let f be a measurable real-valued function on X . We say that the infinite sequence f_1, f_2, f_3, \dots of functions *converges in mean* to the function f if, given any strictly positive real number ε , there exists some positive integer N such that

$$\int_X |f_j - f| d\mu < \varepsilon$$

whenever $j \geq N$.

Definition Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j| d\mu < +\infty$ for all positive integers j . We say that the infinite sequence f_1, f_2, f_3, \dots of functions is *Cauchy in mean* if given any strictly positive real number ε , there exists some positive integer N such that

$$\int_X |f_j - f_k| d\mu < \varepsilon$$

whenever $j \geq N$ and $k \geq N$.

Proposition 9.10 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j| d\mu < +\infty$ for all positive integers j , and let f be a measurable real-valued function on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is Cauchy in mean and also converges almost everywhere on X to the limit function f . Then the infinite sequence f_1, f_2, f_3, \dots of functions on X converges in mean to the function f .*

Proof Let some strictly positive real number ε be given, and let ε_0 satisfy $0 < \varepsilon_0 < \varepsilon$. The infinite sequence f_1, f_2, f_3, \dots of functions is Cauchy in mean, hence there exists some positive integer N such that $\int_X |f_j - f_k| d\mu < \varepsilon_0$ whenever $j \geq N$ and $k \geq N$. Taking limits as $k \rightarrow +\infty$, and applying Fatou's Lemma (Lemma 8.25), we find that

$$\begin{aligned} \int_X |f_j - f| d\mu &= \int_X \left(\lim_{k \rightarrow +\infty} |f_j - f_k| \right) d\mu \\ &\leq \liminf_{k \rightarrow +\infty} \int_X |f_j - f_k| d\mu \leq \varepsilon_0 < \varepsilon \end{aligned}$$

whenever $j \geq N$. Therefore $\lim_{j \rightarrow +\infty} \int_X |f_j - f| d\mu = 0$, and thus f_j converges to f in mean, as required. ■

Lemma 9.11 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j| d\mu < +\infty$ for all positive integers j , and let f be a measurable real-valued function on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is converges in mean to the limit function f . Then this infinite sequence of functions also converges in measure to the function f .*

Proof Let

$$E_{j,\varepsilon} = \{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}$$

and let $\chi_{j,\varepsilon}$ denote the characteristic function of the set $E_{j,\varepsilon}$ for all positive integers j and for all positive real numbers ε . Then $|f_j(x) - f(x)| \geq \varepsilon \chi_{j,\varepsilon}(x)$ for all $x \in X$, positive integers j and positive real numbers ε , and therefore

$$\int_X |f_j - f| dx \geq \varepsilon \mu(E_{j,\varepsilon})$$

for all positive integers j and for all positive real number ε .

Now let positive real numbers ε and δ be given. The sequence f_1, f_2, f_3, \dots is Cauchy in mean. It follows that there exists some positive integer N such that $\int_X |f_j - f| d\mu < \varepsilon \delta$ whenever $j \geq N$. Then $\mu(E_{j,\varepsilon}) < \delta$ whenever $j \geq N$. The result follows. ■

Lemma 9.12 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j| d\mu < +\infty$ for all positive integers j . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is Cauchy in mean. Then this infinite sequence of functions is also Cauchy in measure.*

Proof Let

$$E_{j,k,\varepsilon} = \{x \in X : |f_j(x) - f_k(x)| \geq \varepsilon\}$$

and let $\chi_{j,k,\varepsilon}$ denote the characteristic function of the set $E_{j,k,\varepsilon}$ for all positive integers j and k and for all positive real numbers ε . Then $|f_j(x) - f_k(x)| \geq \varepsilon \chi_{j,k,\varepsilon}(x)$ for all $x \in X$, positive integers j and k and positive real numbers ε , and therefore

$$\int_X |f_j - f_k| d\mu \geq \varepsilon \mu(E_{j,k,\varepsilon})$$

for all positive integers j and k and for all positive real number ε .

Now let positive real numbers ε and δ be given. The sequence f_1, f_2, f_3, \dots is Cauchy in mean. It follows that there exists some positive integer N such that $\int_X |f_j - f_k| d\mu < \varepsilon \delta$ whenever $j \geq N$ and $k \geq N$. Then $\mu(E_{j,k,\varepsilon}) < \delta$ whenever $j \geq N$ and $k \geq N$. The result follows. ■

Proposition 9.13 *Let (X, \mathcal{A}, μ) be a measure space, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j| d\mu < +\infty$ for all positive integers j . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is Cauchy in mean. Then this infinite sequence of functions converges in mean to some measurable real-valued function f for which $\int_X |f| d\mu < +\infty$.*

Proof The infinite sequence f_1, f_2, f_3, \dots of functions is Cauchy in mean. It is therefore Cauchy in measure (Lemma 9.12). It therefore has a subsequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ that converges almost uniformly on X to some measurable real-valued function f on X (Proposition 9.8). This subsequence converges pointwise almost everywhere on X to the limit function f , (Lemma 9.3), and therefore converges in mean to the function f (Proposition 9.10). A positive integer k can then be chosen large enough to ensure that $\int_X |f_{j_k} - f| d\mu \leq 1$. It then follows that

$$\int_X |f| d\mu \leq \int_X |f_{j_k}| d\mu + \int_X |f_{j_k} - f| d\mu \leq \int_X |f_{j_k}| d\mu + 1 < +\infty.$$

To complete the proof we show that the original sequence f_1, f_2, f_3, \dots converges in mean to the limit function f .

Let some strictly positive real number ε be given. Then there exist positive integers M and N that are large enough to ensure that $\int_X |f_{j_k} - f| d\mu < \frac{1}{2}\varepsilon$ whenever $k \geq M$ and $\int_X |f_s - f_t| d\mu < \frac{1}{2}\varepsilon$ whenever $s \geq N$ and $t \geq N$. Let some positive integer k be chosen large enough to ensure that $k \geq M$ and $j_k \geq N$. Then

$$\int_X |f_n - f| d\mu \leq \int_X |f_n - f_{j_k}| d\mu + \int_X |f_{j_k} - f| d\mu < \varepsilon$$

whenever $n \geq N$. Thus $\lim_{n \rightarrow +\infty} \int_X |f_n - f| d\mu = 0$. The result follows. ■

9.6 Hölder's Inequality

Lemma 9.14 (Young's Inequality) *Let a and b be non-negative real numbers, and let p and q be real numbers for which $p > 1$, $q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$. Then*

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

Proof First we show that $e^{(1-t)u+tv} \leq (1-t)e^u + te^v$ for all real numbers u, v and t satisfying $0 \leq t \leq 1$. This is a consequence of the fact that the derivative of the exponential function is increasing. The result clearly holds when $u = v$, and moreover the form of the inequality is preserved on swapping u and v and replacing t by $1-t$. It therefore suffices to verify that the above inequality holds in the case when $u < v$.

Suppose then that $u < v$, and let

$$f(t) = (1-t)e^u + te^v - e^{(1-t)u+tv}.$$

for all real numbers t . Then $f(0) = f(1) = 0$ and

$$f'(t) = e^v - e^u - (v-u)e^{(1-t)u+tv}.$$

From this expression we see that $f'(t)$ is a decreasing function of t . It must therefore be the case that $f'(0) > 0$ and $f'(1) < 0$. This is then a real number t_0 satisfying $0 < t_0 < 1$ for which $f'(t_0) = 0$. The function f is then an increasing function of t on the interval $[0, t_0]$ and a decreasing function of t on the interval $[t_0, 1]$. But $f(0) = f(1) = 0$. It follows that $f(t) > 0$ whenever $0 < t < 1$, and thus $e^{(1-t)u+tv} < (1-t)e^u + te^v$ for all real numbers u, v and t satisfying $u < v$ and $0 < t < 1$. This completes the verification that $e^{(1-t)u+tv} \leq (1-t)e^u + te^v$ in all cases where u, v and t are real numbers and $0 \leq t \leq 1$.

Now let a and b be positive real numbers, and let p and q be real numbers for which $p > 1$, $q > 1$ and $p^{-1} + q^{-1} = 1$. Let $u = p \log a$, $v = q \log b$ and $t = 1/q$. Then $1-t = 1/p$, and therefore

$$e^{(1-t)u+tv} = e^{\log a + \log b} = e^{\log a} e^{\log b} = ab.$$

Also

$$(1-t)e^u = \frac{1}{p}e^{p \log a} = \frac{a^p}{p} \quad \text{and} \quad te^v = \frac{1}{q}e^{q \log b} = \frac{b^q}{q}.$$

The inequality previously established therefore ensures that

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}$$

for all positive real numbers a and b . This inequality also holds when $a \geq 0$, $b \geq 0$ and either $a = 0$ or $b = 0$. The result follows. ■

Alternative Proof Let p and q be real numbers satisfying $p > 1$, $q > 1$ and $p^{-1} + q^{-1} = 1$. Then $q - 1 = 1/(p - 1)$. It follows that if x and y are positive real numbers then $y = x^{p-1}$ if and only if $x = y^{q-1}$. Let a and b be positive real numbers. Then the rectangle in \mathbb{R}^2 with vertices $(0, 0)$, $(a, 0)$, (a, b) and $(0, b)$ is contained in the union of the regions A and B , where

$$\begin{aligned} A &= \{(x, y) \in \mathbb{R}^2 : 0 \leq x \leq a \text{ and } 0 \leq y \leq x^{p-1}\}, \\ B &= \{(x, y) \in \mathbb{R}^2 : 0 \leq y \leq b \text{ and } 0 \leq x \leq y^{q-1}\}. \end{aligned}$$

It follows that

$$ab \leq \int_0^a x^{p-1} dx + \int_0^b y^{q-1} dy = \frac{a^p}{p} + \frac{b^q}{q},$$

as required. ■

Proposition 9.15 (Hölder's Inequality) *Let (X, \mathcal{A}, μ) be a measure space, let p and q be real numbers satisfying the conditions $p > 1$, $q > 1$ and $p^{-1} + q^{-1} = 1$, and let f and g be measurable real-valued functions on X . Suppose that the functions $|f|^p$ and $|g|^q$ are integrable. Then the function fg is integrable and*

$$\int_X |fg| d\mu \leq \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}} \left(\int_X |g|^q d\mu \right)^{\frac{1}{q}}.$$

Proof Let

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}} \quad \text{and} \quad \|g\|_q = \left(\int_X |g|^q d\mu \right)^{\frac{1}{q}}.$$

The identity $ab \leq p^{-1}a^p + q^{-1}b^q$ holds for all non-negative real numbers a and b . (Lemma 9.14). Setting $a = |f(x)|/\|f\|_p$ and $b = |g(x)|/\|g\|_q$, and integrating over the space X we find that

$$\begin{aligned} \frac{1}{\|f\|_p \|g\|_q} \int_X |fg| d\mu &\leq \frac{1}{p \|f\|_p^p} \int_X |f|^p d\mu + \frac{1}{q \|g\|_q^q} \int_X |g|^q d\mu \\ &= \frac{1}{p} + \frac{1}{q} = 1. \end{aligned}$$

Thus $\int_X |fg| d\mu \leq \|f\|_p \|g\|_q$, as required. ■

9.7 Minkowski's Inequality

Proposition 9.16 (Minkowski's Inequality) *Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying the conditions $p \geq 1$, and let f and g be measurable real-valued functions on X . Suppose that the functions $|f|^p$ and $|g|^p$ are integrable. Then the function $|f + g|^p$ is integrable and*

$$\left(\int_X |f + g|^p d\mu \right)^{\frac{1}{p}} \leq \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}} + \left(\int_X |g|^p d\mu \right)^{\frac{1}{p}}.$$

Proof The result in the case $p = 1$ follows immediately on integrating the inequality $|f + g| \leq |f| + |g|$ over the measure space X . It remains therefore to prove the result in the case $p > 1$. There then exists some positive real number q , such that $q > 1$ and $p^{-1} + q^{-1} = 1$. Then $1/q = (p - 1)/p$, and thus $p = (p - 1)q$. Let

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}}, \quad \|g\|_p = \left(\int_X |g|^p d\mu \right)^{\frac{1}{p}}$$

and

$$\|f + g\|_p = \left(\int_X |f + g|^p d\mu \right)^{\frac{1}{p}}.$$

We may assume moreover that $\|f + g\|_p > 0$, since the required inequality is trivially satisfied in cases where $\|f + g\|_p = 0$. Now, applying Hölder's Inequality (Proposition 9.15), we find that

$$\begin{aligned} \|f + g\|_p^p &= \int_X |f + g|^p d\mu \\ &\leq \int_X |f + g|^{p-1} (|f| + |g|) d\mu \\ &\leq \left(\int_X |f + g|^{(p-1)q} d\mu \right)^{\frac{1}{q}} (\|f\|_p + \|g\|_p) \\ &= \left(\int_X |f + g|^p d\mu \right)^{1 - \frac{1}{p}} (\|f\|_p + \|g\|_p) \\ &= \|f + g\|_p^{p-1} (\|f\|_p + \|g\|_p) \end{aligned}$$

Thus $\|f + g\|_p^p \leq \|f + g\|_p^{p-1} (\|f\|_p + \|g\|_p)$. Dividing both sides of the inequality by $\|f + g\|_p^{p-1}$, we find that $\|f + g\|_p \leq \|f\|_p + \|g\|_p$, as required. ■

9.8 Convergence in L^p norm

Definition Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j , and let f be a measurable real-valued function on X . We say that the infinite sequence f_1, f_2, f_3, \dots of functions *converges in L^p norm* to the function f if, given any strictly positive real number ε , there exists some positive integer N such that

$$\int_X |f_j - f|^p d\mu < \varepsilon^p$$

whenever $j \geq N$.

Given a measure space (X, \mathcal{A}, μ) , and given a real number p satisfying $p \geq 1$, it is convenient to define

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}}$$

for all measurable real-valued functions f on X for which $\int_X |f|^p d\mu < +\infty$. With this notation, we can say that if f_1, f_2, f_3, \dots is an infinite sequence of measurable real-valued functions on X , where $\int_X |f|^p d\mu < +\infty$, and if f is a measurable real-valued function on X , then the infinite sequence f_1, f_2, f_3 of functions converges to the limit function f in L^p norm if and only if $\lim_{j \rightarrow +\infty} \|f_j - f\|_p = 0$.

Definition Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j . We say that the infinite sequence f_1, f_2, f_3, \dots of functions is *Cauchy in L^p norm* if given any strictly positive real number ε , there exists some positive integer N such that

$$\int_X |f_j - f_k|^p d\mu < \varepsilon^p$$

whenever $j \geq N$ and $k \geq N$.

Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j . The infinite sequence f_1, f_2, f_3, \dots of functions on X is then *Cauchy in L^p norm* if, given any positive real number ε , there exists some positive integer N such that $\|f_j - f_k\|_p < \varepsilon$ whenever $j \geq N$ and $k \geq N$.

An infinite sequence of measurable real-valued functions on a measure space converges in L^1 norm if and only if it converges in mean. Similarly an infinite sequence of measurable real-valued functions on a measure space is Cauchy in L^1 norm if and only if it is Cauchy in mean. The following results, and their proofs, accordingly generalize those previously stated and proved for sequences of functions that converge in mean or are Cauchy in mean.

Proposition 9.17 *Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j , and let f be a measurable real-valued function on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is Cauchy in L^p norm and also converges almost everywhere on X to the limit function f . Then the infinite sequence f_1, f_2, f_3, \dots of functions on X converges in L^p norm to the function f .*

Proof Let some strictly positive real number ε be given, and let ε_0 satisfy $0 < \varepsilon_0 < \varepsilon$. The infinite sequence f_1, f_2, f_3, \dots of functions is Cauchy in L^p norm, hence there exists some positive integer N such that $\|f_j - f_k\|_p < \varepsilon_0$ whenever $j \geq N$ and $k \geq N$. Taking limits as $k \rightarrow +\infty$, and applying Fatou's Lemma (Lemma 8.25), we find that

$$\begin{aligned} \int_X |f_j - f|^p d\mu &= \int_X \left(\lim_{k \rightarrow +\infty} |f_j - f_k|^p \right) d\mu \\ &\leq \liminf_{k \rightarrow +\infty} \int_X |f_j - f_k|^p d\mu \leq \varepsilon_0^p < \varepsilon^p \end{aligned}$$

whenever $j \geq N$. Therefore $\lim_{j \rightarrow +\infty} \|f_j - f\|_p = 0$, and thus f_j converges to f in L^p norm, as required. \blacksquare

Lemma 9.18 *Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j , and let f be a measurable real-valued function on X . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X converges in L^p norm to the limit function f . Then this infinite sequence of functions also converges in measure to the function f .*

Proof Let

$$E_{j,\varepsilon} = \{x \in X : |f_j(x) - f(x)| \geq \varepsilon\}$$

and let $\chi_{j,\varepsilon}$ denote the characteristic function of the set $E_{j,\varepsilon}$ for all positive integers j and for all positive real numbers ε . Then $|f_j(x) - f(x)|^p \geq \varepsilon^p \chi_{j,\varepsilon}(x)$ for all $x \in X$, positive integers j and positive real numbers ε , and therefore

$$\int_X |f_j - f|^p dx \geq \varepsilon^p \mu(E_{j,\varepsilon})$$

for all positive integers j and for all positive real number ε .

Now let positive real numbers ε and δ be given. The sequence f_1, f_2, f_3, \dots is Cauchy in L^p norm. It follows that there exists some positive integer N such that $\|f_j - f\|_p < \varepsilon \delta^{\frac{1}{p}}$ whenever $j \geq N$. Then $\mu(E_{j,\varepsilon}) < \delta$ whenever $j \geq N$. The result follows. ■

Lemma 9.19 *Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is Cauchy in L^p norm. Then this infinite sequence of functions is also Cauchy in measure.*

Proof Let

$$E_{j,k,\varepsilon} = \{x \in X : |f_j(x) - f_k(x)| \geq \varepsilon\}$$

and let $\chi_{j,k,\varepsilon}$ denote the characteristic function of the set $E_{j,k,\varepsilon}$ for all positive integers j and k and for all positive real numbers ε . Then $|f_j(x) - f_k(x)|^p \geq \varepsilon^p \chi_{j,k,\varepsilon}(x)$ for all $x \in X$, positive integers j and k and positive real numbers ε , and therefore

$$\int_X |f_j - f_k|^p dx \geq \varepsilon^p \mu(E_{j,k,\varepsilon})$$

for all positive integers j and k and for all positive real number ε .

Now let positive real numbers ε and δ be given. The sequence f_1, f_2, f_3, \dots is Cauchy in L^p norm. It follows that there exists some positive integer N such that $\|f_j - f_k\|_p < \varepsilon \delta^{\frac{1}{p}}$ whenever $j \geq N$ and $k \geq N$. Then $\mu(E_{j,k,\varepsilon}) < \delta$ whenever $j \geq N$ and $k \geq N$. The result follows. ■

Proposition 9.20 *Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, let f_1, f_2, f_3, \dots be an infinite sequence of measurable real-valued functions on X , where $\int_X |f_j|^p d\mu < +\infty$ for all positive integers j . Suppose that the infinite sequence f_1, f_2, f_3, \dots of functions on X is Cauchy in L^p norm. Then this infinite sequence of functions converges in L^p norm to some measurable real-valued function f for which $\int_X |f|^p d\mu < +\infty$.*

Proof The infinite sequence f_1, f_2, f_3, \dots of functions is Cauchy in L^p norm. It is therefore Cauchy in measure (Lemma 9.19). It therefore has a subsequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ that converges almost uniformly on X to some measurable real-valued function f on X (Proposition 9.8). The subsequence $f_{j_1}, f_{j_2}, f_{j_3}, \dots$ then converges pointwise almost everywhere on X to the limit function f (Lemma 9.3), and therefore converges in L^p norm to the function f (Proposition 9.17). A positive integer k can then be chosen large enough to ensure that $\|f_{j_k} - f\|_p \leq 1$. It then follows from Minkowski's inequality (Proposition 9.16) that

$$\left(\int_X |f|^p d\mu \right)^{\frac{1}{p}} \leq \|f_{j_k}\|_p + \|f_{j_k} - f\|_p \leq \|f_{j_k}\|_p + 1 < +\infty.$$

To complete the proof we show that the original sequence f_1, f_2, f_3, \dots converges in L^p norm to the limit function f .

Let some strictly positive real number ε be given. Then there exist positive integers M and N that are large enough to ensure that $\|f_{j_k} - f\|_p < \frac{1}{2}\varepsilon$ whenever $k \geq M$ and $\|f_s - f_t\|_p < \frac{1}{2}\varepsilon$ whenever $s \geq N$ and $t \geq N$. Let some positive integer k be chosen large enough to ensure that $k \geq M$ and $j_k \geq N$. Then

$$\|f_n - f\|_p \leq \|f_n - f_{j_k}\|_p + \|f_{j_k} - f\|_p < \varepsilon$$

whenever $n \geq N$. Thus $\lim_{n \rightarrow +\infty} \|f_n - f\|_p = 0$. The result follows. ■

9.9 The L^p Spaces

Let (X, \mathcal{A}, μ) be a measure space, let p be a real number satisfying $p \geq 1$, and let $\mathcal{L}^p(X, \mu)$ denote the collection consisting of all measurable real-valued functions f on X with the property that $\int_X |f|^p d\mu < +\infty$. It follows from Minkowski's Inequality (Proposition 9.16) that the sum of two real-valued functions belonging to $\mathcal{L}^p(X, \mu)$ itself belongs to $\mathcal{L}^p(X, \mu)$, and thus $\mathcal{L}^p(X, \mu)$ is a real vector space.

Let

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}}$$

for all $f \in \mathcal{L}^p(X, \mu)$. Then $\|f\|_p \geq 0$ and $\|cf\|_p = |c| \|f\|_p$ for all $f \in \mathcal{L}^p(X, \mu)$ and for all real numbers c . Also Minkowski's Inequality (Proposition 9.16) ensures that $\|f + g\|_p \leq \|f\|_p + \|g\|_p$ for all $f, g \in \mathcal{L}^p(X, \mu)$.

However it is not the case that $\|f\|_p = 0$ implies that $f = 0$. In fact $\|f\|_p = 0$ if and only if the set $\{x \in X : f(x) \neq 0\}$ has measure zero, so that the function f is equal to zero almost everywhere on X .

Now it is an easy exercise to verify that the relation of being equal almost everywhere is an equivalence relation on the set of functions that belong to $\mathcal{L}^p(X, \mu)$. This relation therefore partitions the set $\mathcal{L}^p(X, \mu)$ into equivalence classes. Let f be a measurable real-valued function on X for which $|f|^p$ is integrable on X . The equivalence class $[f]_p$ of f then consists of those measurable real-valued functions on X that are equal to the function f almost everywhere on X .

Now if f, g, \tilde{f} and \tilde{g} are measurable functions on X , if c is a real number, if f and \tilde{f} are equal almost everywhere on X , and if g and \tilde{g} are equal almost everywhere on X , then cf and $c\tilde{f}$ are equal almost everywhere on X , and $f + g$ and $\tilde{f} + \tilde{g}$ are equal almost everywhere on X . We may therefore add together equivalence classes of functions belonging to $\mathcal{L}^p(X, \mu)$ and multiply them by real scalars so that $[f]_p + [g]_p = [f + g]_p$ and $[cf]_p = c[f]_p$ for all $f, g \in \mathcal{L}^p(X, \mu)$ and for all real numbers c . Also $\|f\|_p = \|\tilde{f}\|_p$ whenever the measurable real-valued functions f and \tilde{f} are equal almost everywhere on X and belong to the vector space $\mathcal{L}^p(X, \mu)$. We may therefore define $\|[f]_p\|_p = \|f\|_p$ for all $f \in \mathcal{L}^p(X, \mu)$.

Now let $L^p(X, \mu)$ denote the set of all equivalence classes $[f]_p$ of measurable real-valued functions on X that belong to $\mathcal{L}^p(X, \mu)$, where two such functions on X belong to the same equivalence class if and only if they are equal almost everywhere on X . Denoting the equivalence class of any member f of $\mathcal{L}^p(X, \mu)$ by $[f]_p$, we note that there are well-defined operations of addition and multiplication-by-scalars on $L^p(X, \mu)$, where

$$[f]_p + [g]_p = [f + g]_p, \quad c[f]_p = [cf]_p$$

for all $f, g \in \mathcal{L}^p(X, \mu)$ and for all real numbers c . Also setting $\|[f]_p\|_p = \|f\|_p$ for all $f \in \mathcal{L}^p(X, \mu)$, where

$$\|f\|_p = \left(\int_X |f|^p d\mu \right)^{\frac{1}{p}},$$

we find that $\|[f]_p\|_p \geq 0$, $\|c[f]_p\|_p = |c| \|[f]_p\|_p$ and $\|[f]_p + [g]_p\|_p \leq \|[f]_p\|_p + \|[g]_p\|_p$ for all $f, g \in \mathcal{L}^p(X, \mu)$ and for all real numbers c . Also $\|[f]_p\|_p = 0$ if and only if $[f]_p = [0]_p$. It follows that the function that maps the equivalence class $[f]_p$ of each member f of $\mathcal{L}^p(X, \mu)$ to the non-negative real number $\|f\|_p$ is a norm on the vector space $L^p(X, \mu)$. We obtain in this fashion a normed vector space $L^p(X, \mu)$ whose elements are equivalence classes of measurable real-valued functions f on X . A measurable real-valued function f on X determines a corresponding element of $L^p(X, \mu)$ if and only if $\int_X |f|^p d\mu < +\infty$. Two such functions determine the same element of $L^p(X, \mu)$ if and only if they are equal almost everywhere on X with respect to the measure μ .

Proposition 9.20 ensures that the normed vector space $L^p(X, \mu)$ is complete for all real numbers p satisfying $p \geq 1$. The space $L^p(X, \mu)$ is thus a *Banach space*. (Banach spaces are complete normed vector spaces.) This result is of fundamental importance in many branches of mathematics developed from the early twentieth century onwards.