

Module MAU34804: Fixed Point Theorems  
and Economic Equilibria  
Hilary Term 2022  
Part I (Sections 1 to 2)

D. R. Wilkins

Copyright © David R. Wilkins 2016–2022

## Contents

<b>1</b>	<b>Review of Basic Results of Analysis in Euclidean Spaces</b>	<b>1</b>
1.1	Basic Properties of Vectors and Norms . . . . .	1
1.2	The Bolzano-Weierstrass Theorem . . . . .	2
1.3	The Multidimensional Extreme Value Theorem . . . . .	6
1.4	The Glueing Lemma . . . . .	6
1.5	Lebesgue Numbers . . . . .	7
<b>2</b>	<b>Correspondences and Hemicontinuity</b>	<b>11</b>
2.1	Correspondences . . . . .	11
2.2	The Graph of a Correspondence . . . . .	15
2.3	Compact-Valued Upper Hemicontinuous Correspondences . . .	22
2.4	A Criterion characterizing Lower Hemicontinuity . . . . .	28
2.5	Intersections of Correspondences . . . . .	30
2.6	Berge's Maximum Theorem . . . . .	31

# 1 Review of Basic Results of Analysis in Euclidean Spaces

## 1.1 Basic Properties of Vectors and Norms

We denote by  $\mathbb{R}^n$  the set consisting of all  $n$ -tuples  $(x_1, x_2, \dots, x_n)$  of real numbers. The set  $\mathbb{R}^n$  represents  $n$ -dimensional *Euclidean space* (with respect to the standard Cartesian coordinate system). Let  $\mathbf{x}$  and  $\mathbf{y}$  be elements of  $\mathbb{R}^n$ , where

$$\mathbf{x} = (x_1, x_2, \dots, x_n), \quad \mathbf{y} = (y_1, y_2, \dots, y_n),$$

and let  $\lambda$  be a real number. We define

$$\begin{aligned} \mathbf{x} + \mathbf{y} &= (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n), \\ \mathbf{x} - \mathbf{y} &= (x_1 - y_1, x_2 - y_2, \dots, x_n - y_n), \\ \lambda \mathbf{x} &= (\lambda x_1, \lambda x_2, \dots, \lambda x_n), \\ \mathbf{x} \cdot \mathbf{y} &= x_1 y_1 + x_2 y_2 + \dots + x_n y_n, \\ |\mathbf{x}| &= \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}. \end{aligned}$$

The quantity  $\mathbf{x} \cdot \mathbf{y}$  is the *scalar product* (or *inner product*) of  $\mathbf{x}$  and  $\mathbf{y}$ , and the quantity  $|\mathbf{x}|$  is the *Euclidean norm* of  $\mathbf{x}$ . Note that  $|\mathbf{x}|^2 = \mathbf{x} \cdot \mathbf{x}$ . The *Euclidean distance* between two points  $\mathbf{x}$  and  $\mathbf{y}$  of  $\mathbb{R}^n$  is defined to be the Euclidean norm  $|\mathbf{y} - \mathbf{x}|$  of the vector  $\mathbf{y} - \mathbf{x}$ .

Let  $\mathbf{x}$  and  $\mathbf{y}$  be elements in  $\mathbb{R}^n$ , Let  $p(t) = |t\mathbf{x} + \mathbf{y}|^2$  for all real numbers  $t$ . Then

$$\begin{aligned} p(t) &= (t\mathbf{x} + \mathbf{y}) \cdot (t\mathbf{x} + \mathbf{y}) \\ &= t^2 |\mathbf{x}|^2 + 2t \mathbf{x} \cdot \mathbf{y} + |\mathbf{y}|^2 \end{aligned}$$

for all real numbers  $t$ . But  $p(t) \geq 0$  for all real numbers  $t$ . It follows that  $|\mathbf{x} \cdot \mathbf{y}| \leq |\mathbf{x}| |\mathbf{y}|$ . This inequality is known as *Schwarz's Inequality*.

Moreover, given any elements  $\mathbf{x}$  and  $\mathbf{y}$  of  $\mathbb{R}^n$ ,

$$\begin{aligned} |\mathbf{x} + \mathbf{y}|^2 &= (\mathbf{x} + \mathbf{y}) \cdot (\mathbf{x} + \mathbf{y}) = |\mathbf{x}|^2 + |\mathbf{y}|^2 + 2\mathbf{x} \cdot \mathbf{y} \\ &\leq |\mathbf{x}|^2 + |\mathbf{y}|^2 + 2|\mathbf{x}| |\mathbf{y}| = (|\mathbf{x}| + |\mathbf{y}|)^2. \end{aligned}$$

It follows that  $|\mathbf{x} + \mathbf{y}| \leq |\mathbf{x}| + |\mathbf{y}|$ . It follows from this inequality that

$$|\mathbf{x} - \mathbf{z}| \leq |\mathbf{x} - \mathbf{y}| + |\mathbf{y} - \mathbf{z}|$$

for all  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$ . This identity is known as the *Triangle Inequality*. It expresses the geometric result that the length of any side of a triangle in a Euclidean space of any dimension is the sum of the lengths of the other two sides of that triangle.

**Definition** A sequence  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  of points in  $\mathbb{R}^n$  is said to *converge* to a point  $\mathbf{p}$  if and only if the following criterion is satisfied:—

given any real number  $\varepsilon$  satisfying  $\varepsilon > 0$  there exists some positive integer  $N$  such that  $|\mathbf{x}_j - \mathbf{p}| < \varepsilon$  whenever  $j \geq N$ .

We refer to  $\mathbf{p}$  as the *limit*  $\lim_{j \rightarrow +\infty} \mathbf{x}_j$  of the sequence  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$ .

**Lemma 1.1** *Let  $\mathbf{p}$  be a point of  $\mathbb{R}^n$ , where  $\mathbf{p} = (p_1, p_2, \dots, p_n)$ . Then a sequence  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  of points in  $\mathbb{R}^n$  converges to  $\mathbf{p}$  if and only if the  $i$ th components of the elements of this sequence converge to  $p_i$  for  $i = 1, 2, \dots, n$ .*

A proof of Lemma 1.1 is to be found in Appendix A.

## 1.2 The Bolzano-Weierstrass Theorem

**Definition** Let  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  be an infinite sequence of points in  $n$ -dimensional Euclidean space  $\mathbb{R}^n$ . A *subsequence* of this infinite sequence is a sequence of the form  $\mathbf{x}_{j_1}, \mathbf{x}_{j_2}, \mathbf{x}_{j_3}, \dots$  where  $j_1, j_2, j_3, \dots$  is an infinite sequence of positive integers with

$$j_1 < j_2 < j_3 < \dots.$$

### Theorem 1.2 (Multidimensional Bolzano-Weierstrass Theorem)

*Every bounded sequence of points in a Euclidean space has a convergent subsequence.*

A proof of Theorem 1.2 is to be found in Appendix A.

**Definition** Let  $X$  be a subset of  $\mathbb{R}^n$ . Given a point  $\mathbf{p}$  of  $X$  and a non-negative real number  $r$ , the *open ball*  $B_X(\mathbf{p}, r)$  in  $X$  of *radius*  $r$  about  $\mathbf{p}$  is defined to be the subset of  $X$  defined so that

$$B_X(\mathbf{p}, r) = \{\mathbf{x} \in X : |\mathbf{x} - \mathbf{p}| < r\}.$$

(Thus  $B_X(\mathbf{p}, r)$  is the set consisting of all points of  $X$  that lie within a sphere of radius  $r$  centred on the point  $\mathbf{p}$ .)

**Definition** Let  $X$  be a subset of  $\mathbb{R}^n$ . A subset  $V$  of  $X$  is said to be *open* in  $X$  if, given any point  $\mathbf{p}$  of  $V$ , there exists some strictly positive real number  $\delta$  such that  $B_X(\mathbf{p}, \delta) \subset V$ , where  $B_X(\mathbf{p}, \delta)$  is the open ball in  $X$  of radius  $\delta$  about on the point  $\mathbf{p}$ . The empty set  $\emptyset$  is also defined to be an open set in  $X$ .

**Lemma 1.3** *Let  $X$  be a subset of  $\mathbb{R}^n$ , and let  $\mathbf{p}$  be a point of  $X$ . Then, for any positive real number  $r$ , the open ball  $B_X(\mathbf{p}, r)$  in  $X$  of radius  $r$  about  $\mathbf{p}$  is open in  $X$ .*

A proof of Lemma 1.3 is to be found in Appendix A.

**Proposition 1.4** *Let  $X$  be a subset of  $\mathbb{R}^n$ . The collection of open sets in  $X$  has the following properties:—*

- (i) *the empty set  $\emptyset$  and the whole set  $X$  are both open in  $X$ ;*
- (ii) *the union of any collection of open sets in  $X$  is itself open in  $X$ ;*
- (iii) *the intersection of any finite collection of open sets in  $X$  is itself open in  $X$ .*

A proof of Proposition 1.4 is to be found in Appendix A.

**Proposition 1.5** *Let  $X$  be a subset of  $\mathbb{R}^n$ , and let  $U$  be a subset of  $X$ . Then  $U$  is open in  $X$  if and only if there exists some open set  $V$  in  $\mathbb{R}^n$  for which  $U = V \cap X$ .*

A proof of Proposition 1.5 is to be found in Appendix A.

**Lemma 1.6** *A sequence  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  of points in  $\mathbb{R}^n$  converges to a point  $\mathbf{p}$  if and only if, given any open set  $U$  which contains  $\mathbf{p}$ , there exists some positive integer  $N$  such that  $\mathbf{x}_j \in U$  for all  $j$  satisfying  $j \geq N$ .*

A proof of Lemma 1.6 is to be found in Appendix A.

**Definition** Let  $X$  be a subset of  $\mathbb{R}^n$ . A subset  $F$  of  $X$  is said to be *closed* in  $X$  if and only if its complement  $X \setminus F$  in  $X$  is open in  $X$ . (Recall that  $X \setminus F = \{\mathbf{x} \in X : \mathbf{x} \notin F\}$ .)

**Proposition 1.7** *Let  $X$  be a subset of  $\mathbb{R}^n$ . The collection of closed sets in  $X$  has the following properties:—*

- (i) *the empty set  $\emptyset$  and the whole set  $X$  are both closed in  $X$ ;*
- (ii) *the intersection of any collection of closed sets in  $X$  is itself closed in  $X$ ;*
- (iii) *the union of any finite collection of closed sets in  $X$  is itself closed in  $X$ .*

A proof of Proposition 1.7 is to be found in Appendix A.

**Lemma 1.8** *Let  $X$  be a subset of  $\mathbb{R}^n$ , and let  $F$  be a subset of  $X$  which is closed in  $X$ . Let  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  be a sequence of points of  $F$  which converges to a point  $\mathbf{p}$  of  $X$ . Then  $\mathbf{p} \in F$ .*

A proof of Lemma 1.8 is to be found in Appendix A.

**Definition** Let  $X$  and  $Y$  be a subsets of  $\mathbb{R}^m$  and  $\mathbb{R}^n$  respectively. A function  $f: X \rightarrow Y$  from  $X$  to  $Y$  is said to be *continuous* at a point  $\mathbf{p}$  of  $X$  if and only if the following criterion is satisfied:—

given any strictly positive real number  $\varepsilon$ , there exists some strictly positive real number  $\delta$  such that  $|f(\mathbf{x}) - f(\mathbf{p})| < \varepsilon$  whenever  $\mathbf{x} \in X$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta$ .

The function  $f: X \rightarrow Y$  is said to be continuous on  $X$  if and only if it is continuous at every point  $\mathbf{p}$  of  $X$ .

**Lemma 1.9** *Let  $X, Y$  and  $Z$  be subsets of  $\mathbb{R}^m, \mathbb{R}^n$  and  $\mathbb{R}^k$  respectively, and let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be functions satisfying  $f(X) \subset Y$ . Suppose that  $f$  is continuous at some point  $\mathbf{p}$  of  $X$  and that  $g$  is continuous at  $f(\mathbf{p})$ . Then the composition function  $g \circ f: X \rightarrow Z$  is continuous at  $\mathbf{p}$ .*

A proof of Lemma 1.9 is to be found in Appendix A.

**Lemma 1.10** *Let  $X$  and  $Y$  be a subsets of  $\mathbb{R}^m$  and  $\mathbb{R}^n$  respectively, and let  $f: X \rightarrow Y$  be a continuous function from  $X$  to  $Y$ . Let  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  be a sequence of points of  $X$  which converges to some point  $\mathbf{p}$  of  $X$ . Then the sequence  $f(\mathbf{x}_1), f(\mathbf{x}_2), f(\mathbf{x}_3), \dots$  converges to  $f(\mathbf{p})$ .*

A proof of Lemma 1.10 is to be found in Appendix A.

Let  $X$  and  $Y$  be a subsets of  $\mathbb{R}^m$  and  $\mathbb{R}^n$  respectively, and let  $f: X \rightarrow Y$  be a function from  $X$  to  $Y$ . Then

$$f(\mathbf{x}) = (f_1(\mathbf{x}), f_2(\mathbf{x}), \dots, f_n(\mathbf{x}))$$

for all  $\mathbf{x} \in X$ , where  $f_1, f_2, \dots, f_n$  are functions from  $X$  to  $\mathbb{R}$ , referred to as the *components* of the function  $f$ .

**Proposition 1.11** *Let  $X$  and  $Y$  be a subsets of  $\mathbb{R}^m$  and  $\mathbb{R}^n$  respectively, and let  $\mathbf{p} \in X$ . A function  $f: X \rightarrow Y$  is continuous at the point  $\mathbf{p}$  if and only if its components are all continuous at  $\mathbf{p}$ .*

A proof of Proposition 1.11 is to be found in Appendix A.

**Proposition 1.12** *Let  $X$  be a subset of  $\mathbb{R}^n$ , and let  $f: X \rightarrow \mathbb{R}$  and  $g: X \rightarrow \mathbb{R}$  be continuous functions from  $X$  to  $\mathbb{R}$ . Then the functions  $f + g$ ,  $f - g$  and  $f \cdot g$  are continuous. If in addition  $g(\mathbf{x}) \neq 0$  for all  $\mathbf{x} \in X$  then the quotient function  $f/g$  is continuous.*

A proof of Proposition 1.12 is to be found in Appendix A.

**Lemma 1.13** *Let  $X$  be a subset of  $\mathbb{R}^m$ , let  $f: X \rightarrow \mathbb{R}^n$  be a continuous function mapping  $X$  into  $\mathbb{R}^n$ , and let  $|f|: X \rightarrow \mathbb{R}$  be defined such that  $|f|(\mathbf{x}) = |f(\mathbf{x})|$  for all  $\mathbf{x} \in X$ . Then the real-valued function  $|f|$  is continuous on  $X$ .*

A proof of Proposition 1.13 is to be found in Appendix A.

Given any function  $f: X \rightarrow Y$ , we denote by  $f^{-1}(V)$  the *preimage* of a subset  $V$  of  $Y$  under the map  $f$ , defined by  $f^{-1}(V) = \{\mathbf{x} \in X : f(\mathbf{x}) \in V\}$ .

**Proposition 1.14** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^m$  and  $\mathbb{R}^n$ , and let  $f: X \rightarrow Y$  be a function from  $X$  to  $Y$ . The function  $f$  is continuous if and only if  $f^{-1}(V)$  is open in  $X$  for every open subset  $V$  of  $Y$ .*

A proof of Proposition 1.14 is to be found in Appendix A.

Let  $X$  be a subset of  $\mathbb{R}^n$ , let  $f: X \rightarrow \mathbb{R}$  be continuous, and let  $c$  be some real number. Proposition 1.14 ensures that the sets  $\{\mathbf{x} \in X : f(\mathbf{x}) > c\}$  and  $\{\mathbf{x} \in X : f(\mathbf{x}) < c\}$  are open in  $X$ . Moreover given real numbers  $a$  and  $b$  satisfying  $a < b$ , the set  $\{\mathbf{x} \in X : a < f(\mathbf{x}) < b\}$  is open in  $X$ .

**Corollary 1.15** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\varphi: X \rightarrow Y$  be a continuous function from  $X$  to  $Y$ . Then  $\varphi^{-1}(F)$  is closed in  $X$  for every subset  $F$  of  $Y$  that is closed in  $Y$ .*

A proof of Corollary 1.15 is to be found in Appendix A.

**Lemma 1.16** *Let  $X$  be a closed subset of  $n$ -dimensional Euclidean space  $\mathbb{R}^n$ . Then a subset of  $X$  is closed in  $X$  if and only if it is closed in  $\mathbb{R}^n$ .*

A proof of Lemma 1.16 is to be found in Appendix A.

### 1.3 The Multidimensional Extreme Value Theorem

**Theorem 1.17 (The Multidimensional Extreme Value Theorem)**

Let  $X$  be a non-empty closed bounded set in  $\mathbb{R}^m$ , and let  $f: X \rightarrow \mathbb{R}$  be a continuous real-valued function defined on  $X$ . Then there exist points  $\mathbf{u}$  and  $\mathbf{v}$  of  $X$  such that  $f(\mathbf{u}) \leq f(\mathbf{x}) \leq f(\mathbf{v})$  for all  $\mathbf{x} \in X$ .

A proof of Theorem 1.17 is to be found in Appendix A.

### 1.4 The Glueing Lemma

The following result, together with its generalizations, is sometimes referred to as the *Glueing Lemma*.

**Lemma 1.18 (Glueing Lemma)** Let  $\varphi: X \rightarrow \mathbb{R}^n$  be a function mapping a subset  $X$  of  $\mathbb{R}^m$  into  $\mathbb{R}^n$ . Let  $F_1, F_2, \dots, F_k$  be a finite collection of subsets of  $X$  such that  $F_i$  is closed in  $X$  for  $i = 1, 2, \dots, k$  and

$$F_1 \cup F_2 \cup \dots \cup F_k = X.$$

Then the function  $\varphi$  is continuous on  $X$  if and only if the restriction of  $\varphi$  to  $F_i$  is continuous on  $F_i$  for  $i = 1, 2, \dots, k$ .

**Proof** Suppose that  $\varphi: X \rightarrow \mathbb{R}^n$  is continuous. Then it follows directly from the definition of continuity that the restriction of  $\varphi$  to each subset of  $X$  is continuous on that subset. Therefore the restriction of  $\varphi$  to  $F_i$  is continuous on  $F_i$  for  $i = 1, 2, \dots, k$ .

Conversely we must prove that if the restriction of the function  $\varphi$  to  $F_i$  is continuous on  $F_i$  for  $i = 1, 2, \dots, k$  then the function  $\varphi: X \rightarrow \mathbb{R}^n$  is continuous. Let  $\mathbf{p}$  be a point of  $X$ , and let some positive real number  $\varepsilon$  be given. Then there exist positive real numbers  $\delta_1, \delta_2, \dots, \delta_k$  satisfying the following conditions:—

- (i) if  $\mathbf{p} \in F_i$ , where  $1 \leq i \leq k$ , and if  $\mathbf{x} \in F_i$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta_i$  then  $|\varphi(\mathbf{x}) - \varphi(\mathbf{p})| < \varepsilon$ ;
- (ii) if  $\mathbf{p} \notin F_i$ , where  $1 \leq i \leq k$ , and if  $\mathbf{x} \in X$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta_i$  then  $\mathbf{x} \notin F_i$ .

Indeed the continuity of the function  $\varphi$  on each set  $F_i$  ensures that  $\delta_i$  may be chosen to satisfy (i) for each integer  $i$  between 1 and  $k$  for which  $\mathbf{p} \in F_i$ . Also the requirement that  $F_i$  be closed in  $X$  ensures that  $X \setminus F_i$  is open in  $X$  and therefore  $\delta_i$  may be chosen to satisfy (ii) for each integer  $i$  between 1 and  $k$  for which  $\mathbf{p} \notin F_i$ .

Let  $\delta$  be the minimum of  $\delta_1, \delta_2, \dots, \delta_k$ . Then  $\delta > 0$ . Let  $\mathbf{x} \in X$  satisfy  $|\mathbf{x} - \mathbf{p}| < \delta$ . If  $\mathbf{p} \notin F_i$  then the choice of  $\delta_i$  ensures that if  $\mathbf{x} \notin F_i$ . But  $X$  is the union of the sets  $F_1, F_2, \dots, F_k$ , and therefore there must exist some integer  $i$  between 1 and  $k$  for which  $\mathbf{x} \in F_i$ . Then  $\mathbf{p} \in F_i$ , and the choice of  $\delta_i$  ensures that  $|\varphi(\mathbf{x}) - \varphi(\mathbf{p})| < \varepsilon$ . We have thus shown that  $|\varphi(\mathbf{x}) - \varphi(\mathbf{p})| < \varepsilon$  for all points  $\mathbf{x}$  of  $X$  that satisfy  $|\mathbf{x} - \mathbf{p}| < \delta$ . It follows that  $\varphi: X \rightarrow \mathbb{R}^n$  is continuous, as required. ■

## 1.5 Lebesgue Numbers

**Definition** Let  $X$  be a subset of  $n$ -dimensional Euclidean space  $\mathbb{R}^n$ . A collection of subsets of  $\mathbb{R}^n$  is said to *cover*  $X$  if and only if every point of  $X$  belongs to at least one of these subsets.

**Definition** Let  $X$  be a subset of  $n$ -dimensional Euclidean space  $\mathbb{R}^n$ . An *open cover* of  $X$  is a collection of subsets of  $X$  that are open in  $X$  and cover the set  $X$ .

**Proposition 1.19** *Let  $X$  be a closed bounded set in  $n$ -dimensional Euclidean space, and let  $\mathcal{V}$  be an open cover of  $X$ . Then there exists a positive real number  $\delta_L$  with the property that, given any point  $\mathbf{u}$  of  $X$ , there exists a member  $V$  of the open cover  $\mathcal{V}$  for which*

$$\{\mathbf{x} \in X : |\mathbf{x} - \mathbf{u}| < \delta_L\} \subset V.$$

**Proof** Let

$$B_X(\mathbf{u}, \delta) = \{\mathbf{x} \in X : |\mathbf{x} - \mathbf{u}| < \delta\}$$

for all  $\mathbf{u} \in X$  and for all positive real numbers  $\delta$ . Suppose that there did not exist any positive real number  $\delta_L$  with the stated property.

Then, given any positive number  $\delta$ , there would exist a point  $\mathbf{u}$  of  $X$  for which the set  $B_X(\mathbf{u}, \delta)$  would not be wholly contained within any open set  $V$  belonging to the open cover  $\mathcal{V}$ . Consequently there would exist an infinite sequence

$$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \dots$$

of points of  $X$  with the property that, for each positive integer  $j$ , the set  $B_X(\mathbf{u}_j, 1/j)$  would not be wholly contained within any open set  $V$  belonging to the open cover  $\mathcal{V}$ . The multidimensional Bolzano-Weierstrass Theorem (Theorem 1.2) would then ensure the existence of a convergent subsequence

$$\mathbf{u}_{j_1}, \mathbf{u}_{j_2}, \mathbf{u}_{j_3}, \dots$$



of this infinite sequence.

Let  $\mathbf{p}$  be the limit of this convergent subsequence. Then the point  $\mathbf{p}$  would then belong to  $X$ , because  $X$  is closed (see Lemma 1.8). But then the point  $\mathbf{p}$  would belong to an open set  $V$  belonging to the open cover  $\mathcal{V}$ . It would then follow from the definition of open sets that there would exist a positive real number  $\delta$  for which  $B_X(\mathbf{p}, 2\delta) \subset V$ . Let  $j = j_k$  for a positive integer  $k$  large enough to ensure that both  $1/j < \delta$  and  $\mathbf{u}_j \in B_X(\mathbf{p}, \delta)$ . The Triangle Inequality would then ensure that every point of  $X$  within a distance  $1/j$  of the point  $\mathbf{u}_j$  would lie within a distance  $2\delta$  of the point  $\mathbf{p}$ , and therefore

$$B_X(\mathbf{u}_j, 1/j) \subset B_X(\mathbf{p}, 2\delta) \subset V.$$

But we supposed that the point  $\mathbf{u}_j$  was chosen so as to ensure that the set  $B_X(\mathbf{u}_j, 1/j)$  was not wholly contained within any open set  $V$  belonging to the open cover  $\mathcal{V}$ . Thus a logical contradiction as resulted from the assumption that there is no positive real number  $\delta_L$  with the property that, given any point  $\mathbf{u}$  of  $X$ , the set  $B_X(\mathbf{u}, \delta_L)$  is not wholly contained within any open set belonging to the open cover  $\mathcal{V}$ . Consequently some positive real number  $\delta_L$  satisfying this property must exist, and thus the required result has been proved. ■

**Definition** Let  $X$  be a subset of  $n$ -dimensional Euclidean space, and let  $\mathcal{V}$  be an open cover of  $X$ . A positive real number  $\delta_L$  is said to be a *Lebesgue number* for the open cover  $\mathcal{V}$  if, given any point  $\mathbf{p}$  of  $X$ , there exists some member  $V$  of the open cover  $\mathcal{V}$  for which

$$\{\mathbf{x} \in X : |\mathbf{x} - \mathbf{p}| < \delta_L\} \subset V.$$

Proposition 1.19 ensures that, given any open cover of a closed bounded subset of  $n$ -dimensional Euclidean space, there exists a positive real number that is a Lebesgue number for that open cover.

**Definition** The *diameter*  $\text{diam}(A)$  of a bounded subset  $A$  of  $n$ -dimensional Euclidean space is defined so that

$$\text{diam}(A) = \sup\{|\mathbf{x} - \mathbf{y}| : \mathbf{x}, \mathbf{y} \in A\}.$$

It follows from this definition that  $\text{diam}(A)$  is the smallest real number  $K$  with the property that  $|\mathbf{x} - \mathbf{y}| \leq K$  for all  $\mathbf{x}, \mathbf{y} \in A$ .

**Lemma 1.20** *Let  $X$  be a bounded subset of  $n$ -dimensional Euclidean space, and let  $\delta$  be a positive real number. Then there exists a finite collection  $A_1, A_2, \dots, A_s$  of subsets of  $X$  such that the  $\text{diam}(A_i) < \delta$  for  $i = 1, 2, \dots, s$  and*

$$X = A_1 \cup A_2 \cup \dots \cup A_s.$$

**Proof** Let  $b$  be a real number satisfying  $0 < \sqrt{n}b < \delta$  and, for each  $n$ -tuple  $(j_1, j_2, \dots, j_n)$  of integers, let  $H_{(j_1, j_2, \dots, j_n)}$  denote the hypercube in  $\mathbb{R}^n$  defined such that

$$\begin{aligned} H_{(j_1, j_2, \dots, j_n)} \\ = \{ (x_1, x_2, \dots, x_n) \in \mathbb{R}^n : j_i b \leq x_i \leq (j_i + 1)b \text{ for } i = 1, 2, \dots, n \}. \end{aligned}$$

Note that if  $\mathbf{u}$  and  $\mathbf{v}$  are points of  $H_{(j_1, j_2, \dots, j_n)}$  for some  $n$ -tuple  $(j_1, j_2, \dots, j_n)$  of integers then  $|u_i - v_i| < b$  for  $i = 1, 2, \dots, n$ , and therefore  $|\mathbf{u} - \mathbf{v}| \leq \sqrt{n}b < \delta$ . Therefore the diameter of each hypercube  $H_{(j_1, j_2, \dots, j_n)}$  is less than  $\delta$ .

The boundedness of the set  $X$  ensures that there are only finitely many  $n$ -tuples  $(j_1, j_2, \dots, j_n)$  of integers for which  $X \cap H_{(j_1, j_2, \dots, j_n)}$  is non-empty. It follows that  $X$  is covered by a finite collection  $A_1, A_2, \dots, A_k$  of subsets of  $X$ , where each of these subsets is of the form  $X \cap H_{(j_1, j_2, \dots, j_n)}$  for some  $n$ -tuple  $(j_1, j_2, \dots, j_n)$  of integers. These subsets all have diameter less than  $\delta$ . The result follows. ■

**Definition** Let  $\mathcal{V}$  and  $\mathcal{W}$  be open covers of some subset  $X$  of a Euclidean space. Then  $\mathcal{W}$  is said to be a *subcover* of  $\mathcal{V}$  if and only if every open set belonging to  $\mathcal{W}$  also belongs to  $\mathcal{V}$ .

**Definition** A subset  $X$  of a Euclidean space is said to be *compact* if and only if every open cover of  $X$  possesses a finite subcover.

**Theorem 1.21 (The Multidimensional Heine-Borel Theorem)** *A subset of  $n$ -dimensional Euclidean space  $\mathbb{R}^n$  is compact if and only if it is both closed and bounded.*

**Proof** Let  $X$  be a compact subset of  $\mathbb{R}^n$  and let

$$V_j = \{\mathbf{x} \in X : |\mathbf{x}| < j\}$$

for all positive integers  $j$ . Then the sets  $V_1, V_2, V_3, \dots$  constitute an open cover of  $X$ . This open cover has a finite subcover, and therefore there exist positive integers  $j_1, j_2, \dots, j_k$  such that

$$X \subset V_{j_1} \cup V_{j_2} \cup \dots \cup V_{j_k}.$$

Let  $M$  be the largest of the positive integers  $j_1, j_2, \dots, j_k$ . Then  $|\mathbf{x}| \leq M$  for all  $\mathbf{x} \in X$ . Thus the set  $X$  is bounded.

Let  $\mathbf{q}$  be a point of  $\mathbb{R}^n$  that does not belong to  $X$ , and let

$$W_j = \left\{ \mathbf{x} \in X : |\mathbf{x} - \mathbf{q}| > \frac{1}{j} \right\}$$

for all positive integers  $j$ . Then the sets  $W_1, W_2, W_3, \dots$  constitute an open cover of  $X$ . This open cover has a finite subcover, and therefore there exist positive integers  $j_1, j_2, \dots, j_k$  such that

$$X \subset W_{j_1} \cup W_{j_2} \cup \dots \cup W_{j_k}.$$

Let  $\delta = 1/M$ , where  $M$  is the largest of the positive integers  $j_1, j_2, \dots, j_k$ . Then  $|\mathbf{x} - \mathbf{q}| \geq \delta$  for all  $\mathbf{x} \in X$  and thus the open ball of radius  $\delta$  about the point  $\mathbf{q}$  does not intersect the set  $X$ . We conclude that the set  $X$  is closed. We have now shown that every compact subset of  $\mathbb{R}^n$  is both closed and bounded.

We now prove the converse. Let  $X$  be a closed bounded subset of  $\mathbb{R}^n$ , and let  $\mathcal{V}$  be an open cover of  $X$ . It follows from Proposition 1.19 that there exists a Lebesgue number  $\delta_L$  for the open cover  $\mathcal{V}$ . It then follows from Lemma 1.20 that there exist subsets  $A_1, A_2, \dots, A_s$  of  $X$  such that  $\text{diam}(A_i) < \delta_L$  for  $i = 1, 2, \dots, s$  and

$$X = A_1 \cup A_2 \cup \dots \cup A_s.$$

We may suppose that  $A_i$  is non-empty for  $i = 1, 2, \dots, s$  (because if  $A_i = \emptyset$  then  $A_i$  could be deleted from the list). Choose  $\mathbf{p}_i \in A_i$  for  $i = 1, 2, \dots, s$ . Then  $A_i \subset B_X(\mathbf{p}_i, \delta_L)$  for  $i = 1, 2, \dots, s$ . The definition of the Lebesgue number  $\delta_L$  then ensures that there exist members  $V_1, V_2, \dots, V_s$  of the open cover  $\mathcal{V}$  such that  $B_X(\mathbf{p}_i, \delta_L) \subset V_i$  for  $i = 1, 2, \dots, s$ . Then  $A_i \subset V_i$  for  $i = 1, 2, \dots, s$ , and therefore

$$X \subset V_1 \cup V_2 \cup \dots \cup V_s.$$

Thus  $V_1, V_2, \dots, V_s$  constitute a finite subcover of the open cover  $\mathcal{U}$ . We have therefore proved that every closed bounded subset of  $n$ -dimensional Euclidean space is compact, as required. ■

## 2 Correspondences and Hemicontinuity

### 2.1 Correspondences

Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A *correspondence*  $\Phi: X \rightrightarrows Y$  assigns to each point  $\mathbf{x}$  of  $X$  a subset  $\Phi(\mathbf{x})$  of  $Y$ .

The *power set*  $\mathcal{P}(Y)$  of  $Y$  is the set whose elements are the subsets of  $Y$ . A correspondence  $\Phi: X \rightrightarrows Y$  may be regarded as a function from  $X$  to  $\mathcal{P}(Y)$ .

Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Then the following definitions apply:–

- the correspondence  $\Phi: X \rightarrow Y$  is said to be *non-empty-valued* if  $\Phi(\mathbf{x})$  is a non-empty subset of  $Y$  for all  $\mathbf{x} \in X$ ;
- the correspondence  $\Phi: X \rightarrow Y$  is said to be *closed-valued* if  $\Phi(\mathbf{x})$  is a closed subset of  $Y$  for all  $\mathbf{x} \in X$ ;
- the correspondence  $\Phi: X \rightarrow Y$  is said to be *compact-valued* if  $\Phi(\mathbf{x})$  is a compact subset of  $Y$  for all  $\mathbf{x} \in X$ .

The multidimensional Heine-Borel Theorem (Theorem 1.21) ensures that the correspondence  $\Phi: X \rightarrow Y$  is compact-valued if and only if  $\Phi(\mathbf{x})$  is a closed bounded subset of  $\mathbb{R}^m$  for all  $\mathbf{x} \in X$ .

**Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A correspondence  $\Phi: X \rightrightarrows Y$  is said to be *upper hemicontinuous* at a point  $\mathbf{p}$  of  $X$  if, given any set  $V$  in  $Y$  that is open in  $Y$  and satisfies  $\Phi(\mathbf{p}) \subset V$ , there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \subset V$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . The correspondence  $\Phi$  is upper hemicontinuous on  $X$  if it is upper hemicontinuous at each point of  $X$ .

**Example** Let  $F: \mathbb{R} \rightrightarrows \mathbb{R}$  and  $G: \mathbb{R} \rightrightarrows \mathbb{R}$  be the correspondences from  $\mathbb{R}$  to  $\mathbb{R}$  defined such that

$$F(x) = \begin{cases} [1, 2] & \text{if } x < 0, \\ [0, 3] & \text{if } x \geq 0, \end{cases}$$

and

$$G(x) = \begin{cases} [1, 2] & \text{if } x \leq 0, \\ [0, 3] & \text{if } x > 0, \end{cases}$$

The correspondences  $F$  and  $G$  are upper hemicontinuous at  $x$  for all non-zero real numbers  $x$ . The correspondence  $F$  is also upper hemicontinuous at 0,

for if  $V$  is an open set in  $\mathbb{R}$  and if  $F(0) \subset V$  then  $[0, 3] \subset V$  and therefore  $F(x) \in V$  for all real numbers  $x$ .

However the correspondence  $G$  is not upper hemicontinuous at 0. Indeed let

$$V = \{y \in \mathbb{R} : \frac{1}{2} < y < \frac{5}{2}\}.$$

Then  $G(0) \subset V$ , but  $G(x)$  is not contained in  $V$  for any positive real number  $x$ . Therefore there cannot exist any positive real number  $\delta$  such that  $G(x) \subset V$  whenever  $|x| < \delta$ .

Let

$$\text{Graph}(F) = \{(x, y) \in \mathbb{R}^2 : y \in F(x)\}$$

and

$$\text{Graph}(G) = \{(x, y) \in \mathbb{R}^2 : y \in G(x)\}.$$

Then  $\text{Graph}(F)$  is a closed subset of  $\mathbb{R}^2$  but  $\text{Graph}(G)$  is not a closed subset of  $\mathbb{R}^2$ .

**Example** Let  $S^1$  be the unit circle in  $\mathbb{R}^2$ , defined such that

$$S^1 = \{(u, v) \in \mathbb{R}^2 : u^2 + v^2 = 1\},$$

let  $Z$  be the closed square with corners at  $(1, 1)$ ,  $(-1, 1)$ ,  $(-1, -1)$  and  $(1, -1)$ , so that

$$Z = \{(x, y) \in \mathbb{R}^2 : -1 \leq x \leq 1 \text{ and } -1 \leq y \leq 1\}.$$

Let  $g_{(u,v)}: \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined for all  $(u, v) \in S^1$  such that

$$g_{(u,v)}(x, y) = ux + vy,$$

and let  $\Phi: S^1 \rightrightarrows \mathbb{R}^2$  be defined such that, for all  $(u, v) \in S^1$ ,  $\Phi(u, v)$  is the subset of  $\mathbb{R}^2$  consisting of the point of points of  $Z$  at which the linear functional  $g_{(u,v)}$  attains its maximum value on  $Z$ . Thus a point  $(x, y)$  of  $Z$  belongs to  $\Phi(u, v)$  if and only if  $g_{(u,v)}(x, y) \geq g_{(u,v)}(x', y')$  for all  $(x', y') \in Z$ . Then

$$\Phi(u, v) = \begin{cases} \{(1, 1)\} & \text{if } u > 0 \text{ and } v > 0; \\ \{(x, 1) : -1 \leq x \leq 1\} & \text{if } u = 0 \text{ and } v > 0; \\ \{(-1, 1)\} & \text{if } u < 0 \text{ and } v > 0; \\ \{(-1, y) : -1 \leq y \leq 1\} & \text{if } u < 0 \text{ and } v = 0; \\ \{(-1, -1)\} & \text{if } u < 0 \text{ and } v < 0; \\ \{(x, -1) : -1 \leq x \leq 1\} & \text{if } u = 0 \text{ and } v < 0; \\ \{(1, -1)\} & \text{if } u > 0 \text{ and } v < 0; \\ \{(1, y) : -1 \leq y \leq 1\} & \text{if } u > 0 \text{ and } v = 0. \end{cases}$$

It is a straightforward exercise to verify that the correspondence  $\Phi: S^1 \rightrightarrows \mathbb{R}^2$  is upper hemicontinuous.

Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence between  $X$  and  $Y$ . Given any subset  $V$  of  $Y$ , we denote by  $\Phi^+(V)$  the subset of  $X$  defined such that

$$\Phi^+(V) = \{\mathbf{x} \in X : \Phi(\mathbf{x}) \subset V\}.$$

**Lemma 2.1** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A correspondence  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous on  $X$  if and only if, given any set  $V$  in  $Y$  that is open in  $Y$ , the set  $\Phi^+(V)$  is open in  $X$ .*

**Proof** First suppose that  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at each point of  $X$ . Let  $V$  be an open set in  $Y$  and let  $\mathbf{p} \in \Phi^+(V)$ . Then  $\Phi(\mathbf{p}) \subset V$ . It then follows from the definition of upper hemicontinuity that there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \subset V$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . But then  $\mathbf{x} \in \Phi^+(V)$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . It follows that  $\Phi^+(V)$  is open in  $X$ .

Conversely suppose that  $\Phi: X \rightrightarrows Y$  is a correspondence with the property that, for all subsets  $V$  of  $Y$  that are open in  $Y$ ,  $\Phi^+(V)$  is open in  $X$ . Let  $\mathbf{p} \in X$ , and let  $V$  be an open set in  $Y$  satisfying  $\Phi(\mathbf{p}) \subset V$ . Then  $\Phi^+(V)$  is open in  $X$  and  $\mathbf{p} \in \Phi^+(V)$ , and therefore there exists some positive number  $\delta$  such that

$$\{\mathbf{x} \in X : |\mathbf{x} - \mathbf{p}| < \delta\} \subset \Phi^+(V).$$

Then  $\Phi(\mathbf{x}) \subset V$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . Thus  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at  $\mathbf{p}$ . The result follows. ■

**Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A correspondence  $\Phi: X \rightrightarrows Y$  is said to be *lower hemicontinuous* at a point  $\mathbf{p}$  of  $X$  if, given any set  $V$  in  $Y$  that is open in  $Y$  and satisfies  $\Phi(\mathbf{p}) \cap V \neq \emptyset$ , there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \cap V \neq \emptyset$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . The correspondence  $\Phi$  is lower hemicontinuous on  $X$  if it is lower hemicontinuous at each point of  $X$ .

**Example** Let  $F: \mathbb{R} \rightrightarrows \mathbb{R}$  and  $G: \mathbb{R} \rightrightarrows \mathbb{R}$  be the correspondences from  $\mathbb{R}$  to  $\mathbb{R}$  defined such that

$$F(x) = \begin{cases} [1, 2] & \text{if } x < 0, \\ [0, 3] & \text{if } x \geq 0, \end{cases}$$

and

$$G(x) = \begin{cases} [1, 2] & \text{if } x \leq 0, \\ [0, 3] & \text{if } x > 0, \end{cases}$$

The correspondences  $F$  and  $G$  are lower hemicontinuous at  $x$  for all non-zero real numbers  $x$ . The correspondence  $G$  is also lower hemicontinuous at 0, for

if  $V$  is an open set in  $\mathbb{R}$  and if  $G(0) \cap V \neq \emptyset$  then  $[1, 2] \cap V \neq \emptyset$  and therefore  $G(x) \cap V \neq \emptyset$  for all real numbers  $x$ .

However the correspondence  $F$  is not lower hemicontinuous at 0. Indeed let

$$V = \{y \in \mathbb{R} : 0 < y < \frac{1}{2}\}.$$

Then  $F(0) \cap V \neq \emptyset$ , but  $F(x) \cap V = \emptyset$  for all negative real numbers  $x$ . Therefore there cannot exist any positive real number  $\delta$  such that  $F(x) \cap V \neq \emptyset$  whenever  $|x| < \delta$ .

Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence between  $X$  and  $Y$ . Given any subset  $V$  of  $Y$ , we denote by  $\Phi^-(V)$  the subset of  $X$  defined such that

$$\Phi^-(V) = \{\mathbf{x} \in X : \Phi(\mathbf{x}) \cap V \neq \emptyset\}.$$

**Lemma 2.2** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A correspondence  $\Phi: X \rightrightarrows Y$  is lower hemicontinuous on  $X$  if and only if, given any set  $V$  in  $Y$  that is open in  $Y$ , the set  $\Phi^-(V)$  is open in  $X$ .*

**Proof** First suppose that  $\Phi: X \rightrightarrows Y$  is lower hemicontinuous at each point of  $X$ . Let  $V$  be an open set in  $Y$  and let  $\mathbf{p} \in \Phi^-(V)$ . Then  $\Phi(\mathbf{p}) \cap V$  is non-empty. It then follows from the definition of lower hemicontinuity that there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \cap V$  is non-empty for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . But then  $\mathbf{x} \in \Phi^-(V)$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . It follows that  $\Phi^-(V)$  is open in  $X$ .

Conversely suppose that  $\Phi: X \rightrightarrows Y$  is a correspondence with the property that, for all subsets  $V$  of  $Y$  that are open in  $Y$ ,  $\Phi^-(V)$  is open in  $X$ . Let  $\mathbf{p} \in X$ , and let  $V$  be an open set in  $Y$  satisfying  $\Phi(\mathbf{p}) \cap V \neq \emptyset$ . Then  $\Phi^-(V)$  is open in  $X$  and  $\mathbf{p} \in \Phi^-(V)$ , and therefore there exists some positive number  $\delta$  such that

$$\{\mathbf{x} \in X : |\mathbf{x} - \mathbf{p}| < \delta\} \subset \Phi^-(V).$$

Then  $\Phi(\mathbf{x}) \cap V \neq \emptyset$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . Thus  $\Phi: X \rightrightarrows Y$  is lower hemicontinuous at  $\mathbf{p}$ . The result follows.  $\blacksquare$

**Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A correspondence  $\Phi: X \rightrightarrows Y$  is said to be *continuous* at a point  $\mathbf{p}$  of  $X$  if it is both upper hemicontinuous and lower hemicontinuous at  $\mathbf{p}$ . The correspondence  $\Phi$  is continuous on  $X$  if it is continuous at each point of  $X$ .

**Lemma 2.3** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, let  $\varphi: X \rightarrow Y$  be a function from  $X$  to  $Y$ , and let  $\Phi: X \rightrightarrows Y$  be the correspondence defined such that  $\Phi(\mathbf{x}) = \{\varphi(\mathbf{x})\}$  for all  $\mathbf{x} \in X$ . Then  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous if and only if  $\varphi: X \rightarrow Y$  is continuous. Similarly  $\Phi: X \rightrightarrows Y$  is lower hemicontinuous if and only if  $\varphi: X \rightarrow Y$  is continuous.*

**Proof** The function  $\varphi: X \rightarrow Y$  is continuous if and only if

$$\{\mathbf{x} \in X : \varphi(\mathbf{x}) \in V\}$$

is open in  $X$  for all subsets  $V$  of  $Y$  that are open in  $Y$  (see Proposition 1.14). Let  $V$  be a subset of  $Y$  that is open in  $Y$ . Then  $\Phi(\mathbf{x}) \subset V$  if and only if  $\varphi(\mathbf{x}) \in V$ . Also  $\Phi(\mathbf{x}) \cap V \neq \emptyset$  if and only if  $\varphi(\mathbf{x}) \in V$ . The result therefore follows from the definitions of upper and lower hemicontinuity. ■

## 2.2 The Graph of a Correspondence

Let  $m$  and  $n$  be integers. Then the Cartesian product  $\mathbb{R}^n \times \mathbb{R}^m$  of the Euclidean spaces  $\mathbb{R}^n$  and  $\mathbb{R}^m$  of dimensions  $n$  and  $m$  is itself a Euclidean space of dimension  $n + m$  whose Euclidean norm is characterized by the property that

$$|(\mathbf{x}, \mathbf{y})|^2 = |\mathbf{x}|^2 + |\mathbf{y}|^2$$

for all  $\mathbf{x} \in \mathbb{R}^n$  and  $\mathbf{y} \in \mathbb{R}^m$ .

**Lemma 2.4** *Let  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  and  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$  be infinite sequences of points in  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\mathbf{p} \in \mathbb{R}^n$  and  $\mathbf{q} \in \mathbb{R}^m$ . Then the infinite sequence*

$$(\mathbf{x}_1, \mathbf{y}_1), (\mathbf{x}_2, \mathbf{y}_2), (\mathbf{x}_3, \mathbf{y}_3), \dots$$

*converges in  $\mathbb{R}^n \times \mathbb{R}^m$  to the point  $(\mathbf{p}, \mathbf{q})$  if and only if the infinite sequence Let  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  converges to the point  $\mathbf{p}$  and the infinite sequence  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$  converges to the point  $\mathbf{q}$ .*

**Proof** Suppose that the infinite sequence

$$(\mathbf{x}_1, \mathbf{y}_1), (\mathbf{x}_2, \mathbf{y}_2), (\mathbf{x}_3, \mathbf{y}_3), \dots$$

converges in  $\mathbb{R}^n \times \mathbb{R}^m$  to the point  $(\mathbf{p}, \mathbf{q})$ . Let some strictly positive real number  $\varepsilon$  be given. Then there exists some positive integer  $N$  such that

$$|\mathbf{x}_j - \mathbf{p}|^2 + |\mathbf{y}_j - \mathbf{q}|^2 < \varepsilon^2$$



whenever  $j \geq N$ . But then

$$|\mathbf{x}_j - \mathbf{p}| < \varepsilon \quad \text{and} \quad |\mathbf{y}_j - \mathbf{q}| < \varepsilon$$

whenever  $j \geq N$ . It follows that  $\mathbf{x}_j \rightarrow \mathbf{p}$  and  $\mathbf{y}_j \rightarrow \mathbf{q}$  as  $j \rightarrow +\infty$ .

Conversely suppose that  $\mathbf{x}_j \rightarrow \mathbf{p}$  and  $\mathbf{y}_j \rightarrow \mathbf{q}$  as  $j \rightarrow +\infty$ . Let some positive real number  $\varepsilon$  be given. Then there exist positive integers  $N_1$  and  $N_2$  such that  $|\mathbf{x}_j - \mathbf{p}| < \varepsilon/\sqrt{2}$  whenever  $j \geq N_1$  and  $|\mathbf{y}_j - \mathbf{q}| < \varepsilon/\sqrt{2}$  whenever  $j \geq N_2$ . Let  $N$  be the maximum of  $N_1$  and  $N_2$ . Then

$$|\mathbf{x}_j - \mathbf{p}|^2 + |\mathbf{y}_j - \mathbf{q}|^2 < \varepsilon^2$$

whenever  $j \geq N$ . It follows that  $(\mathbf{x}_j, \mathbf{y}_j) \rightarrow (\mathbf{p}, \mathbf{q})$  as  $j \rightarrow +\infty$ , as required. ■

**Lemma 2.5** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $V$  be a subset of  $X \times Y$ . Then  $V$  is open in  $X \times Y$  if and only if, given any point  $(\mathbf{p}, \mathbf{q})$  of  $V$ , where  $\mathbf{p} \in X$  and  $\mathbf{q} \in Y$ , there exist subsets  $W_X$  and  $W_Y$  of  $X$  and  $Y$  respectively such that  $\mathbf{p} \in W_X$ ,  $\mathbf{q} \in W_Y$ ,  $W_X$  is open in  $X$ ,  $W_Y$  is open in  $Y$  and  $W_X \times W_Y \subset V$ .*

**Proof** Let  $V$  be a subset of  $X \times Y$  and let  $(\mathbf{p}, \mathbf{q}) \in V$ , where  $\mathbf{p} \in X$  and  $\mathbf{q} \in Y$ .

Suppose that  $V$  is open in  $X \times Y$ . Then there exists a positive real number  $\delta$  such that  $(\mathbf{x}, \mathbf{y}) \in V$  for all  $\mathbf{x} \in X$  and  $\mathbf{y} \in Y$  satisfying

$$|\mathbf{x} - \mathbf{p}|^2 + |\mathbf{y} - \mathbf{q}|^2 < \delta^2.$$

Let

$$W_X = \left\{ \mathbf{x} \in X : |\mathbf{x} - \mathbf{p}| < \frac{\delta}{\sqrt{2}} \right\}$$

and

$$W_Y = \left\{ \mathbf{y} \in Y : |\mathbf{y} - \mathbf{q}| < \frac{\delta}{\sqrt{2}} \right\}$$

If  $\mathbf{x} \in W_X$  and  $\mathbf{y} \in W_Y$  then

$$|\mathbf{x} - \mathbf{p}|^2 + |\mathbf{y} - \mathbf{q}|^2 < 2 \left( \frac{\delta}{\sqrt{2}} \right)^2 = \delta^2$$

and therefore  $(\mathbf{x}, \mathbf{y}) \in V$ . It follows that  $W_X \times W_Y \subset V$ .

Conversely suppose that there exist open sets  $W_X$  and  $W_Y$  in  $X$  and  $Y$  respectively such that  $\mathbf{p} \in W_X$ ,  $\mathbf{q} \in W_Y$  and  $W_X \times W_Y \subset V$ . Then there exists some positive real number  $\delta$  such that  $\mathbf{x} \in W_X$  for all  $\mathbf{x} \in X$  satisfying

$|\mathbf{x} - \mathbf{p}| < \delta$  and also  $\mathbf{y} \in W_Y$  for all  $\mathbf{y} \in Y$  satisfying  $|\mathbf{y} - \mathbf{q}| < \delta$ . If  $(\mathbf{x}, \mathbf{y})$  is a point of  $X \times Y$  that lies within a distance  $\delta$  of  $(\mathbf{p}, \mathbf{q})$  then  $|\mathbf{x} - \mathbf{p}| < \delta$  and  $|\mathbf{y} - \mathbf{q}| < \delta$ , and therefore  $(\mathbf{x}, \mathbf{y}) \in W_X \times W_Y$ . But  $W_X \times W_Y \subset V$ . It follows that the open ball of radius  $\delta$  about the point  $(\mathbf{p}, \mathbf{q})$  is wholly contained within the subset  $V$  of  $X \times Y$ . The result follows. ■

**Proposition 2.6** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $G$  be a subset of  $X \times Y$ . Then  $G$  is closed in  $X \times Y$  if and only if*

$$(\lim_{j \rightarrow \infty} \mathbf{x}_j, \lim_{j \rightarrow \infty} \mathbf{y}_j) \in G$$

*for all convergent infinite sequences  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  in  $X$  and for all convergent infinite sequences  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$  in  $Y$  with the property that  $(\mathbf{x}_j, \mathbf{y}_j) \in G$  for all positive integers  $j$ .*

**Proof** Suppose that  $G$  is closed in  $X \times Y$ . Let  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  be an infinite sequence in  $X$  converging to some point  $\mathbf{p}$  of  $X$  and let  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$  be an infinite sequence in  $Y$  converging to a point  $\mathbf{q}$  of  $Y$ , where  $(\mathbf{x}_j, \mathbf{y}_j) \in G$  for all positive integers  $j$ . We must prove that  $(\mathbf{p}, \mathbf{q}) \in G$ . Now the infinite sequence consisting of the ordered pairs  $(\mathbf{x}_j, \mathbf{y}_j)$  converges in  $X \times Y$  to  $(\mathbf{p}, \mathbf{q})$  (see Lemma 2.4). Now every infinite sequence contained in  $G$  that converges to a point of  $X \times Y$  must converge to a point of  $G$ , because  $G$  is closed in  $X \times Y$  (see Lemma 1.8). It follows that  $(\mathbf{p}, \mathbf{q}) \in G$ .

Now suppose that  $G$  is not closed in  $X \times Y$ . Then the complement of  $G$  in  $X \times Y$  is not open, and therefore there exists a point  $(\mathbf{p}, \mathbf{q})$  of  $X \times Y$  that does not belong to  $G$  though every open ball of positive radius about the point  $(\mathbf{p}, \mathbf{q})$  intersects  $G$ . It follows that, given any positive integer  $j$ , the open ball of radius  $1/j$  about the point  $(\mathbf{p}, \mathbf{q})$  intersects  $G$  and therefore there exist  $\mathbf{x}_j \in X$  and  $\mathbf{y}_j \in Y$  for which  $|\mathbf{x}_j - \mathbf{p}| < 1/j$ ,  $|\mathbf{y}_j - \mathbf{q}| < 1/j$  and  $(\mathbf{x}_j, \mathbf{y}_j) \in G$ . Then  $\lim_{j \rightarrow +\infty} \mathbf{x}_j = \mathbf{p}$  and  $\lim_{j \rightarrow +\infty} \mathbf{y}_j = \mathbf{q}$  and therefore

$$(\lim_{j \rightarrow \infty} \mathbf{x}_j, \lim_{j \rightarrow \infty} \mathbf{y}_j) \notin G.$$

The result follows. ■

**Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\varphi: X \rightarrow Y$  be a function from  $X$  to  $Y$ . The *graph*  $\text{Graph}(\varphi)$  of the function  $\varphi$  is the subset of  $\mathbb{R}^n \times \mathbb{R}^m$  defined so that

$$\text{Graph}(\varphi) = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^n \times \mathbb{R}^m : \mathbf{x} \in X \text{ and } \mathbf{y} = \varphi(\mathbf{x})\}.$$

**Definition** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence between  $X$  and  $Y$ . The *graph*  $\text{Graph}(\Phi)$  of the correspondence  $\Phi$  is the subset of  $\mathbb{R}^n \times \mathbb{R}^m$  defined so that

$$\text{Graph}(\Phi) = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^n \times \mathbb{R}^m : \mathbf{x} \in X \text{ and } \mathbf{y} \in \Phi(\mathbf{x})\}.$$

**Lemma 2.7** Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\varphi: X \rightarrow Y$  be a function from  $X$  to  $Y$ . Suppose that  $\varphi: X \rightarrow Y$  is continuous. Then the graph  $\text{Graph}(\varphi)$  of the function  $\varphi$  is closed in  $X \times Y$ .

**Proof** Let  $\psi: X \times Y \rightarrow Y$  be the function defined such that

$$\psi(\mathbf{x}, \mathbf{y}) = \mathbf{y} - \varphi(\mathbf{x})$$

for all  $\mathbf{x} \in X$  and  $\mathbf{y} \in Y$ . Then  $\text{Graph}(\varphi) = \psi^{-1}(\{\mathbf{0}\})$ , and  $\{\mathbf{0}\}$  is closed in  $\mathbb{R}^m$ . It follows that  $\text{Graph}(\varphi)$  is closed in  $X \times Y$  (see Corollary 1.15). ■

**Example** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be defined such that

$$f(x) = \begin{cases} \frac{1}{x} & \text{if } x > 0; \\ 0 & \text{if } x \leq 0. \end{cases}$$

Then the graph  $\text{Graph}(f)$  of the function  $f$  satisfies  $\text{Graph}(f) = Z \cup H$ , where

$$Z = \{(x, y) \in \mathbb{R}^2 : x \leq 0 \text{ and } y = 0\}$$

and

$$H = \{(x, y) \in \mathbb{R}^2 : x \geq 0 \text{ and } xy = 1\}.$$

Each of the sets  $Z$  and  $H$  is a closed set in  $\mathbb{R}^2$ . It follows that  $\text{Graph}(f)$  is a closed set in  $\mathbb{R}^2$ . However the function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is not continuous at 0.

**Lemma 2.8** Let  $X$  be a subset of  $n$ -dimensional Euclidean space  $\mathbb{R}^n$ , let  $S$  be a non-empty subset of  $X$ , and let

$$d(\mathbf{x}, S) = \inf\{|\mathbf{x} - \mathbf{s}| : \mathbf{s} \in S\}$$

for all  $\mathbf{x} \in X$ . Then the function sending  $\mathbf{x}$  to  $d(\mathbf{x}, S)$  for all  $\mathbf{x} \in X$  is a continuous function on  $X$ .

**Proof** Let  $f(\mathbf{x}) = d(\mathbf{x}, S) = \inf\{|\mathbf{x} - \mathbf{s}| : \mathbf{s} \in S\}$  for all  $\mathbf{x} \in X$ .

Let  $\mathbf{x}$  and  $\mathbf{x}'$  be points of  $X$ . It follows from the Triangle Inequality that

$$f(\mathbf{x}) \leq |\mathbf{x} - \mathbf{s}| \leq |\mathbf{x} - \mathbf{x}'| + |\mathbf{x}' - \mathbf{s}|$$

for all  $\mathbf{s} \in S$ , and therefore

$$|\mathbf{x}' - \mathbf{s}| \geq f(\mathbf{x}) - |\mathbf{x} - \mathbf{x}'|$$

for all  $\mathbf{s} \in S$ . Thus  $f(\mathbf{x}) - |\mathbf{x} - \mathbf{x}'|$  is a lower bound for the quantities  $|\mathbf{x}' - \mathbf{s}|$  as  $\mathbf{s}$  ranges over the set  $S$ , and therefore cannot exceed the greatest lower bound of these quantities. It follows that

$$f(\mathbf{x}') = \inf\{|\mathbf{x}' - \mathbf{s}| : \mathbf{s} \in S\} \geq f(\mathbf{x}) - |\mathbf{x} - \mathbf{x}'|,$$

and thus

$$f(\mathbf{x}) - f(\mathbf{x}') \leq |\mathbf{x} - \mathbf{x}'|.$$

Interchanging  $\mathbf{x}$  and  $\mathbf{x}'$ , it follows that

$$f(\mathbf{x}') - f(\mathbf{x}) \leq |\mathbf{x} - \mathbf{x}'|.$$

Thus

$$|f(\mathbf{x}) - f(\mathbf{x}')| \leq |\mathbf{x} - \mathbf{x}'|$$

for all  $\mathbf{x}, \mathbf{x}' \in X$ . It follows that the function  $f: X \rightarrow \mathbb{R}$  is continuous, as required. ■

The multidimensional Heine-Borel Theorem (Theorem 1.21) ensures that a subset of a Euclidean space is compact if and only if it is both closed and bounded.

**Proposition 2.9** *Let  $X$  be a subset of  $n$ -dimensional Euclidean space  $\mathbb{R}^n$ , let  $V$  be a subset of  $X$  that is open in  $X$ , and let  $K$  be a compact subset of  $\mathbb{R}^n$  satisfying  $K \subset V$ . Then there exists some positive real number  $\varepsilon$  with the property that  $B_X(K, \varepsilon) \subset V$ , where  $B_X(K, \varepsilon)$  denotes the subset of  $X$  consisting of those points of  $X$  that lie within a distance less than  $\varepsilon$  of some point of  $K$ .*

**Proof of Proposition 2.9 using the Extreme Value Theorem** Let  $f: K \rightarrow \mathbb{R}$  be defined such that

$$f(\mathbf{x}) = \inf\{|\mathbf{z} - \mathbf{x}| : \mathbf{z} \in X \setminus V\}.$$

for all  $\mathbf{x} \in K$ . It follows from Lemma 2.8 that the function  $f$  is continuous on  $K$ .

Now  $K \subset V$  and therefore, given any point  $\mathbf{x} \in K$ , there exists some positive real number  $\delta$  such that the open ball of radius  $\delta$  about the point  $\mathbf{x}$  is contained in  $V$ , and therefore  $f(\mathbf{x}) \geq \delta$ . It follows that  $f(\mathbf{x}) > 0$  for all  $\mathbf{x} \in K$ .

It follows from the Extreme Value Theorem for continuous real-valued functions on closed bounded subsets of Euclidean spaces (Theorem 1.17) that the function  $f: K \rightarrow \mathbb{R}$  attains its minimum value at some point of  $K$ . Let that minimum value be  $\varepsilon$ . Then  $f(\mathbf{x}) \geq \varepsilon > 0$  for all  $\mathbf{x} \in K$ , and therefore  $|\mathbf{x} - \mathbf{z}| \geq \varepsilon > 0$  for all  $\mathbf{x} \in K$  and  $\mathbf{z} \in X \setminus V$ . It follows that  $B_X(K, \varepsilon) \subset V$ , as required. ■

**Example** Let

$$F = \{(x, y) \in \mathbb{R}^2 : x \geq 0, y \geq 0 \text{ and } xy \geq 1\}.$$

and let

$$V = \{(x, y) \in \mathbb{R}^2 : y > 0\}.$$

Note that if  $(x, y) \in F$  then  $x > 0$  and  $y > 0$ , because  $xy = 1$ . It follows that  $F \subset V$ . Also  $F$  is a closed set in  $\mathbb{R}^2$  and  $V$  is an open set in  $\mathbb{R}^2$ . However  $F$  is not a compact subset of  $\mathbb{R}^2$  because  $F$  is not bounded.

We now show that there does not exist any positive real number  $\varepsilon$  with the property that  $B_{\mathbb{R}^2}(F, \varepsilon) \subset V$ , where  $B_{\mathbb{R}^2}(F, \varepsilon)$  denotes the set of points of  $\mathbb{R}^2$  that lie within a distance  $\varepsilon$  of some point of  $F$ . Indeed let some positive real number  $\varepsilon$  be given, let  $x$  be a positive real number satisfying  $x > 2\varepsilon^{-1}$ , and let  $y = x^{-1} - \frac{1}{2}\varepsilon$ . Then  $y < 0$ , and therefore  $(x, y) \notin V$ . But  $(x, y + \frac{1}{2}\varepsilon) \in F$ , and therefore  $(x, y) \in B_{\mathbb{R}^2}(F, \varepsilon)$ . This shows that there does not exist any positive real number  $\varepsilon$  for which  $B_{\mathbb{R}^2}(F, \varepsilon) \subset V$ .

**Proposition 2.10** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, let  $K$  be a non-empty compact subset of  $Y$ , and let  $U$  be a subset in  $X \times Y$  that is open in  $X \times Y$ . Let*

$$d_Y(\mathbf{y}, K) = \inf\{|\mathbf{y} - \mathbf{z}| : \mathbf{z} \in K\}$$

*for all  $\mathbf{y} \in Y$ . Let  $\mathbf{p}$  be a point of  $X$  with the property that  $(\mathbf{p}, \mathbf{z}) \in U$  for all  $\mathbf{z} \in K$ . Then there exists some positive number  $\delta$  such that  $(\mathbf{x}, \mathbf{y}) \in U$  for all  $\mathbf{x} \in X$  and  $\mathbf{y} \in Y$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$  and  $d(\mathbf{y}, K) < \delta$ .*

**Proof** Let

$$\tilde{K} = \{(\mathbf{p}, \mathbf{z}) : \mathbf{z} \in K\}.$$

Then  $\tilde{K}$  is a closed bounded subset of  $\mathbb{R}^n \times \mathbb{R}^m$ . It follows from Proposition 2.9 that there exists some positive real number  $\varepsilon$  such that

$$B_{X \times Y}(\tilde{K}, \varepsilon) \subset U$$

where  $B_{X \times Y}(\tilde{K}, \varepsilon)$  denotes that subset of  $X \times Y$  consisting of those points  $(\mathbf{x}, \mathbf{y})$  of  $X \times Y$  that lie within a distance  $\varepsilon$  of a point of  $\tilde{K}$ . Now a point

$(\mathbf{x}, \mathbf{y})$  of  $X \times Y$  belongs to  $B_{X \times Y}(\tilde{K}, \varepsilon)$  if and only if there exists some point  $\mathbf{z}$  of  $K$  for which

$$|\mathbf{x} - \mathbf{p}|^2 + |\mathbf{y} - \mathbf{z}|^2 < \varepsilon^2.$$

Let  $\delta = \varepsilon/\sqrt{2}$ . If  $\mathbf{x} \in X$  and  $\mathbf{y} \in Y$  satisfy  $|\mathbf{x} - \mathbf{p}| < \delta$  and  $d_Y(\mathbf{y}, K) < \delta$  then there exists some point  $\mathbf{z}$  of  $K$  for which  $|\mathbf{y} - \mathbf{z}| < \delta$ . But then

$$|\mathbf{x} - \mathbf{p}|^2 + |\mathbf{y} - \mathbf{z}|^2 < 2\delta^2 = \varepsilon^2,$$

and therefore  $(\mathbf{x}, \mathbf{y}) \in U$ , as required.  $\blacksquare$

**Proposition 2.11** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Suppose that  $\Phi(\mathbf{x})$  is closed in  $Y$  for every  $\mathbf{x} \in X$ . Suppose also that  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous. Then the graph  $\text{Graph}(\Phi)$  of  $\Phi: X \rightrightarrows Y$  is closed in  $X \times Y$ .*

**Proof** Let  $(\mathbf{p}, \mathbf{q})$  be a point of the complement  $X \times Y \setminus \text{Graph}(\Phi)$  of the graph  $\text{Graph}(\Phi)$  of  $\Phi$  in  $X \times Y$ . Then  $\Phi(\mathbf{p})$  is closed in  $Y$  and  $\mathbf{q} \notin \Phi(\mathbf{p})$ . It follows that there exists some positive real number  $\delta_Y$  such that  $|\mathbf{y} - \mathbf{q}| > \delta_Y$  for all  $\mathbf{y} \in \Phi(\mathbf{p})$ .

Let

$$V = \{\mathbf{y} \in Y : |\mathbf{y} - \mathbf{q}| > \delta_Y\}$$

and

$$W = \{\mathbf{x} \in X : \Phi(\mathbf{x}) \subset V\}.$$

Then  $V$  is open in  $Y$  and  $\Phi(\mathbf{p}) \subset V$ . Now the correspondence  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous. It therefore follows from the definition of upper hemicontinuity that the subset  $W$  of  $X$  is open in  $X$ . Moreover  $\mathbf{p} \in W$ . It follows that there exists some positive real number  $\delta_X$  such that  $\mathbf{x} \in W$  for all points  $\mathbf{x}$  of  $X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta_X$ . Then  $\Phi(\mathbf{x}) \subset V$  for all points  $\mathbf{x}$  of  $X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta_X$ . Let  $\delta$  be the minimum of  $\delta_X$  and  $\delta_Y$ , and let  $(\mathbf{x}, \mathbf{y})$  be a point of  $X \times Y$  whose distance from the point  $(\mathbf{p}, \mathbf{q})$  is less than  $\delta$ . Then  $|\mathbf{x} - \mathbf{p}| < \delta_X$  and therefore  $\Phi(\mathbf{x}) \subset V$ . Also  $|\mathbf{y} - \mathbf{q}| < \delta_Y$ , and therefore  $\mathbf{y} \notin V$ . It follows that  $\mathbf{y} \notin \Phi(\mathbf{x})$ , and therefore  $(\mathbf{x}, \mathbf{y}) \notin \text{Graph}(\Phi)$ . We conclude from this that the complement of  $\text{Graph}(\Phi)$  is open in  $X \times Y$ . It follows that  $\text{Graph}(\Phi)$  itself is closed in  $X \times Y$ , as required.  $\blacksquare$

**Proposition 2.12** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Suppose that the graph  $\text{Graph}(\Phi)$  of the correspondence  $\Phi$  is closed in  $X \times Y$ . Suppose also that  $Y$  is a compact subset of  $\mathbb{R}^m$ . Then the correspondence  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous.*

**Proof of Proposition 2.12 using Proposition 2.10** Let  $\mathbf{p}$  be a point of  $X$ , let  $V$  be an open set satisfying  $\Phi(\mathbf{p}) \subset V$ , and let  $K = Y \setminus V$ . The compact set  $Y$  is closed and bounded in  $\mathbb{R}^m$ . Also  $K$  is closed in  $Y$ . It follows that  $K$  is a closed bounded subset of  $\mathbb{R}^m$  (see Lemma 1.16). Let  $U$  be the complement of  $\text{Graph}(\Phi)$  in  $X \times Y$ . Then  $U$  is open in  $X \times Y$ , because  $\text{Graph}(\Phi)$  is closed in  $X \times Y$ . Also  $(\mathbf{p}, \mathbf{y}) \in U$  for all  $\mathbf{y} \in K$ , because  $\Phi(\mathbf{p}) \cap K = \emptyset$ . It follows from Proposition 2.10 that there exists some positive number  $\delta$  such that  $(\mathbf{x}, \mathbf{y}) \in U$  for all  $\mathbf{x} \in X$  and  $\mathbf{y} \in K$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . Thus if  $\mathbf{x} \in X$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta$  then  $\mathbf{y} \notin \Phi(\mathbf{x})$  for all  $\mathbf{y} \in K$ , and therefore  $\Phi(\mathbf{x}) \subset V$ , where  $V = Y \setminus K$ . Thus the correspondence  $\Phi$  is upper hemicontinuous at  $\mathbf{p}$ , as required. ■

**Corollary 2.13** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\varphi: X \rightarrow Y$  be a function from  $X$  to  $Y$ . Suppose that the graph  $\text{Graph}(\varphi)$  of the function  $\varphi$  is closed in  $X \times Y$ . Suppose also that  $Y$  is a compact subset of  $\mathbb{R}^m$ . Then the function  $\varphi: X \rightarrow Y$  is continuous.*

**Proof** Let  $\Phi: X \rightrightarrows Y$  be the correspondence defined such that  $\Phi(\mathbf{x}) = \{\varphi(\mathbf{x})\}$  for all  $\mathbf{x} \in X$ . Then  $\text{Graph}(\Phi) = \text{Graph}(\varphi)$ , and therefore  $\text{Graph}(\Phi)$  is closed in  $X \times Y$ . The subset  $Y$  of  $\mathbb{R}^m$  is compact. It therefore follows from Proposition 2.12 that the correspondence  $\Phi$  is upper hemicontinuous. It then follows from Lemma 2.3 that the function  $\varphi: X \rightarrow Y$  is continuous, as required. ■

## 2.3 Compact-Valued Upper Hemicontinuous Correspondences

**Lemma 2.14** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Suppose that  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous. Then*

$$\{\mathbf{x} \in X : \Phi(\mathbf{x}) \neq \emptyset\}$$

*is closed in  $X$ .*

**Proof** Given any open set  $V$  in  $Y$ , let

$$\Phi^+(V) = \{\mathbf{x} \in X : \Phi(\mathbf{x}) \subset V\}.$$

It follows from the upper hemicontinuity of  $\Phi$  that  $\Phi^+(V)$  is open in  $X$  for all open sets  $V$  in  $Y$  (see Lemma 2.1). Now the empty set  $\emptyset$  is open in  $Y$ . It follows that  $\Phi^+(\emptyset)$  is open in  $X$ . But

$$\Phi^+(\emptyset) = \{\mathbf{x} \in X : \Phi(\mathbf{x}) \subset \emptyset\} = \{\mathbf{x} \in X : \Phi(\mathbf{x}) = \emptyset\}.$$

It follows that the set of point  $\mathbf{x}$  in  $X$  for which  $\Phi(\mathbf{x}) = \emptyset$  is open in  $X$ , and therefore the set of points  $\mathbf{x} \in X$  for which  $\Phi(\mathbf{x}) \neq \emptyset$  is closed in  $X$ , as required. ■

Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Given any subset  $S$  of  $X$ , we define the *image*  $\Phi(S)$  of  $S$  under the correspondence  $\Phi$  to be the subset of  $Y$  defined such that

$$\Phi(S) = \bigcup_{\mathbf{x} \in S} \Phi(\mathbf{x})$$

**Lemma 2.15** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$  that is compact-valued and upper hemicontinuous. Let  $K$  be a compact subset of  $X$ . Then  $\Phi(K)$  is a compact subset of  $Y$ .*

**Proof** Let  $\mathcal{V}$  be collection of open sets in  $Y$  that covers  $\Phi(K)$ . Given any point  $\mathbf{p}$  of  $K$ , there exists a finite subcollection  $\mathcal{W}_{\mathbf{p}}$  of  $\mathcal{V}$  that covers the compact set  $\Phi(\mathbf{p})$ . Let  $U_{\mathbf{p}}$  be the union of the open sets belonging to this subcollection  $\mathcal{W}_{\mathbf{p}}$ . Then  $\Phi(\mathbf{p}) \subset U_{\mathbf{p}}$ . Now it follows from the upper hemicontinuity of  $\Phi: X \rightrightarrows Y$  that there exists an open set  $N_{\mathbf{p}}$  in  $X$  such that  $\Phi(\mathbf{x}) \subset U_{\mathbf{p}}$  for all  $\mathbf{x} \in N_{\mathbf{p}}$ . Moreover, given any  $\mathbf{p} \in K$ , the finite collection  $\mathcal{W}_{\mathbf{p}}$  of open sets in  $Y$  covers  $\Phi(N_{\mathbf{p}})$ . It then follows from the compactness of  $K$  that there exist points

$$\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$$

of  $K$  such that

$$K \subset N_{\mathbf{p}_1} \cup N_{\mathbf{p}_2} \cup \dots \cup N_{\mathbf{p}_k}.$$

Let

$$\mathcal{W} = \mathcal{W}_{\mathbf{p}_1} \cup \mathcal{W}_{\mathbf{p}_2} \cup \dots \cup \mathcal{W}_{\mathbf{p}_k}.$$

Then  $\mathcal{W}$  is a finite subcollection of  $\mathcal{V}$  that covers  $\Phi(K)$ . The result follows. ■

**Proposition 2.16** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a compact-valued correspondence from  $X$  to  $Y$ . Let  $\mathbf{p}$  be a point of  $X$  for which  $\Phi(\mathbf{p})$  is non-empty. Then the correspondence  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at  $\mathbf{p}$  if and only if, given any positive real number  $\varepsilon$ , there exists some positive real number  $\delta$  such that*

$$\Phi(\mathbf{x}) \subset B_Y(\Phi(\mathbf{p}), \varepsilon)$$

*for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ , where  $B_Y(\Phi(\mathbf{p}), \varepsilon)$  denotes the subset of  $Y$  consisting of all points of  $Y$  that lie within a distance  $\varepsilon$  of some point of  $\Phi(\mathbf{p})$ .*



**Proof** Let  $\Phi: X \rightrightarrows Y$  is a compact-valued correspondence, and let  $\mathbf{p}$  be a point of  $X$  for which  $\Phi(\mathbf{p}) \neq \emptyset$ .

First suppose that, given any positive real number  $\varepsilon$ , there exists some positive real number  $\delta$  such that

$$\Phi(\mathbf{x}) \subset B_Y(\Phi(\mathbf{p}), \varepsilon)$$

for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . We must prove that  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at  $\mathbf{p}$ .

Let  $V$  be an open set in  $Y$  that satisfies  $\Phi(\mathbf{p}) \subset V$ . Now  $\Phi(\mathbf{p})$  is a compact subset of  $Y$ , because  $\Phi: X \rightarrow Y$  is compact-valued. It follows that there exists some positive real number  $\varepsilon$  such that  $B_Y(\Phi(\mathbf{p}), \varepsilon) \subset V$  (see Proposition 2.9). There then exists some positive number  $\delta$  such that

$$\Phi(\mathbf{x}) \subset B_Y(\Phi(\mathbf{p}), \varepsilon) \subset V$$

whenever  $|\mathbf{x} - \mathbf{p}| < \delta$ . Thus  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at  $\mathbf{p}$ .

Conversely suppose that the correspondence  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at the point  $\mathbf{p}$ . Now  $\Phi(\mathbf{p})$  is a non-empty subset of  $Y$ . Let some positive number  $\varepsilon$  be given. Then  $B_Y(\Phi(\mathbf{p}), \varepsilon)$  is open in  $Y$  and  $\Phi(\mathbf{p}) \subset B_Y(\Phi(\mathbf{p}), \varepsilon)$ . It follows from the upper hemicontinuity of  $\Phi$  at  $\mathbf{p}$  that there exists some positive number  $\delta$  such that  $\Phi(\mathbf{x}) \subset B_Y(\Phi(\mathbf{p}), \varepsilon)$  whenever  $|\mathbf{x} - \mathbf{p}| < \delta$ . The result follows. ■

**Proposition 2.17** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Then the correspondence is both compact-valued and upper hemicontinuous at a point  $\mathbf{p} \in X$  if and only if, given any infinite sequences*

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

and

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

in  $X$  and  $Y$  respectively, where  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$  and  $\lim_{j \rightarrow +\infty} \mathbf{x}_j = \mathbf{p}$ , there exists a subsequence of

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

which converges to a point of  $\Phi(\mathbf{p})$ .

**Proof** Throughout this proof, let us say that the correspondence  $\Phi$  satisfies the *constrained convergent subsequence criterion* if (and only if), given any infinite sequences

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

and

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

in  $X$  and  $Y$  respectively, where  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$  and  $\lim_{j \rightarrow +\infty} \mathbf{x}_j = \mathbf{p}$ , there exists a subsequence of

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

which converges to a point of  $\Phi(\mathbf{p})$ .

We must prove that the correspondence  $\Phi: X \rightrightarrows Y$  satisfies the constrained convergent subsequence criterion if and only if it is compact-valued and upper hemicontinuous.

Suppose first that the correspondence  $\Phi: X \rightrightarrows Y$  satisfies the constrained convergent subsequence criterion. Applying this criterion when  $\mathbf{x}_j = \mathbf{p}$  for all positive integers  $j$ , we conclude that every infinite sequence  $(\mathbf{y}_j : j \in \mathbb{N})$  of points of  $\Phi(\mathbf{p})$  has a convergent subsequence, and therefore  $\Phi(\mathbf{p})$  is compact.

Let

$$D = \{\mathbf{x} \in X : \Phi(\mathbf{x}) \neq \emptyset\}.$$

We show that  $D$  is closed in  $X$ . Let

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

be a sequence of points of  $D$  converging to some point of  $\mathbf{p}$  of  $X$ . Then  $\Phi(\mathbf{x}_j)$  is non-empty for all positive integers  $j$ , and therefore there exists an infinite sequence

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

of points of  $Y$  such that  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$ . The constrained convergent subsequence criterion ensures that this infinite sequence in  $Y$  must have a subsequence that converges to some point of  $\Phi(\mathbf{p})$ . It follows that  $\Phi(\mathbf{p})$  is non-empty, and thus  $\mathbf{p} \in D$ .

Let  $\mathbf{p}$  be a point of the complement of  $D$ . Then  $\Phi(\mathbf{p}) = \emptyset$ . There then exists  $\delta > 0$  such that  $\Phi(\mathbf{x}) = \emptyset$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . But then  $\Phi(\mathbf{x}) \subset V$  for all open sets  $V$  in  $Y$ . It follows that the correspondence  $\Phi$  is upper hemicontinuous at those points  $\mathbf{p}$  for which  $\Phi(\mathbf{p}) = \emptyset$ .

Now consider the situation in which  $\Phi: X \rightrightarrows Y$  satisfies the constrained convergent subsequence criterion and  $\mathbf{p}$  is some point of  $X$  for which  $\Phi(\mathbf{p}) \neq \emptyset$ . Let  $K = \Phi(\mathbf{p})$ . Then  $K$  is a compact non-empty subset of  $Y$ . Let  $V$  be an open set in  $Y$  that satisfies  $\Phi(\mathbf{p}) \subset V$ . Suppose that there did not exist any positive real number  $\delta$  with the property that  $\Phi(\mathbf{x}) \subset V$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . It would then follow that there would exist infinite sequences

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

and

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

in  $X$  and  $Y$  respectively for which  $|\mathbf{x}_j - \mathbf{p}| < 1/j$ ,  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  and  $\mathbf{y}_j \notin V$ . Then  $\lim_{j \rightarrow +\infty} \mathbf{x}_j = \mathbf{p}$ , and thus the constrained convergent subsequence criterion satisfied by the correspondence  $\Phi$  would ensure the existence of a subsequence

$$\mathbf{y}_{k_1}, \mathbf{y}_{k_2}, \mathbf{y}_{k_3}, \dots$$

of  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$  converging to some point  $\mathbf{q}$  of  $\Phi(\mathbf{p})$ . But then  $\mathbf{q} \notin V$ , because  $\mathbf{y}_{k_j} \notin V$  for all positive integers  $j$ , and the complement  $Y \setminus V$  of  $V$  is closed in  $Y$ . But  $\Phi(\mathbf{p}) \subset V$ , and  $\mathbf{q} \in \Phi(\mathbf{p})$ , and therefore  $\mathbf{q} \in V$ . Thus a contradiction would arise were there not to exist a positive real number  $\delta$  with the property that  $\Phi(\mathbf{x}) \subset V$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . Thus such a real number  $\delta$  must exist, and thus the constrained convergent subsequence criterion ensures that the correspondence  $\Phi: X \rightrightarrows Y$  is upper hemicontinuous at  $\mathbf{p}$ .

It remains to show that any compact-valued upper hemicontinuous correspondence  $\Phi: X \rightrightarrows Y$  satisfies the constrained convergent subsequence criterion. Let  $\Phi: X \rightrightarrows Y$  be compact-valued and upper hemicontinuous. It follows from Lemma 2.14 that

$$\{\mathbf{x} \in X : \Phi(\mathbf{x}) \neq \emptyset\}$$

is closed in  $X$ .

Let

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

and

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

be infinite sequences in  $X$  and  $Y$  respectively, where  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$  and  $\lim_{j \rightarrow +\infty} \mathbf{x}_j = \mathbf{p}$ . Then  $\Phi(\mathbf{p})$  is non-empty, because

$$\{\mathbf{x} \in X : \Phi(\mathbf{x}) \neq \emptyset\}$$

is closed in  $X$  (see Lemma 2.14). Let  $K = \Phi(\mathbf{p})$ . Then  $K$  is compact, because  $\Phi: X \rightrightarrows Y$  is compact-valued by assumption. For each integer  $j$  let  $d(\mathbf{y}_j, K)$  denote the greatest lower bound on the distances from  $\mathbf{y}_j$  to points of  $K$ . There then exists an infinite sequence

$$\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \dots$$

of points of  $K$  such that  $|\mathbf{y}_j - \mathbf{z}_j| \leq 2d(\mathbf{y}_j, K)$  for all positive integers  $j$ . (Indeed if  $d(\mathbf{y}_j, K) = 0$  then  $\mathbf{y}_j \in K$ , because the compact set  $K$  is closed,

and in that case we can take  $\mathbf{z}_j = \mathbf{y}_j$ . Otherwise  $2d(\mathbf{y}_j, K)$  is strictly greater than the greatest lower bound on the distances from  $\mathbf{y}_j$  to points of  $K$ , and we can therefore find  $\mathbf{z}_j \in K$  with  $|\mathbf{y}_j - \mathbf{z}_j| \leq 2d(\mathbf{y}_j, K)$ .)

Now the upper hemicontinuity of  $\Phi: X \rightrightarrows Y$  ensures that  $d(\mathbf{y}_j, K) \rightarrow 0$  as  $j \rightarrow +\infty$ . Indeed, given any positive real number  $\varepsilon$ , the set  $B_Y(K, \varepsilon)$  of points of  $Y$  that lie within a distance  $\varepsilon$  of a point of  $K$  is an open set containing  $\Phi(\mathbf{p})$ . It follows from the upper hemicontinuity of  $\Phi$  that there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \subset B_Y(K, \varepsilon)$  whenever  $|\mathbf{x} - \mathbf{p}| < \delta$ . Now  $\mathbf{x}_j \rightarrow \mathbf{p}$  as  $j \rightarrow +\infty$ . It follows that there exists some positive integer  $N$  such that  $|\mathbf{x}_j - \mathbf{p}| < \delta$  whenever  $j \geq N$ . But then  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  and therefore  $d(\mathbf{y}_j, K) < \varepsilon$  whenever  $j \geq N$ . Now the compactness of  $K$  ensures that the infinite sequence

$$\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \dots$$

of points of  $K$  has a subsequence

$$\mathbf{z}_{k_1}, \mathbf{z}_{k_2}, \mathbf{z}_{k_3}, \dots$$

that converges to some point  $\mathbf{q}$  of  $K$ . Now  $|\mathbf{y}_j - \mathbf{z}_j| \leq 2d(\mathbf{y}_j, K)$  for all positive integers  $j$ , and  $d(\mathbf{y}_j, K) \rightarrow 0$  as  $j \rightarrow +\infty$ . It follows that  $\mathbf{y}_{k_j} \rightarrow \mathbf{q}$  as  $j \rightarrow +\infty$ . Moreover  $\mathbf{q} \in \Phi(\mathbf{p})$ . We have therefore verified that the constrained convergent subsequence criterion is satisfied by any correspondence  $\Phi: X \rightrightarrows Y$  that is compact-valued and upper hemicontinuous. This completes the proof. ■

**Proposition 2.18** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$  that is both upper hemicontinuous and compact-valued. Let  $U$  be an open set in  $X \times Y$ . Then*

$$\{\mathbf{x} \in X : (\mathbf{x}, \mathbf{y}) \in U \text{ for all } \mathbf{y} \in \Phi(\mathbf{x})\}$$

*is open in  $X$ .*

**Proof of Proposition 2.18 using Proposition 2.17** Let

$$W = \{\mathbf{x} \in X : (\mathbf{x}, \mathbf{y}) \in U \text{ for all } \mathbf{y} \in \Phi(\mathbf{x})\},$$

and let  $\mathbf{p} \in W$ . Suppose that there did not exist any strictly positive real number  $\delta$  with the property that  $\mathbf{x} \in W$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . Then, given any positive real number  $\delta$ , there would exist points  $\mathbf{x}$  of  $X$  and  $\mathbf{y}$  of  $Y$  such that  $|\mathbf{x} - \mathbf{p}| < \delta$ ,  $\mathbf{y} \in \Phi(\mathbf{x})$  and  $(\mathbf{x}, \mathbf{y}) \notin U$ . Therefore there would exist infinite sequences

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

and

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

in  $X$  and  $Y$  respectively such that  $\mathbf{x}_j \rightarrow \mathbf{p}$  as  $j \rightarrow +\infty$  and  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  and  $(\mathbf{x}_j, \mathbf{y}_j) \notin U$  for all positive integers  $j$ . The correspondence  $\Phi: X \rightrightarrows Y$  is compact-valued and upper hemicontinuous. Proposition 2.17 would therefore ensure the existence of a subsequence

$$\mathbf{y}_{k_1}, \mathbf{y}_{k_2}, \mathbf{y}_{k_3}, \dots$$

of  $Y$  converging to some point  $\mathbf{q}$  of  $\Phi(\mathbf{p})$ . Now the complement of  $U$  in  $X \times Y$  is closed in  $X \times Y$ , because  $U$  is open in  $X \times Y$  and  $(\mathbf{x}_j, \mathbf{y}_j) \notin U$ . It would therefore follow that  $(\mathbf{p}, \mathbf{q}) \notin U$  (see Proposition 2.6). But this gives rise to a contradiction, because  $\mathbf{q} \in \Phi(\mathbf{p})$  and  $(\mathbf{p}, \mathbf{y}) \in U$  for all  $\mathbf{y} \in \Phi(\mathbf{p})$ . In order to avoid the contradiction, there must exist some positive real number  $\delta$  with the property that with the property that  $(\mathbf{x}, \mathbf{y}) \in U$  for all  $\mathbf{x} \in X$  and  $\mathbf{y} \in Y$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$  and  $\mathbf{y} \in \Phi(\mathbf{x})$ . The result follows. ■

**Remark** It should be noted that other results proved in this section do not necessarily generalize to correspondences  $\Phi: X \rightrightarrows Y$  mapping the topological space  $X$  into an arbitrary topological space  $Y$ . For example all closed-valued upper hemicontinuous correspondences between metric spaces have closed graphs. The appropriate generalization of this result states that any closed-valued upper hemicontinuous correspondence  $\Phi: X \rightrightarrows Y$  from a topological space  $X$  to a regular topological space  $Y$  has a closed graph. To interpret this, one needs to know the definition of what is meant by saying that a topological space is *regular*. A topological space  $Y$  is said to be *regular* if, given any closed subset  $F$  of  $Y$ , and given any point  $p$  of the complement  $Y \setminus F$  of  $F$ , there exist open sets  $V$  and  $W$  in  $Y$  such that  $F \subset V$ ,  $p \in W$  and  $V \cap W = \emptyset$ . Metric spaces are regular. Also compact Hausdorff spaces are regular.

## 2.4 A Criterion characterizing Lower Hemicontinuity

**Proposition 2.19** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively. A correspondence  $\Phi: X \rightrightarrows Y$  is lower hemicontinuous at a point  $\mathbf{p}$  of  $X$  if and only if given any infinite sequence*

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

*in  $X$  for which  $\lim_{j \rightarrow +\infty} \mathbf{x}_j = \mathbf{p}$  and given any point  $\mathbf{q}$  of  $\Phi(\mathbf{p})$ , there exists an infinite sequence*

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

of points of  $Y$  such that  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$  and  $\lim_{j \rightarrow +\infty} \mathbf{y}_j = \mathbf{q}$ .

**Proof** First suppose that  $\Phi: X \rightarrow Y$  is lower hemicontinuous at some point  $\mathbf{p}$  of  $X$ . Let  $\mathbf{q} \in \Phi(\mathbf{p})$ , and let some positive number  $\varepsilon$  be given. Then the open ball  $B_Y(\mathbf{q}, \varepsilon)$  in  $Y$  of radius  $\varepsilon$  centred on the point  $\mathbf{q}$  is an open set in  $Y$ . It follows from the lower hemicontinuity of  $\Phi: X \rightarrow Y$  that there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \cap B_Y(\mathbf{q}, \varepsilon)$  is non-empty whenever  $|\mathbf{x} - \mathbf{p}| < \delta$ . Then, given any point  $\mathbf{x}$  of  $X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$  there exists some  $\mathbf{y} \in \Phi(\mathbf{x})$  that satisfies  $|\mathbf{y} - \mathbf{q}| < \varepsilon$ . In particular, given any positive integer  $s$ , there exists some positive integer  $\delta_s$  such that, given any point  $\mathbf{x}$  of  $X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta_s$ , there exists some  $\mathbf{y} \in \Phi(\mathbf{x})$  that satisfies  $|\mathbf{y} - \mathbf{q}| < 1/s$ .

Now  $\mathbf{x}_j \rightarrow \mathbf{p}$  as  $j \rightarrow +\infty$ . It follows that there exist positive integers  $k(1), k(2), k(3), \dots$ , where

$$k(1) < k(2) < k(3) < \dots$$

such that  $|\mathbf{x}_j - \mathbf{p}| < \delta_s$  for all positive integers  $j$  satisfying  $j \geq k(s)$ . There then exists an infinite sequence

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

such that  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$  and  $|\mathbf{y}_j - \mathbf{q}| < 1/s$  for all positive integers  $j$  and  $s$  satisfying  $k(s) \leq j < k(s+1)$ . Then  $\lim_{j \rightarrow +\infty} \mathbf{y}_j = \mathbf{q}$ .

We have thus shown that if  $\Phi: X \rightarrow Y$  is lower hemicontinuous at the point  $\mathbf{p}$ , if  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$  is a sequence in  $X$  converging to the point  $\mathbf{p}$ , and if  $\mathbf{q} \in \Phi(\mathbf{p})$ , then there exists an infinite sequence  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$  in  $Y$  such that  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integer  $j$  and  $\lim_{j \rightarrow +\infty} \mathbf{y}_j = \mathbf{q}$ .

Next suppose that the correspondence  $\Phi: X \rightrightarrows Y$  is not lower hemicontinuous at  $\mathbf{p}$ . Then there exists an open set  $V$  in  $Y$  such that  $\Phi(\mathbf{p}) \cap V$  is non-empty but there does not exist any positive real number  $\delta$  with the property that  $\Phi(\mathbf{x}) \cap V \neq \emptyset$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{p} - \mathbf{x}| < \delta$ . Let  $\mathbf{q} \in \Phi(\mathbf{p}) \cap V$ . There then exists an infinite sequence

$$\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$$

converging to the point  $\mathbf{p}$  with the property that  $\Phi(\mathbf{x}_j) \cap V = \emptyset$  for all positive integers  $j$ . It is not then possible to construct an infinite sequence

$$\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots$$

such that  $\mathbf{y}_j \in \Phi(\mathbf{x}_j)$  for all positive integers  $j$  and  $\lim_{j \rightarrow +\infty} \mathbf{y}_j = \mathbf{q}$ . The result follows. ■

## 2.5 Intersections of Correspondences

Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  and  $\Psi: X \rightarrow Y$  be correspondences between  $X$  and  $Y$ . The *intersection*  $\Phi \cap \Psi$  of the correspondences  $\Phi$  and  $\Psi$  is defined such that

$$(\Phi \cap \Psi)(\mathbf{x}) = \Phi(\mathbf{x}) \cap \Psi(\mathbf{x})$$

for all  $\mathbf{x} \in X$ .

**Proposition 2.20** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, let  $\Phi: X \rightrightarrows Y$  and  $\Psi: X \rightarrow Y$  be correspondences from  $X$  to  $Y$ , where the correspondence  $\Phi: X \rightrightarrows Y$  is compact-valued and upper hemicontinuous and the correspondence  $\Psi: X \rightarrow Y$  has closed graph. Let  $\Phi \cap \Psi: X \rightrightarrows Y$  be the correspondence defined such that*

$$(\Phi \cap \Psi)(\mathbf{x}) = \Phi(\mathbf{x}) \cap \Psi(\mathbf{x})$$

*for all  $\mathbf{x} \in X$ . Then the correspondence  $\Phi \cap \Psi: X \rightrightarrows Y$  is compact-valued and upper hemicontinuous.*

**Proof** Let

$$W = \{(\mathbf{x}, \mathbf{y}) \in X \times Y : \mathbf{y} \notin \Psi(\mathbf{x})\}.$$

Then  $W$  is the complement of the graph  $\text{Graph}(\Psi)$  of  $\Psi$  in  $X \times Y$ . The graph of  $\Psi$  is closed in  $X \times Y$ , by assumption. It follows that  $W$  is open in  $X \times Y$ .

Let  $\mathbf{x} \in X$ . The subset  $\Psi(\mathbf{x})$  of  $Y$  is closed in  $Y$ , because the graph of the correspondence  $\Psi$  is closed. It follows from the compactness of  $\Phi(\mathbf{x})$  that  $\Phi(\mathbf{x}) \cap \Psi(\mathbf{x})$  is a closed subset of the compact set  $\Phi(\mathbf{x})$ , and must therefore be compact. Thus the correspondence  $\Phi \cap \Psi$  is compact-valued.

Now let  $\mathbf{p}$  be a point of  $X$ , and let  $V$  be any open set in  $Y$  for which  $\Phi(\mathbf{p}) \cap \Psi(\mathbf{p}) \subset V$ . In order to prove that  $\Phi \cap \Psi$  is upper hemicontinuous we must show that there exists some positive real number  $\delta$  such that  $\Phi(\mathbf{x}) \cap \Psi(\mathbf{x}) \subset V$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$ . Let

$$U = \{(\mathbf{x}, \mathbf{y}) \in X \times Y : \text{either } \mathbf{y} \in V \text{ or else } \mathbf{y} \notin \Psi(\mathbf{x})\}.$$

Then  $U$  is the union of the subsets  $X \times V$  and  $W$  of  $X \times Y$ , where both these subsets are open in  $X \times Y$ . It follows that  $U$  is open in  $X \times Y$ . Moreover if  $\mathbf{y} \in \Phi(\mathbf{p})$  then either  $\mathbf{y} \in \Phi(\mathbf{p}) \cap \Psi(\mathbf{p})$ , in which case  $\mathbf{y} \in V$ , or else  $\mathbf{y} \notin \Psi(\mathbf{p})$ . It follows that  $(\mathbf{p}, \mathbf{y}) \in U$  for all  $\mathbf{y} \in \Phi(\mathbf{p})$ .

Now it follows from Proposition 2.18 that

$$\{\mathbf{x} \in X : (\mathbf{x}, \mathbf{y}) \in U \text{ for all } \mathbf{y} \in \Phi(\mathbf{x})\}$$

is open in  $X$ . Therefore there exists some positive real number  $\delta$  such that  $(\mathbf{x}, \mathbf{y}) \in U$  for all  $(\mathbf{x}, \mathbf{y}) \in X \times Y$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta$  and  $\mathbf{y} \in \Phi(\mathbf{x})$ . Now if  $(\mathbf{x}, \mathbf{y})$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta$  and  $\mathbf{y} \in \Phi(\mathbf{x}) \cap \Psi(\mathbf{x})$  then  $(\mathbf{x}, \mathbf{y}) \in U$  but  $(\mathbf{x}, \mathbf{y}) \notin W$ . It follows from the definition of  $U$  that  $\mathbf{y} \in V$ . Thus  $\Phi(\mathbf{x}) \cap \Psi(\mathbf{x}) \subset V$  whenever  $|\mathbf{x} - \mathbf{p}| < \delta$ . The result follows. ■

## 2.6 Berge's Maximum Theorem

**Lemma 2.21** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$  that is both upper hemicontinuous and compact-valued. Let  $f: X \times Y \rightarrow \mathbb{R}$  be a continuous real-valued function on  $X \times Y$ , and let  $c$  be a real number. Then*

$$\{\mathbf{x} \in X : f(\mathbf{x}, \mathbf{y}) < c \text{ for all } \mathbf{y} \in \Phi(\mathbf{x})\}$$

*is open in  $X$ .*

**Proof** Let

$$U = \{(\mathbf{x}, \mathbf{y}) \in X \times Y : f(\mathbf{x}, \mathbf{y}) < c\}.$$

It follows from the continuity of the function  $f$  that  $U$  is open in  $X \times Y$ . It then follows from Proposition 2.18 that

$$\{\mathbf{x} \in X : (\mathbf{x}, \mathbf{y}) \in U \text{ for all } \mathbf{y} \in \Phi(\mathbf{x})\}$$

is open in  $X$ . The result follows. ■

**Lemma 2.22** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$  that is lower hemicontinuous. Let  $f: X \times Y \rightarrow \mathbb{R}$  be a continuous real-valued function on  $X \times Y$ , and let  $c$  be a real number. Then*

$$\{\mathbf{x} \in X : \text{there exists } \mathbf{y} \in \Phi(\mathbf{x}) \text{ for which } f(\mathbf{x}, \mathbf{y}) > c\}$$

*is open in  $X$ .*

**Proof** Let

$$U = \{(\mathbf{x}, \mathbf{y}) \in X \times Y : f(\mathbf{x}, \mathbf{y}) > c\},$$

and let

$$W = \{\mathbf{x} \in X : \text{there exists } \mathbf{y} \in \Phi(\mathbf{x}) \text{ for which } f(\mathbf{x}, \mathbf{y}) > c\},$$

Let  $\mathbf{p} \in W$ . Then there exists  $\mathbf{y} \in \Phi(\mathbf{p})$  for which  $(\mathbf{p}, \mathbf{y}) \in U$ . There then exist subsets  $W_X$  of  $X$  and  $W_Y$  of  $Y$ , where  $W_X$  is open in  $X$  and  $W_Y$  is



open in  $Y$ , such that  $\mathbf{p} \in W_X$ ,  $\mathbf{y} \in W_Y$  and  $W_X \times W_Y \subset U$  (see Lemma 2.5). There then exists some positive real number  $\delta_1$  such that  $\mathbf{x} \in W_X$  whenever  $|\mathbf{x} - \mathbf{p}| < \delta_1$ .

Now  $\Phi(\mathbf{p}) \cap W_Y \neq \emptyset$ , because  $\mathbf{y} \in \Phi(\mathbf{p}) \cap W_Y$ . It follows from the lower hemicontinuity of the correspondence  $\Phi$  that there exists some positive real number  $\delta_2$  such that  $\Phi(\mathbf{x}) \cap W_Y \neq \emptyset$  whenever  $|\mathbf{x} - \mathbf{p}| < \delta_2$ . Let  $\delta$  be the minimum of  $\delta_1$  and  $\delta_2$ . If  $\mathbf{x} \in X$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta$  then there exists  $\mathbf{y} \in \Phi(\mathbf{x})$  for which  $\mathbf{y} \in W_Y$ . But then  $(\mathbf{x}, \mathbf{y}) \in W_X \times W_Y$  and therefore  $(\mathbf{x}, \mathbf{y}) \in U$ , and thus  $f(\mathbf{x}, \mathbf{y}) > c$ . The result follows. ■

**Theorem 2.23 (Berge's Maximum Theorem)** *Let  $X$  and  $Y$  be subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively, let  $f: X \times Y \rightarrow \mathbb{R}$  be a continuous real-valued function on  $X \times Y$ , and let  $\Phi: X \rightrightarrows Y$  be a correspondence from  $X$  to  $Y$ . Suppose that  $\Phi(\mathbf{x})$  is both non-empty and compact for all  $\mathbf{x} \in X$  and that the correspondence  $\Phi: X \rightarrow Y$  is both upper hemicontinuous and lower hemicontinuous. Let*

$$m(\mathbf{x}) = \sup\{f(\mathbf{x}, \mathbf{y}) : \mathbf{y} \in \Phi(\mathbf{x})\}$$

for all  $\mathbf{x} \in X$ , and let

$$M(\mathbf{x}) = \{\mathbf{y} \in \Phi(\mathbf{x}) : f(\mathbf{x}, \mathbf{y}) = m(\mathbf{x})\}$$

for all  $\mathbf{x} \in X$ . Then  $m: X \rightarrow \mathbb{R}$  is continuous,  $M(\mathbf{x})$  is a non-empty compact subset of  $Y$  for all  $\mathbf{x} \in X$ , and the correspondence  $M: X \rightrightarrows Y$  is upper hemicontinuous.

**Proof** Let  $\mathbf{x} \in X$ . Then  $\Phi(\mathbf{x})$  is a non-empty compact subset of  $Y$ . It is thus a closed bounded subset of  $\mathbb{R}^m$ . It follows from the Extreme Value Theorem (Theorem 1.17) that there exists at least one point  $\mathbf{y}^*$  of  $\Phi(\mathbf{x})$  with the property that  $f(\mathbf{x}, \mathbf{y}^*) \geq f(\mathbf{x}, \mathbf{y})$  for all  $\mathbf{y} \in \Phi(\mathbf{x})$ . Then  $m(\mathbf{x}) = f(\mathbf{x}, \mathbf{y}^*)$  and  $\mathbf{y}^* \in M(\mathbf{x})$ . Moreover

$$M(\mathbf{x}) = \{\mathbf{y} \in \Phi(\mathbf{x}) : f(\mathbf{x}, \mathbf{y}) = m(\mathbf{x})\}.$$

It follows from the continuity of  $f$  that the set  $M(\mathbf{x})$  is closed in  $Y$  (see Corollary 1.15). It is thus a closed subset of the compact set  $\Phi(\mathbf{x})$  and must therefore itself be compact.

Let some positive number  $\varepsilon$  be given. Then  $f(\mathbf{p}, \mathbf{y}) < m(\mathbf{p}) + \varepsilon$  for all  $\mathbf{y} \in \Phi(\mathbf{p})$ . It follows from Lemma 2.21 that

$$\{\mathbf{x} \in X : f(\mathbf{x}, \mathbf{y}) < m(\mathbf{p}) + \varepsilon \text{ for all } \mathbf{y} \in \Phi(\mathbf{x})\}$$

is open in  $X$ , and thus there exists some positive real number  $\delta_1$  such that  $f(\mathbf{x}, \mathbf{y}) < m(\mathbf{p}) + \varepsilon$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta_1$  and  $\mathbf{y} \in \Phi(\mathbf{x})$ . Then  $m(\mathbf{x}) < m(\mathbf{p}) + \varepsilon$  for all  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta_1$ .

The correspondence  $\Phi: X \rightrightarrows Y$  is also lower hemicontinuous. It therefore follows from Lemma 2.22 that there exists some positive real number  $\delta_2$  such that, given any  $\mathbf{x} \in X$  satisfying  $|\mathbf{x} - \mathbf{p}| < \delta_2$ , there exists some  $\mathbf{y} \in \Phi(\mathbf{x})$  for which  $f(\mathbf{x}, \mathbf{y}) > m(\mathbf{p}) - \varepsilon$ . It follows that  $m(\mathbf{x}) > m(\mathbf{p}) - \varepsilon$  whenever  $\mathbf{x} \in X$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta_2$ . Let  $\delta$  be the minimum of  $\delta_1$  and  $\delta_2$ . Then  $\delta > 0$ , and

$$m(\mathbf{p}) - \varepsilon < m(\mathbf{x}) < m(\mathbf{p}) + \varepsilon$$

whenever  $\mathbf{x} \in X$  satisfies  $|\mathbf{x} - \mathbf{p}| < \delta$ . Thus the function  $m: X \rightarrow \mathbb{R}$  is continuous on  $X$ .

It only remains to prove that the correspondence  $M: X \rightrightarrows Y$  is upper hemicontinuous. Let

$$\Psi(\mathbf{x}) = \{\mathbf{y} \in Y : f(\mathbf{x}, \mathbf{y}) = m(\mathbf{x})\}$$

for all  $\mathbf{x} \in X$ . Then

$$\text{Graph}(\Psi) = \{(\mathbf{x}, \mathbf{y}) \in X \times Y : f(\mathbf{x}, \mathbf{y}) = m(\mathbf{x})\}$$

Thus  $\text{Graph}(\Psi)$  is the preimage of zero under the continuous real-valued function that sends  $(\mathbf{x}, \mathbf{y}) \in X \times Y$  to  $f(\mathbf{x}, \mathbf{y}) - m(\mathbf{x})$ . It follows that  $\text{Graph}(\Psi)$  is a closed subset of  $X \times Y$ .

Now  $M(\mathbf{x}) = \Phi(\mathbf{x}) \cap \Psi(\mathbf{x})$  for all  $\mathbf{x} \in X$ , where the correspondence  $\Phi$  is compact-valued and upper hemicontinuous and the correspondence  $\Psi$  has closed graph. It follows from Proposition 2.20 that the correspondence  $M$  must itself be both compact-valued and upper hemicontinuous. This completes the proof of Berge's Maximum Theorem.  $\blacksquare$