

2009/10 Schol paper 1, my questions, outline solutions.

1. The Laplace transform of a function $f(t)$, $t \geq 0$ is a function of s defined as

$$\mathcal{L}[f(t)] = \int_0^{\infty} f(t)e^{-st} dt$$

Show

$$\mathcal{L}[e^{at}] = \frac{1}{s-a}$$

where a is a constant and you may assume $s > a$. Show that,

$$\mathcal{L}\left[\frac{df}{dt}\right] = s\mathcal{L}[f(t)] - f(0)$$

where you can assume

$$\lim_{t \rightarrow \infty} f(t)e^{-st} = 0$$

By taking the Laplace transform of both sides of the equation solve

$$\frac{df}{dt} = 2f$$

where $f(0) = -1$.

Solution: So you just have to follow this question along; first

$$\mathcal{L}[e^{at}] = \int_0^{\infty} e^{(a-s)t} dt = \frac{1}{a-s} e^{(a-s)t} \Big|_0^{\infty} = \frac{1}{s-a} \quad (1)$$

where we have assumed that $s > a$ in order to evaluate the exponential at infinity.

Next, from the definition

$$\mathcal{L}\left[\frac{df}{dt}\right] = \int_0^{\infty} \frac{df}{dt} e^{-st} dt \quad (2)$$

Now use integration by parts

$$\mathcal{L}\left[\frac{df}{dt}\right] = \int_0^{\infty} \frac{df}{dt} e^{-st} dt$$

$$= f(t)e^{-st} \Big|_0^\infty - s \int_0^\infty \frac{df}{dt} e^{-st} dt = s\mathcal{L}[f(t)] - f(0) \quad (3)$$

Finally taking the Laplace transform of both sides of the differential equation gives

$$s\mathcal{L}[f(t)] + 1 = 2\mathcal{L}[f(t)] \quad (4)$$

so, solving gives

$$\mathcal{L}[f(t)] + 1 = -\frac{1}{s-2} \quad (5)$$

and, hence, by the above, and noting that the transform is linear

$$f(t) = -e^{2t} \quad (6)$$

2. Write the on-off pulse

$$f(t) = \begin{cases} 1 & -\pi < t < 0 \\ -1 & 0 < t < \pi \\ 0 & \text{otherwise} \end{cases}$$

as a Fourier integral.

Solution: Well lets apply the definition and integrate

$$\begin{aligned} \widetilde{f(k)} &= \frac{1}{2\pi} \int_{-\infty}^{\infty} f(t)e^{-ikt} dt \\ &= \frac{1}{2\pi} \int_{-\pi}^0 e^{-ikt} dt - \frac{1}{2\pi} \int_0^{\pi} e^{-ikt} dt \\ &= \frac{1}{2\pi} \left[\frac{1}{-ik} e^{-ikt} \right]_{-\pi}^0 - \frac{1}{2\pi} \left[\frac{1}{-ik} e^{-ikt} \right]_0^{\pi} \\ &= \frac{i}{2\pi k} (1 - e^{ik\pi}) - \frac{i}{2\pi k} (e^{-ik\pi} - 1) \end{aligned} \quad (7)$$

Now, putting all that together

$$\widetilde{f(k)} = \frac{i}{\pi k} - \frac{i}{\pi k} \cos \pi k \quad (8)$$

3. Find the general solution of

$$\frac{d^3 y}{dt^3} - 3\frac{d^2 y}{dt^2} + 3\frac{dy}{dt} - y = e^t$$

*Solution:*First off, substitute into the homogenous equation

$$y = e^{\lambda t} \quad (9)$$

giving

$$\lambda^3 - 3\lambda^2 + 3\lambda - 1 = 0 \quad (10)$$

which factorizes as

$$(\lambda - 1)^3 = 0 \quad (11)$$

so the homogenous equation has general solution

$$y = (C_1 + tC_2 + t^2C_3)e^t \quad (12)$$

Hence, to match the right hand side we need to substitute

$$y = Ct^3e^t \quad (13)$$

so

$$\begin{aligned} y' &= Ct^3e^t + 3Ct^2e^t \\ y'' &= Ct^3e^t + 6Ct^2e^t + 6Cte^t \\ y''' &= Ct^3e^t + 9Ct^2e^t + 18Cte^t + 6Ce^t \end{aligned} \quad (14)$$

All the terms with t outside the exponential in them cancel and we get

$$6C = 1 \quad (15)$$

and hence the solution to the inhomogenous equation is

$$y = \left(C_1 + tC_2 + t^2C_3 + \frac{1}{6}t^3 \right) e^t \quad (16)$$

4. Find the general solution of

$$t^2\ddot{y} + 3t\dot{y} + y = t^2$$

*Solution:*So this is an Euler equation, let

$$t = e^z \quad (17)$$

so

$$\frac{dy}{dt} = \frac{dy}{dz} \frac{dz}{dt} = \frac{1}{t} \frac{dy}{dz} \quad (18)$$

and

$$\frac{d^2y}{dt^2} = \frac{d}{dt} \left(\frac{1}{t} \frac{dy}{dz} \right) = -\frac{1}{t^2} \frac{dy}{dz} + \frac{1}{t^2} \frac{d^2y}{dz^2} \quad (19)$$

Now, the equation becomes

$$\frac{d^2y}{dz^2} + 2\frac{dy}{dz} + y = e^{2z} \quad (20)$$

which can be solved using the normal substitution to give

$$y = (C_1 + C_2 z)e^{-z} + \frac{1}{9}e^{2z} \quad (21)$$

or

$$y = (C_1 + C_2 \log t) \frac{1}{t} + \frac{1}{9}t^2 \quad (22)$$