

2E2 Tutorial Sheet 20, Solutions¹

30 April 2004

Vectors calculus. With $f(x, y, z)$ a scalar field the grad of f , ∇f , is

$$\text{grad } f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} + \frac{\partial f}{\partial z} \mathbf{k} \quad (1)$$

1. (2) $f = x^2 + 2x^2yz$, find $\text{grad } f$. If $g = x^2yz + 2z$, what is $\nabla f \cdot \nabla g$.

Solution: So

$$\begin{aligned} \nabla f &= \frac{\partial}{\partial x} (x^2 + 2x^2yz) \mathbf{i} + \frac{\partial}{\partial y} (x^2 + 2x^2yz) \mathbf{j} + \frac{\partial}{\partial z} (x^2 + 2x^2yz) \mathbf{k} \\ &= (2x + 4xyz) \mathbf{i} + 2x^2z \mathbf{j} + 2x^2y \mathbf{k} \end{aligned} \quad (2)$$

and

$$\begin{aligned} \nabla g &= \frac{\partial}{\partial x} (x^2yz + 2z) \mathbf{i} + \frac{\partial}{\partial y} (x^2yz + 2z) \mathbf{j} + \frac{\partial}{\partial z} (x^2yz + 2z) \mathbf{k} \\ &= 2xyzi + x^2z \mathbf{j} + (x^2y + 2) \mathbf{k} \end{aligned} \quad (3)$$

and

$$\begin{aligned} \nabla f \cdot \nabla g &= 2xyz(2x + 4xyz) + 2x^2zx^2z + 2x^2y(2 + x^2y) \\ &= 4x^2yz + 8x^2y^2z^2 + 2x^4z^2 + 4x^2y + 2x^4y^2 \end{aligned} \quad (4)$$

2. (2) Find the directional derivative of $z/(x^2 + y^2)$ in the direction \mathbf{i} .

Solution: Well the directional derivative along \mathbf{i} is $\text{grad } f \cdot \mathbf{i}$, in other words the one component of the grad. Furthermore

$$\frac{\partial}{\partial x} \frac{z}{x^2 + y^2} = -\frac{2xz}{(x^2 + y^2)^2} \quad (5)$$

3. (2) $f = xyz$, work out $\text{grad } f$. What is the value of $\text{grad } f$ at $(2, 1, 2)$. What is the directional derivative of f in the \mathbf{i} -direction at $(2, 1, 2)$.

Solution: So,

$$\begin{aligned} \text{grad } xyz &= \frac{\partial xyz}{\partial x} \mathbf{i} + \frac{\partial xyz}{\partial y} \mathbf{j} + \frac{\partial xyz}{\partial z} \mathbf{k} \\ &= yz \mathbf{i} + xz \mathbf{j} + xy \mathbf{k} \end{aligned} \quad (6)$$

¹Conor Houghton, houghton@maths.tcd.ie and <http://www.maths.tcd.ie/~houghton/2E2.html>

At $(2, 1, 2)$, $x = 2$, $y = 1$ and $z = 2$, so

$$\text{grad } f = 2\mathbf{i} + 4\mathbf{j} + 2\mathbf{k} \quad (7)$$

The direction derivative in the \mathbf{i} direction is given by $\mathbf{i} \cdot \text{grad } f = 2$.

4. (2) $f = x^2 + y^2 + z^2$, find $\text{grad } f$. What is the directional derivative of f in the direction of $\mathbf{b} = (1, 1, 1)$ at the point $(1, 1, 1)$. Remember to use a unit vector when working out the directional derivative. What is the directional derivative in the \mathbf{j} direction at $(1, 0, 0)$; what about the \mathbf{j} direction?

Solution: So,

$$\nabla f = 2x\mathbf{i} + 2y\mathbf{j} + 2z\mathbf{k} \quad (8)$$

The directional derivative of f in the direction of $\mathbf{b} = (1, 1, 1)$ is given by $\hat{\mathbf{b}} \cdot \nabla f$ where

$$\hat{\mathbf{b}} = \frac{\mathbf{b}}{|\mathbf{b}|} \quad (9)$$

is the unit vector in the \mathbf{b} direction. Here $|\mathbf{b}| = \sqrt{3}$ so

$$\hat{\mathbf{b}} = \left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right) \quad (10)$$

At $(1, 1, 1)$, we have $x = y = z = 1$ so $\nabla f = 2\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ and hence

$$D_{\hat{\mathbf{b}}} f = \frac{2}{\sqrt{3}} + \frac{2}{\sqrt{3}} + \frac{2}{\sqrt{3}} = 2\sqrt{3} \quad (11)$$

At $(1, 0, 0)$ $\text{grad } f = 2\mathbf{i}$ so the direction derivative in the \mathbf{i} direction is two and in the \mathbf{j} direction is zero.