

2E2 Tutorial Sheet 2 First Term, Solutions¹

17 October 2003

1. Using the Laplace transform solve the differential equation

$$f'' - 4f' + 3f = 1 \quad (1)$$

with boundary conditions $f(0) = f'(0) = 0$.

Solution: First, take the Laplace transform of the equation. Since $f'(0) = f(0) = 0$, if $\mathcal{L}(f) = F(s)$ then $\mathcal{L}(f') = sF(s)$ and $\mathcal{L}(f'') = s^2F(s)$. Thus, the subsidiary equation is

$$s^2F - 4sF + 3F = \frac{1}{s} \quad (2)$$

and so

$$\begin{aligned} (s^2 - 4s + 3)F &= \frac{1}{s} \\ F &= \frac{1}{s} \frac{1}{s^2 - 4s + 3} \end{aligned} \quad (3)$$

and, since $s^2 - 4s + 3 = (s - 3)(s - 1)$, this gives

$$F = \frac{1}{s(s - 3)(s - 1)} \quad (4)$$

Before we can invert this, we need to do a partial fraction expansion.

$$\begin{aligned} \frac{1}{s(s - 3)(s - 1)} &= \frac{A}{s} + \frac{B}{s - 3} + \frac{C}{s - 1} \\ 1 &= A(s - 3)(s - 1) + Bs(s - 1) + Cs(s - 3) \end{aligned} \quad (5)$$

So substituting in $s = 0$ we get $A = 1/3$, $s = 3$ gives $B = 1/6$ and $s = 1$ gives $C = -1/2$. Hence

$$F = \frac{1}{3s} + \frac{1}{6(s - 3)} - \frac{1}{2(s - 1)} \quad (6)$$

and so

$$f(t) = \frac{1}{3} + \frac{1}{6}e^{3t} - \frac{1}{2}e^t \quad (7)$$

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/2E2.html>

2. Using the Laplace transform solve the differential equation

$$f'' - 4f' + 3f = 2e^t \quad (8)$$

with boundary conditions $f(0) = f'(0) = 0$.

Solution: This time we have $\mathcal{L}(2e^t) = 2/(s - 1)$ on the right hand side. This means that the subsidiary equation is

$$(s^2 - 4s + 3)F = \frac{2}{s - 1} \quad (9)$$

so

$$F = \frac{2}{(s - 1)^2(s - 3)} \quad (10)$$

We need to do partial fractions again, but this is one of those cases with a repeated root:

$$\frac{1}{(s - 1)^2(s - 3)} = \frac{A}{s - 1} + \frac{B}{(s - 1)^2} + \frac{C}{s - 3} \quad (11)$$

and multiplying across

$$1 = A(s - 1)(s - 3) + B(s - 3) + C(s - 1)^2 \quad (12)$$

so $s = 1$ gives $B = -1/2$ and $s = 3$ gives $C = 1/4$. No value of s gives A on its own so we try $s = 2$:

$$1 = -A + \frac{1}{2} + \frac{1}{4} \quad (13)$$

which means that $A = -1/4$. Hence

$$F = -\frac{1}{2(s - 1)} - \frac{1}{(s - 1)^2} + \frac{1}{2(s - 3)} \quad (14)$$

and

$$f = -\frac{1}{2}e^t - te^t + \frac{1}{2}e^{3t} \quad (15)$$

3. Using the Laplace transform solve the differential equation

$$f'' - 4f' + 3f = 0 \quad (16)$$

with boundary conditions $f(0) = 1$ and $f'(0) = 1$.

Solution: In this example there are non-zero boundary conditions. Since

$$\mathcal{L}(f') = sF - f(0) \quad (17)$$

$$\mathcal{L}(f'') = s^2F - sf(0) - f'(0) \quad (18)$$

the subsidiary equation in this case is

$$s^2F - s - 1 - 4sF + 4 + 3F = 0 \quad (19)$$

so

$$(s^2 - 4s + 3)F = s - 3. \quad (20)$$

Hence

$$F = \frac{1}{s-1} \quad (21)$$

and

$$f(t) = e^t \quad (22)$$

4. Using the Laplace transform solve the differential equation

$$y'' - 2ay' + a^2y = 0 \quad (23)$$

with boundary conditions $y'(0) = 1$ and $y(0) = 0$. a is some real constant.

Solution: Taking the Laplace transform we get

$$s^2Y - 1 - 2aY + a^2Y = 0 \quad (24)$$

and hence

$$Y = \frac{1}{(s-a)^2} \quad (25)$$

which means that

$$y = te^{at} \quad (26)$$