2E2 Tutorial Sheet 8 Second Term¹

28 November 2003

1. (3) Find the eigenvectors and eigenvalues of the following matrices

$$(i) \quad \begin{pmatrix} 4 & 0 \\ 0 & -6 \end{pmatrix} \qquad (ii) \quad \begin{pmatrix} 10 & -4 \\ 18 & -12 \end{pmatrix} \qquad (iii) \quad \begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} \qquad (1)$$

Solution: In (i) the characteristic equation is

$$\begin{vmatrix} 4-\lambda & 0\\ 0 & -6-\lambda \end{vmatrix} = 0$$
 (2)

 \mathbf{SO}

$$(4-\lambda)(-6-\lambda) = 0 \tag{3}$$

So $\lambda = 4$ or lambda = -6. Taking the $\lambda = 4$ first

$$\begin{pmatrix} 4 & 0 \\ 0 & -6 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 4 \begin{pmatrix} a \\ b \end{pmatrix}$$
(4)

so 4a = 4a and -6b = 4b, hence b = 0 and a is arbitrary, taking a = 1 an eigenvalue 4 eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1\\0 \end{pmatrix} \tag{5}$$

Taking the $\lambda = -6$

$$\begin{pmatrix} 4 & 0 \\ 0 & -6 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -6 \begin{pmatrix} a \\ b \end{pmatrix}$$
(6)

so 4a = -6a and -6b = -6b, hence a = 0 and b is arbitrary, taking b = 1 a eigenvalue -6 eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 0\\1 \end{pmatrix} \tag{7}$$

(9)

In (ii) the characteristic equation is

$$\begin{vmatrix} 10 - \lambda & -4 \\ 18 & -12 - \lambda \end{vmatrix} = 0 \tag{8}$$

 \mathbf{SO}

$$(10 - \lambda)(-12 - \lambda) - (-4)18 = 0$$

¹Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/2E2.html

or

 $\lambda^2 + 2\lambda - 48 = 0$

(1(

Solve this gives us $\lambda = 6$ or $\lambda = -8$. Taking the $\lambda = 6$ first

$$\begin{pmatrix} 10 & -4\\ 18 & -12 \end{pmatrix} \begin{pmatrix} a\\ b \end{pmatrix} = 6 \begin{pmatrix} a\\ b \end{pmatrix}$$
(11)

so the first equation is 10a - 4b = 6a or a = b, the other equation is 18a - 12b = 6 which is also a = b. Taking a = 1 an eigenvalue 6 eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1\\1 \end{pmatrix} \tag{12}$$

Taking the $\lambda = -8$ next

$$\begin{pmatrix} 10 & -4\\ 18 & -12 \end{pmatrix} \begin{pmatrix} a\\ b \end{pmatrix} = -8 \begin{pmatrix} a\\ b \end{pmatrix}$$
(13)

so the first equation is 10a - 4b = -8a or 9a = 2b, the other equation is 18a - 12b = -8b which is also 9a = 2b. Taking a = 2, b = 9 an eigenvalue -8 eigenvector is,

 $\mathbf{x} = \begin{pmatrix} 2\\ 9 \end{pmatrix} \tag{14}$

In (iii) the characteristic equation is

$$\begin{vmatrix} -\lambda & r \\ r & -\lambda \end{vmatrix} = 0 \tag{15}$$

 \mathbf{SO}

$$\lambda^2 - r^2 = 0 \tag{1}$$

or

$$\lambda = \pm r \tag{1}$$

Taking the $\lambda = r$ first

$$\begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = r \begin{pmatrix} a \\ b \end{pmatrix}$$
(18)

so the equation is rb = ra or a = b. Taking a = 1 an eigenvalue r eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1\\1 \end{pmatrix} \tag{19}$$

Taking the $\lambda = -r$

$$\begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -r \begin{pmatrix} a \\ b \end{pmatrix}$$
(20)

so the equation is rb = -ra or a = -b. Taking a = 1 an eigenvalue -r eigenvector is,

$$\mathbf{x} = \begin{pmatrix} 1\\ -1 \end{pmatrix} \tag{2}$$

2. (2) Rewrite

$$\begin{array}{rcl}
10y_1 - 4y_2 &=& 2\\
18y_1 - 12y_2 &=& 3
\end{array} \tag{22}$$

in the matrix form $A\mathbf{y} = \mathbf{a}$ where

$$\mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \tag{23}$$

Solution: This is equivalent to the matrix equation

$$\begin{pmatrix} 10 & -4\\ 18 & -12 \end{pmatrix} \begin{pmatrix} y_1\\ y_2 \end{pmatrix} = \begin{pmatrix} 2\\ 3 \end{pmatrix}$$
(24)

You can check this by multiplying it out.



Figure 1: Two containers with flow between them.

3. (3) As illustrated in Fig. 1, two large containers are connected and American style sandwich spead is pumped between them at a rate of $1/2m^3s^{-1}$. One container has volume $5m^3$, the other $7m^3$. Both are full of spread. Initially the smaller container contains pure jam, the second container has $5m^3$ of jam and $2m^3$ of peanut butter. Assume perfect mixing and so on.

(i) Write down the differential equation for $y_1(t)$ and $y_2(t)$, the amount of peanut butter in the first and second container.

(ii) Solve it to find $y_1(t)$ and $y_2(t)$ explicitly.

(iii) The amount of peanut butter in each container will change less and less as time passes, what values are the tending towards.

Solution: Well if there is y_1 peanut butter in the small container then the concentration of the spead in the small container is $y_1/5$ and so $y_1/10$ is flowing out per

second. In the same way $y_2/7$ is the concentration of peanut butter in the secont tank and so $y_2/14$ per second is going from the large tank to the small one. The means the equations are

$$y_1' = -\frac{1}{10}y_1 + \frac{1}{14}y_2 \tag{2}$$

$$y_2' = \frac{1}{10}y_1 - \frac{1}{14}y_2 \tag{2}$$

This equation can be rewritten

$$\mathbf{y}' = \begin{pmatrix} -\frac{1}{10} & \frac{1}{14} \\ \frac{1}{10} & -\frac{1}{14} \end{pmatrix} \mathbf{y}$$
(2)

We work out the eigenvalues

$$\begin{vmatrix} -\frac{1}{10} - \lambda & \frac{1}{14} \\ \frac{1}{10} & -\frac{1}{14} - \lambda \end{vmatrix} = \left(\frac{1}{10} + \lambda\right) \left(\frac{1}{14} + \lambda\right) - \frac{1}{140}$$
(28)

$$= \lambda^2 + \frac{6}{35}\lambda = 0 \tag{29}$$

This means that there are two eigenvalues, $\lambda_1 = 0$ and $\lambda_2 = -6/35$. The corresponding eigenvectors are given by

$$\begin{pmatrix} -\frac{1}{10} & \frac{1}{14} \\ \frac{1}{10} & -\frac{1}{14} \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = 0$$
(30)

which has solutions of the form a = 10 and b = 14 for λ_1 and

$$\begin{pmatrix} -\frac{1}{10} & \frac{1}{14} \\ \frac{1}{10} & -\frac{1}{14} \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -\frac{6}{35} \begin{pmatrix} a \\ b \end{pmatrix}$$
(31)

for λ_2 . This has solution a = -1 and b = 1. Thus, the general solution is

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = c_1 \begin{pmatrix} 10 \\ 14 \end{pmatrix} + c_2 \begin{pmatrix} -1 \\ 1 \end{pmatrix} e^{-\frac{6}{35}t}$$
(32)

For part (iii), matching with $y_1(0) = 0$ and $y_2(0) = 2$, this gives $c_1 = 1/12$ and $c_2 = 5/6$ and hence

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \frac{1}{12} \begin{pmatrix} 10 \\ 14 \end{pmatrix} + \frac{5}{6} \begin{pmatrix} -1 \\ 1 \end{pmatrix} e^{-\frac{6}{35}t}$$
(33)

By the way, not the end result, clearly the exponetially decaying part goes away wit time so that (5, 5)

4

$$\lim_{t \to \infty} \mathbf{y} = \begin{pmatrix} \frac{5}{6} \\ \frac{7}{6} \end{pmatrix} \tag{34}$$