

# READING WEEK EXERCISES FOR COURSE MA3431

## Recommended textbooks for the Course MA3431 in Classical Field Theory

- Classical Electrodynamics, J. D. Jackson, John Wiley, 1998 (3rd ed) [537.12 K23]
- The Classical Theory of Fields, E. M. Lifshitz and L. D. Landau [530.14 L52]
- Classical Field Theory, Francis E. Low, John Wiley & Sons, 1997 [530.14 N71]
- Classical Mechanics, Herbert Goldstein, Addison-Wesley, 1980 [531 J0\*1]

1. A Lagrangian density  $\mathcal{L}(\phi, \partial_\mu\phi)$  depends upon a scalar field  $\phi(x)$  and its partial derivative  $\partial_\mu\phi = \partial\phi/\partial x^\mu$ . The space-time point  $x$  refers to  $x^\mu$  with components  $(x^0, x^1, x^2, x^3)$ . Consider the following Lagrangian density where  $\sigma$  is a constant

$$\mathcal{L} = \frac{1}{2} \partial_\lambda\phi \partial^\lambda\phi + \frac{1}{3} \sigma \phi^3.$$

- (a) Write down the Euler-Lagrange equation of motion of the scalar field  $\phi(x)$ .
- (b) Evaluate the equation of motion for the particular Lagrangian density  $\mathcal{L}$  provided above.
- (c) Calculate from the above Lagrangian density  $\mathcal{L}$  the following stress tensor

$$T^{\mu\nu} = \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \partial^\nu\phi - g^{\mu\nu}\mathcal{L}$$

where the metric tensor  $g^{\mu\nu}$  is diagonal with matrix entries  $(+1, -1, -1, -1)$ .

- (d) Find the 4-divergence of the stress tensor  $T^{\mu\nu}$  you obtained, namely, determine

$$\partial_\mu T^{\mu\nu}.$$

- (e) Show that the stress tensor  $T^{\mu\nu}$  is conserved by demonstrating that its 4-divergence is zero when the scalar field  $\phi(x)$  obeys its equation of motion

$$\partial_\mu T^{\mu\nu} = 0.$$

2. (a) Show that the electric and magnetic induction fields transform according to

$$\begin{aligned} E'_1 &= E_1 & B'_1 &= B_1 \\ E'_2 &= \gamma E_2 - \gamma\beta B_3 & B'_2 &= \gamma B_2 + \gamma\beta E_3 \\ E'_3 &= \gamma E_3 + \gamma\beta B_2 & B'_3 &= \gamma B_3 - \gamma\beta E_2 \end{aligned}$$

namely, Eq. (11.148) of J. D. Jackson's *Classical Electrodynamics*, using

$$F'^{\mu\nu} = \Lambda^\mu{}_\rho \Lambda^\nu{}_\sigma F^{\rho\sigma}$$

when the velocity  $\beta c$  of frame  $K'$  is directed along the  $x^1$  axis of frame  $K$  with

$$\Lambda^\mu{}_\rho = \begin{pmatrix} \gamma & -\beta\gamma & 0 & 0 \\ -\beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Assume that the elements of the matrix  $F_{\mu\nu}$  may be written in terms of the fields  $\vec{E}$  and  $\vec{B}$  in a form given by Eq. (11.137) of J. D. Jackson's *CED*, viz.

$$F^{\mu\nu} = \begin{pmatrix} 0 & -E^1 & -E^2 & -E^3 \\ E^1 & 0 & -B^3 & B^2 \\ E^2 & B^3 & 0 & -B^1 \\ E^3 & -B^2 & B^1 & 0 \end{pmatrix}.$$

- (b) Show that the electric and magnetic fields in frame  $K$  of the electrostatic field of a stationary charge  $q$  in a frame  $K'$  moving with velocity  $v = \beta c$  along the  $x^1$  axis of frame  $K$  are given by Eq. (11.152) of J. D. Jackson's *CED*,  $3e$

$$\begin{aligned} E^1 &= -\frac{\gamma qvt}{(b^2 + \gamma^2 v^2 t^2)^{3/2}} & B^1 &= 0 \\ E^2 &= \frac{\gamma qb}{(b^2 + \gamma^2 v^2 t^2)^{3/2}} & B^2 &= 0 \\ E^3 &= 0 & B^3 &= \frac{\beta\gamma qb}{(b^2 + \gamma^2 v^2 t^2)^{3/2}} \end{aligned}$$

$t$  referring to the time since the origins of the frames  $K$  and  $K'$  overlapped and  $b$  being the closest distance of approach of the charge, assumed fixed on the  $x'^2$  axis. J. D. Jackson uses ordinary Cartesian spatial components in Eq. (11.148) and in Eq. (11.152). Contravariant indices are used here.