

**2E1 (Timoney) Tutorial sheet 12**  
[Tutorials January 24 – 25, 2007]

**Name:** Solutions

1. Evaluate


$$\int_0^3 \int_{y=0}^{y=\sqrt{3}x} \frac{1}{\sqrt{x^2 + y^2}} dy dx$$

by making a change of variables to polar coordinates. [Hint: Sketch the region first. Then do the  $dr$  integral first, before  $d\theta$ .]

*Solution:* This iterated integral is the same (by Fubini's theorem) as the double integral

$$\iint_R \frac{1}{\sqrt{x^2 + y^2}} dx dy$$

where  $R$  is the triangle in the plane bounded by the  $x$ -axis, the line  $x = 3$  and the line  $y = \sqrt{3}x$ .



The limit  $x = 3$  is  $r \cos \theta = 3$  or  $r = 3 / \cos \theta$  and the angle is  $\pi/3$  (because  $\tan(\pi/3) = \sqrt{3}$ ).

Changing this integral to polar coordinates, and remembering that  $dx dy = r dr d\theta$ , we get

$$\begin{aligned}
 \int_{\theta=0}^{\theta=\pi/3} \left( \int_{r=0}^{r=3/\cos\theta} \frac{1}{r} r dr \right) d\theta &= \int_{\theta=0}^{\theta=\pi/3} [r]_{r=0}^{r=3/\cos\theta} d\theta \\
 &= \int_{\theta=0}^{\theta=\pi/3} \frac{3}{\cos\theta} d\theta \\
 &= \int_{\theta=0}^{\theta=\pi/3} 3 \sec\theta d\theta \\
 &= [3 \ln |\sec\theta + \tan\theta|]_0^{\pi/3} \\
 &= 3 \ln(2 + \sqrt{3})
 \end{aligned}$$

2. Find the mass of a solid object occupying the region in space bounded by the coordinate planes and the plane  $x + y + z = 2$  if its density function is  $\delta(x, y, z) = x^2$ . [Hint: Calculations are easier if you leave the  $dx$  integral to last.]

*Solution:* We know that the answer is

$$\text{Mass} = \iiint dm = \iiint \delta(x, y, z) dx dy dz$$

with the triple integral extending over the object.

That gives

$$\begin{aligned}
 &\int_{x=0}^2 \left( \int_{y=0}^{2-x} \left( \int_{z=0}^{z=2-x-y} x^2 dz \right) dy \right) dx \\
 &= \int_{x=0}^2 \left( \int_{y=0}^{2-x} [x^2 z]_{z=0}^{z=2-x-y} dy \right) dx \\
 &= \int_{x=0}^2 \left( \int_{y=0}^{2-x} x^2(2-x-y) dy \right) dx \\
 &= \int_{x=0}^2 [x^2((2-x)y - y^2/2)]_{y=0}^{2-x} dx \\
 &= \int_{x=0}^2 x^2((2-x)^2 - (2-x)^2/2) - 0 dx \\
 &= \int_{x=0}^2 x^2(2-x)^2/2 dx = \frac{1}{2} \int_{x=0}^2 x^2(4-4x+x^2) dx \\
 &= \frac{1}{2} \int_{x=0}^2 4x^2 - 4x^3 + x^4 dx \\
 &= \frac{1}{2} [(4/3)x^3 - x^4 + x^5/5]_{x=0}^2 \\
 &= \frac{1}{2} \left( \frac{32}{3} - 16 + \frac{32}{5} - 0 \right) = \frac{8}{15}
 \end{aligned}$$