

Course 141: MECHANICS

Problem Set 3

Date Issued: November 7, 2007

Date Due: November 14, 2007

1. (5 points) Simplify: $\vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b})$.
2. (5 points) Two particles 1 and 2 are moving along x- and y-axis with respective velocities $\vec{v}_1 = 2m\hat{i}/s$ and $\vec{v}_2 = 3m\hat{j}/s$. At $t = 0$ they are at $x_1 = -3m$, $y_1 = 0$ and $x_2 = 0$, $y_2 = -3m$, respectively (here s and m are parameters).
 - (a) Sketch the positions of the particles at $t = 0$, the direction of their velocities, and the vector $\vec{r}_2 - \vec{r}_1$ which represents the position of the particle 2 with respect to particle 1.
 - (b) Find $\vec{r}_2 - \vec{r}_1$ as a function of time.
 - (c) When and where are these particles closest?
3. (5 points) A uniformly accelerated car passes two telephone poles with velocities 10 km/hr and 20 km/hr, respectively. Calculate its velocity when it is halfway between the poles. Let a = car's acceleration.
4. (10 points) A spring gun fires two shots, the velocity in each case being v_0 . In the first case the shot is fired at an angle of elevation α and in the second case at an angle α' . Show that the time interval between the two firing such that two shots will collide in mid-air is

$$\delta t = \frac{2v_0}{g} \frac{\sin(\alpha - \alpha')}{\cos \alpha + \cos \alpha'}$$

5. (5 points bonus problem) Three small snails are each at a vertex of an equilateral triangle of side 60 cm. The first sets out towards the second, the second towards the third, and the third towards the first, with a uniform speed of 5 cm/min. During their motion each of them always heads towards its respective target snail. How long does it take for them to meet?
6. (5 points bonus problem) In problem 5 how many revolutions do the snails make before they meet? Assume point-like snails.