Beiträge zur Algebra und Geometrie
Contributions to Algebra and Geometry
Volume 46 (2005), No. 2, 537-544.

A Note on the Existence of $\{k, k\}$-equivelar Polyhedral Maps

Basudeb Datta

Department of Mathematics, Indian Institute of Science,
Bangalore 560 012, India
e-mail: dattab@math.iisc.ernet.in

Abstract. A polyhedral map is called $\{p, q\}$-equivelar if each face has p edges and each vertex belongs to q faces. In [12], it was shown that there exist infinitely many geometrically realizable $\{p, q\}$-equivelar polyhedral maps if $q > p = 4$, $p > q = 4$ or $q - 3 > p = 3$. It was shown in [6] that there exist infinitely many $\{4, 4\}$- and $\{3, 6\}$-equivelar polyhedral maps. In [1], it was shown that $\{5, 5\}$- and $\{6, 6\}$-equivelar polyhedral maps exist. In this note, examples are constructed, to show that infinitely many self dual $\{k, k\}$-equivelar polyhedral maps exist for each $k \geq 5$. Also vertex-minimal non-singular $\{p, p\}$-patterns are constructed for all odd primes p.

MSC 2000: 52B70, 51M20, 57M20
Keywords: Polyhedral maps, equivelar maps, non-singular patterns

1. Introduction and results

A *polyhedral complex* (of dimension 2) is collection of cycles (finite connected 2-regular graphs) together with the edges and the vertices in the cycles such that the intersection of any two cycles is empty, a vertex or an edge. The cycles are called the *faces* of the polyhedral complex. For a polyhedral complex K, $V(K)$ denotes its vertex-set and $\mathrm{EG}(K)$ denotes its edge-graph or 1-skeleton. We say K *finite* if $V(K)$ is finite. If $\mathrm{EG}(K)$ is connected then K is said to be *connected*.

A polyhedral complex is called a *polyhedral 2-manifold* (or an *abstract polyhedron*) if for each vertex v the faces containing v are of the form $F_1, \ldots, F_m$, where $F_1 \cap F_2, \ldots, F_{m-1} \cap$

0138-4821/93 $ 2.50 © 2005 Heldermann Verlag

$F_m, F_m \cap F_1$ are edges for some $m \geq 3$. A connected finite polyhedral 2-manifold is called a *polyhedral map*. A *combinatorial 2-manifold* is a polyhedral 2-manifold whose faces are 3-cycles. A polyhedral map is called $\{p, q\}$-*equivelar* if each face is a p-cycle and each vertex is in q faces. A polyhedral map is called *equivelar* if it is $\{p, q\}$-equivelar for some p, q (cf. [10, 3, 4, 11]).

To each polyhedral complex K, we associate a pure 2-dimensional simplicial complex $B(K)$ (called the *barycentric subdivision* of K) whose 2-faces are of the form ueF, where (u, e, F) is a flag (i.e., e is an edge of the face F and u is a vertex of e) in K. The geometric carrier of $B(K)$ is called the *geometric carrier* of K and is denoted by $|K|$. Clearly, K is a polyhedral 2-manifold if and only if $B(K)$ is a combinatorial 2-manifold (equivalently, $|K|$ is a 2-manifold). A polyhedral 2-manifold K is called *orientable* if $|K|$ is orientable.

An *isomorphism* between two polyhedral complexes K and L is a bijection $\varphi\colon V(K) \to V(L)$ such that $(v_1, \ldots, v_m)$ is a face of K if and only if $(\varphi(v_1), \ldots, \varphi(v_m))$ is a face of L. Two complexes are called *isomorphic* if there is an isomorphism between them. We identify two isomorphic polyhedral complexes. An isomorphism from K to itself is called an *automorphism* of K. The set $\Gamma(K)$ of automorphisms of K forms a group. A polyhedral 2-manifold K is called *combinatorially regular* if $\Gamma(K)$ is transitive on flags (cf. [10]).

For a polyhedral 2-manifold K, consider the polyhedral complex $\widetilde{K}$ whose vertices are the faces of K and $(F_1, \ldots, F_m)$ is a face of $\widetilde{K}$ if $F_1, \ldots, F_m$ have a common vertex and $F_1 \cap F_2, \ldots, F_{m-1} \cap F_m, F_m \cap F_1$ are edges. Then $\widetilde{K}$ is a polyhedral 2-manifold and called the *dual* of K. If $\widetilde{K}$ is isomorphic to K then K is called *self dual*.

A *pattern* is an ordered pair (M, G), where M is a connected closed surface in some Euclidean space and G is a finite graph on M such that each component of $M \setminus G$ is simply connected. The closure of each component of $M \setminus G$ is called a *face* of (M, G). For a face F, the closed path (in G) consisting of all the edges and the vertices in F is called the *boundary* of F. A pattern (M, G) is said to be *non-singular* if the boundary of each face is a cycle. A non-singular pattern is said to be a *polyhedral pattern* if the intersection of any two faces is empty, a vertex or an edge. A pattern (M, G) is called a $\{p, q\}$-*pattern* if each face contains p edges and the degree of each vertex in G is q (cf. [7]).

If (M, G) is a polyhedral pattern then clearly the boundaries of the faces of (M, G) form a polyhedral map. Conversely, for a polyhedral map K, let $M = |K|$ and $G = \mathrm{EG}(K)$. Then (M, G) is a polyhedral pattern and the faces of K are the boundaries of the faces of (M, G). This pattern (M, G) is called a *geometric realization* of K. A geometric realization (M, G) (in some $\mathbb{R}^n$) is called *linear* if each face of M is a convex polygon and no two adjacent faces (i.e., faces which share a common edge) lie in the same plane. If a polyhedral map has a linear geometric realization in $\mathbb{R}^3$ then it is called *geometrically realizable*.

If $f_0(K)$, $f_1(K)$ and $f_2(K)$ are the number of vertices, edges and faces respectively of a polyhedral complex K then the number $\chi(K) := f_0(K) - f_1(K) + f_2(K)$ is called the *Euler characteristic* of K. Observe that $\chi(B(K)) = \chi(K)$. If u and v are vertices of a face F and uv is not an edge of F then uv is called a *diagonal*. Clearly, if $d(K)$ is the number of diagonals of a polyhedral complex K then $d(K) + f_1(K) \leq \binom{f_0(K)}{2}$ and in the case of equality each pair of vertices belongs to a face. A polyhedral map K is called a *weakly neighbourly polyhedral map* (in short, *wnp map*) if each pair of vertices belongs to a common face.

We know (cf. [6]) that there exists a unique $\{p, q\}$-equivelar polyhedral map if $(p, q) = (3, 3)$,

$(3,4)$ or $(4,3)$ and there are exactly two $\{p,q\}$-equivelar polyhedral maps if $(p,q) = (3,5)$ or $(5,3)$. In [12], McMullen et al. constructed infinitely many geometrically realizable $\{p,q\}$-equivelar polyhedral maps for each $(p,q) \in \{(r,4) : r \geq 5\} \cup \{(4,s) : s \geq 5\} \cup \{(3,k) : k \geq 7\}$. In [6], it was shown that there exist infinitely many $\{4,4\}$- and $\{3,6\}$-equivelar polyhedral maps. It was also shown that there are exactly two neighbourly $\{3,8\}$-equivelar polyhedral maps and there are exactly 14 neighbourly $\{3,9\}$-equivelar polyhedral maps.

In [5], Coxeter constructed a geometrically realizable combinatorially regular infinite polyhedral 2-manifold whose faces are hexagons and each vertex is in six faces (namely, $\{6,6\,|\,3\}$). In [9], Grünbaum constructed another combinatorially regular infinite polyhedral 2-manifold of type $\{6,6\}$ (namely, $\{6,6\}_4$) (cf. [10]). In [8], Gott constructed a geometrically realizable infinite polyhedral 2-manifold whose faces are pentagons and each vertex is in five faces. If K is a $\{p,q\}$-equivelar polyhedral map on n vertices then $d(K) = nq(p-3)/2$ and $f_1(K) = nq/2$. Therefore, if K is an n-vertex $\{p,p\}$-equivelar polyhedral map then $np(p-3)/2 + np/2 \leq n(n-1)/2$ and hence $n \geq (p-1)^2$. Clearly, equality holds if and only if K is a wnp map. Let $\alpha(p)$ denote the smallest n such that there exists an n-vertex $\{p,p\}$-equivelar polyhedral map. Clearly, the 4-vertex 2-sphere (the boundary of a 3-simplex) is the unique $\{3,3\}$-equivelar wnp map. In [1], Brehm proved that there exist exactly three $\{4,4\}$-equivelar wnp maps and constructed the 16-vertex $\{5,5\}$-equivelar polyhedral map $M_{5,16}$ (of Example 1). It was shown in [2] that $M_{5,16}$ is the unique $\{5,5\}$-equivelar polyhedral map on 16 vertices. So, $\alpha(k) = (k-1)^2$ for $k \leq 5$. In [1], Brehm also constructed the 26-vertex $\{6,6\}$-equivelar polyhedral map $M_{6,26}$ (of Example 1). Here we show:

Theorem 1. *For each $m \geq 3$ and $n \geq 0$, there exist a $2(3^{m-1}+2n-1)$-vertex self dual $\{2m-1,2m-1\}$-equivelar polyhedral map and a (3^m+2n-1)-vertex self dual $\{2m,2m\}$-equivelar polyhedral map.*

Thus $(2m-2)^2 \leq \alpha(2m-1) \leq 2(3^{m-1}-1)$ and $(2m-1)^2 \leq \alpha(2m) \leq 3^m-1$ for all $m \geq 3$. In [13], using a computer, Nilakantan has shown that there does not exist any 25-vertex $\{6,6\}$-equivelar polyhedral map. So, $\alpha(6) = 26$ and hence there does not exist any $\{6,6\}$-equivelar wnp map. We believe the following is true:

Conjecture 1. *There does not exist any $\{k,k\}$-equivelar wnp map for $k \geq 7$.*

For the existence of an n-vertex $\{k,k\}$-pattern n must be at least $k+1$. Here we show:

Theorem 2. *There exists a $(p+1)$-vertex non-singular $\{p,p\}$-pattern for each odd prime p.*

2. Examples and proofs of the results

We first construct infinitely many $\{k,k\}$-equivelar polyhedral maps. We need these to prove our results. We identify a polyhedral complex with the set of faces in it.

Example 1. For $m \geq 3$ and $n \geq 0$, let

$$\begin{aligned} M_{2m-1,2(3^{m-1}+2n-1)} &= \{F_{i,2m-1} : 1 \leq i \leq 2(3^{m-1}+2n-1)\}, \\ M_{2m,3^m+2n-1} &= \{F_{i,2m} : 1 \leq i \leq 3^m+2n-1\}, \end{aligned}$$

where $b_{2l-1} = 3^{l-1} - 1$, $b_{2l} = 2 \times 3^{l-1} - 1$, for $l \geq 1$ and

$$\begin{aligned} F_{i,2m-1} &= (i + b_1, i + b_2, \ldots, i + b_{2m-3}, i + b_{2m-2} + n, i + b_{2m-1} + 2n), \\ F_{i,2m} &= (i + b_1, i + b_2, \ldots, i + b_{2m-2}, i + b_{2m-1}, i + b_{2m} + n) \end{aligned}$$

are cycles ($(2m-1)$-cycles and $(2m)$-cycles respectively) with vertices from $\mathbb{Z}_{2(3^{m-1}+2n-1)}$ and $\mathbb{Z}_{3^m+2n-1}$ respectively. Clearly, there are $2m$–1 faces through each vertex in $M_{2m-1,2(3^{m-1}+2n-1)}$ and there are $2m$ faces through each vertex in $M_{2m,3^m+2n-1}$. So, $f_1(M_{2m-1,2(3^{m-1}+2n-1)}) = (3^{m-1}+2n-1)(2m-1)$ and $f_1(M_{2m,3^m+2n-1}) = (3^m+2n-1)m$. Thus, $\chi(M_{2m-1,2(3^{m-1}+2n-1)}) = (3^{m-1}+2n-1)(5-2m)$ and $\chi(M_{2m,3^m+2n-1}) = (3^m+2n-1)(2-m)$. By Lemma 2 below, $M_{2m+1,2(3^{m-1}+2n-1)}$ and $M_{2m,3^m+2n-1}$ are polyhedral maps. But, by Lemma 4, none of these polyhedral maps are combinatorially regular.



$M_{5,4r}$

Lemma 1. *For a collection $\mathcal{C}$ of cycles, let $\bar{\mathcal{C}}$ be the 2-dimensional pure simplicial complex whose 2-faces are of the form xyF, where $F \in \mathcal{C}$ and xy is an edge in F. If $B(\mathcal{C})$ is as defined earlier then the following three are equivalent.*

(i) *$B(\mathcal{C})$ is a combinatorial 2-manifold.*

(ii) *$\bar{\mathcal{C}}$ is a combinatorial 2-manifold.*

(iii) *For any vertex v, the cycles containing v are of the form $F_1 = (v, v_{1,1}, \ldots, v_{1,n_1}), \ldots, F_m = (v, v_{m,1}, \ldots, v_{m,n_m})$ such that $v_{1,n_1} = v_{2,1}, \ldots, v_{m-1,n_{m-1}} = v_{m,1}$, $v_{m,n_m} = v_{1,1}$ for some $m \geq 2$.*

Proof. Clearly, $B(\mathcal{C})$ is a subdivision of $\bar{\mathcal{C}}$. Therefore, (i) and (ii) are equivalent.

For a 2-dimensional pure simplicial complex X, the link of a vertex v is the graph $\mathrm{lk}_X(v)$ whose vertex-set is $\{u \in V(X) : uv \in X\}$ and edge-set is $\{xy : xyv \in X\}$. Clearly, X is a combinatorial 2-manifold if and only if $\mathrm{lk}_X(v)$ is a cycle for each $v \in V(X)$.

Let v be a vertex of $\bar{\mathcal{C}}$. If $v = F \in \mathcal{C}$ then $\mathrm{lk}_{\bar{\mathcal{C}}}(v)$ is F itself. Let v be a vertex of $\bar{\mathcal{C}}$ which is not a cycle in $\mathcal{C}$. If the cycles containing v are of the form $F_1 = (v, v_{1,1}, \ldots, v_{1,n_1}), \ldots, F_m = (v, v_{m,1}, \ldots, v_{m,n_m})$ such that $v_{1,n_1} = v_{2,1}, \ldots, v_{m-1,n_{m-1}} = v_{m,1}$, $v_{m,n_m} = v_{1,1}$ for some $m \geq 2$ then $\mathrm{lk}_{\bar{\mathcal{C}}}(v)$ is the cycle $v_{1,1}F_1v_{2,1}F_2 \cdots v_{m,1}F_m$. Conversely, if $\mathrm{lk}_{\bar{\mathcal{C}}}(v)$ is a cycle then, from the definition of $\bar{\mathcal{C}}$, $\mathrm{lk}_{\bar{\mathcal{C}}}(v)$ must be of the form $v_{1,1}F_1v_{2,1}F_2 \cdots v_{m,1}F_m$, where $F_1 = (v, v_{1,1}, \ldots, v_{1,n_1}), \ldots, F_m = (v, v_{m,1}, \ldots, v_{m,n_m})$ such that $v_{1,n_1} = v_{2,1}, \ldots, v_{m-1,n_{m-1}} = v_{m,1}$, $v_{m,n_m} = v_{1,1}$. This proves that (ii) and (iii) are equivalent. □

Lemma 2. *$M_{2m-1,2(3^{m-1}+2n-1)}$ and $M_{2m,3^m+2n-1}$ are polyhedral maps for $m \geq 3$, $n \geq 0$.*

Proof. Since $\{i, i+1\}$ is an edge in $M_{2m-1,2(3^{m-1}+2n-1)}$ for each i, $\mathrm{EG}(M_{2m-1,2(3^{m-1}+2n-1)})$ is connected. Similarly, $\mathrm{EG}(M_{2m,3^m+2n-1})$ is connected.

Observe that the faces in $M_{2m-1,2(3^{m-1}+2n-1)}$ containing i are $F_i, F_{i-b_2}, F_{i-b_3}, F_{i-b_4}, \ldots, F_{i-b_{2m-3}}, F_{i-b_{2m-2}-n}, F_{i-b_{2m-1}-2n}$, where $F_i = F_{i,2m-1} = (i+b_1, i+b_2, \ldots, i+b_{2m-3}, i+b_{2m-2}+n, i+b_{2m-1}+2n)$. Clearly, $F_i \cap F_{i-b_3} = \cdots = F_i \cap F_{i-b_{2m-2}-n} = \cdots = F_{i-b_{2m-1}-2n} \cap F_{i-b_2} = \cdots = F_{i-b_{2m-1}-2n} \cap F_{i-b_{2m-3}} = \{i\}$.

Since $b_{2j+1} = 2b_{2j} - b_{2j-1}$ for all j, $F_{i-b_{2l-1}} \cap F_{i-b_{2l}}$ is the edge $\{i, i+b_{2l}-b_{2l-1}\}$, $F_{i-b_{2l}} \cap F_{i-b_{2l+1}}$ is the edge $\{i+b_{2l}-b_{2l+1}, i\}$ for $1 \le l \le m-2$, $F_{i-b_{2m-3}} \cap F_{i-b_{2m-2}-n}$ is the edge $\{i, i+b_{2m-1}-b_{2m-2}+n\}$ and $F_{i-b_{2m-2}-n} \cap F_{i-b_{2m-1}-2n}$ is the edge $\{i+b_{2m-3}-b_{2m-2}-n, i\}$. Again, since $2b_{2m-1}+4n \equiv 0 \pmod{2(3^{m-1}+2n-1)}$, $F_{i-b_{2m-1}-2n} \cap F_i$ is the edge $\{i, i+b_{2m-1}+2n\}$. Thus, any pair of faces containing i intersects in either at i or on an edge through i and the faces containing i form a single cycle of adjacent faces (sharing a common edge). Therefore, $M_{2m-1,2(3^{m-1}+2n-1)}$ is a polyhedral map.

The faces in $M_{2m,3^m+2n-1}$ containing i are $C_i, C_{i-b_2}, C_{i-b_3}, \ldots, C_{i-b_{2m-1}}, C_{i-b_{2m}-n}$, where $C_i = F_{i,2m} = (i+b_1, i+b_2, \ldots, i+b_{2m-1}, i+b_{2m}+n)$ and $C_i \cap C_{i-b_3} = \cdots = C_i \cap C_{i-b_{2m-1}} = \cdots = C_{i-b_{2m}-n} \cap C_{i-b_2} = \cdots = C_{i-b_{2m}-n} \cap C_{i-b_{2m-2}} = \{i\}$. Also, since $2b_{2m} - b_{2m-1} + 2n \equiv 0 \pmod{3^m+2n-1}$, $C_{i-b_{2l-1}} \cap C_{i-b_{2l}}$ is the edge $\{i, i+b_{2l}-b_{2l-1}\}$, $C_{i-b_{2l}} \cap C_{i-b_{2l+1}}$ is the edge $\{i+b_{2l}-b_{2l+1}, i\}$ for $1 \le l \le m-1$, $C_{i-b_{2m-1}} \cap C_{i-b_{2m}-n}$ is the edge $\{i, i-b_{2m}-n\}$ and $C_{i-b_{2m}-n} \cap C_i$ is the edge $\{i+b_{2m}+n, i\}$. Thus, any pair of faces containing i intersects in either at i or on an edge through i and the faces containing i form a single cycle of adjacent faces. Therefore, $M_{2m,3^m+2n-1}$ is a polyhedral map. □

From the uniqueness of 16-vertex $\{5,5\}$-equivelar polyhedral map it follows that $M_{5,16}$ is self dual. Here we prove.

Lemma 3. *$M_{2m-1,2(3^{m-1}+2n-1)}$ and $M_{2m,3^m+2n-1}$ are self dual for $m \ge 3$ and $n \ge 0$.*

Proof. Let $\varphi\colon M_{2m-1,2(3^{m-1}+2n-1)} \to \widetilde{M}_{2m-1,2(3^{m-1}+2n-1)}$ be the mapping given by $\varphi(i) = F_i := F_{-i,2m-1}$. Clearly φ is a bijection. Consider the face $F_i = (i+b_1, \ldots, i+b_{2m-3}, i+b_{2m-2}+n, i+b_{2m-1}+2n)$. Now, $(\varphi(i+b_1), \ldots, \varphi(i+b_{2m-3}), \varphi(i+b_{2m-2}+n), \varphi(i+b_{2m-1}+2n)) = (F_{-i-b_1}, \ldots, F_{-i-b_{2m-3}}, F_{-i-b_{2m-2}-n}, F_{-i-b_{2m-1}-2n}) = \widetilde{F}_{-i}$ (say). From the proof of Lemma 2, $\widetilde{F}_{-i}$ is a cycle of adjacent faces (sharing a common edge) containing the common vertex $-i$. Therefore, by the definition, $\widetilde{F}_{-i}$ is a face of $\widetilde{M}_{2m-1,2(3^{m-1}+2n-1)}$. This implies that $\widetilde{M}_{2m-1,2(3^{m-1}+2n-1)}$ is isomorphic to $M_{2m-1,2(3^{m-1}+2n-1)}$. Similarly, $\psi\colon M_{2m,3^m+2n-1} \to \widetilde{M}_{2m,3^m+2n-1}$, given by $\psi(i) = F_{-i,2m}$ defines an isomorphism. □

Clearly, $\Gamma(M_{2m-1,2(3^{m-1}+2n-1)})$ and $\Gamma(M_{2m,3^m+2n-1})$ are transitive on the vertices and the faces. Here we prove.

Lemma 4. *$M_{2m-1,2(3^{m-1}+2n-1)}$ and $M_{2m,3^m+2n-1}$ are not combinatorially regular for all $m \ge 3$ and $n \ge 0$.*

Proof. Let $\mu = 2(3^{m-1}+2n-1)$. If $m > 3$ then consider the flags $\mathcal{F}_1 = (0, \{0, b_m - b_{m+1}\}, F_{-b_{m+1}})$ and $\mathcal{F}_2 = (0, \{0, b_{m+2} - b_{m+1}\}, F_{-b_{m+1}})$ in $M_{2m-1,\mu}$. If possible let there exist $\varphi \in \Gamma(M_{2m-1,\mu})$ such that $\varphi(\mathcal{F}_1) = \mathcal{F}_2$. Then $\varphi(0) = 0$, $\varphi(F_{-b_{m+1}}) = F_{-b_{m+1}}$ and hence

$\varphi(1-b_{m+1}) = -b_{m+1}$ and $\varphi(1) = 1$. If $m > 5$ then, by considering the faces containing 1, $\varphi(F_{1-b_{m+2}}) = F_{1-b_{m+2}}$, $\varphi(F_{1-b_{m+1}}) = F_{1-b_{m+3}}$. These imply $1 + b_4 - b_{m+3} = \varphi(1-b_{m+1}) = -b_{m+1}$ in $\mathbb{Z}_\mu$, a contradiction. If $m = 5$ then $\varphi(F_{1-b_6}) = F_{1-b_8-n}$ and hence $1 + b_4 - b_8 - n = \varphi(1-b_6) = -b_6$ in $\mathbb{Z}_\mu$. This is not possible. If $m = 4$ then $\varphi(F_{1-b_5}) = F_{1-b_7-2n}$ and hence $1 + b_4 - b_7 - 2n = \varphi(1-b_{-5}) = -b_5$ in $\mathbb{Z}_\mu$, a contradiction.

For $m = 3$, if $\psi \in \Gamma(M_{5,\mu})$ such that $\psi((0,\{0,3+n\},F_{-b_4-n})) = (0,\{0,13+3n\},F_{-b_4-n})$ then $\psi(12+3n) = 11+3n$ and $\psi(F_{1-b_4-n}) = F_1$ and hence $3 = \psi(12+3n) = 11+3n$ in $\mathbb{Z}_\mu$. This is also not possible.

Thus, $M_{2m-1,2(3^{m-1}+2n-1)}$ always has a pair of flags $\mathcal{F}_1$ and $\mathcal{F}_2$ such that $\varphi(\mathcal{F}_1) \neq \mathcal{F}_2$ for all $\varphi \in \Gamma(M_{2m-1,2(3^{m-1}+2n-1)})$. So, $M_{2m-1,2(3^{m-1}+2n-1)}$ is not combinatorially regular.

Let $\eta = 3^m + 2n - 1$ and $C_i = F_{i,2m}$. Consider the flags $\mathcal{C}_1 = (0,\{0,(-1)^m(b_{m+2} - b_{m+1})\},C_{-b_{m+1}})$ and $\mathcal{C}_2 = (0,\{0,(-1)^m(b_{m+2}-b_{m+1})\},C_{-b_{m+2}})$ in $M_{2m,\eta}$. If $\varphi \in \Gamma(M_{2m,\eta})$ such that $\varphi(\mathcal{C}_1) = \mathcal{C}_2$, then $\varphi(\mathcal{C}_2) = \mathcal{C}_1$, $\varphi(1-b_{m+2}) = -b_{m+1}$ and $\varphi(1) = 1$. If $m > 3$ then $\varphi(C_{1-b_{m+2}}) = C_{1-b_{m+3}}$ and hence $1+b_2-b_{m+3} = \varphi(1-b_{m+2}) = -b_{m+1}$ in $\mathbb{Z}_\eta$, a contradiction. If $m = 3$ then $\varphi(C_{1-b_5}) = C_{1-b_6-n}$ and hence $1+b_2-b_6+n = \varphi(1-b_5) = -b_4$ in $\mathbb{Z}_\eta$. This is not possible. Therefore, by similar argument as before, $M_{2m,3^m+2n-1}$ is not combinatorially regular. □

Example 2. Let $\mathcal{C}_4$ be the collection of 4-cycles of the complete graph K_5 on the vertex set $\mathbb{Z}_4 \cup \{u\}$ given by $\mathcal{C}_4 = \{(0,1,2,3),(u,i,i+1,i+3) \ : i \in \mathbb{Z}_4\}$. Then $|\bar{\mathcal{C}_4}|$ is the torus and hence $(|\bar{\mathcal{C}_4}|, K_5)$ is a non-singular $\{4,4\}$-pattern.

Lemma 5. *Suppose* $\mathcal{C}(\pi_p) = \{(0,1,\ldots,p-1),(u,i+\pi_p(1),\ldots,i+\pi_p(p-1)) \ : \ i \in \mathbb{Z}_p\}$ *is a collection of cycles of the complete graph* K_{p+1} *on the vertex set* $\mathbb{Z}_p \cup \{u\}$*, where* p *is an odd prime and* π_p *is a permutation of* $\mathbb{Z}_p \setminus \{0\} = \{1,\ldots,p-1\}$*. If*

(pp1) $\pi_p(i) + \pi_p(p-i) = p$ *for* $1 \leq i \leq p-1$,

(pp2) $\pi_p(\frac{p-1}{2}) = \frac{p-1}{2}$ *and*

(pp3) *exactly one of* j, $-j$ *is in* $\{\pi_p(2)-\pi_p(1), \pi_p(3)-\pi_p(2),\ldots,\pi_p(\frac{p+1}{2})-\pi_p(\frac{p-1}{2})\}$

then $\bar{\mathcal{C}}(\pi_p)$ *is a connected combinatorial 2-manifold.*

Proof. Since edges of cycles of $\mathcal{C}(\pi_p)$ form a connected graph, $\mathrm{EG}(\bar{\mathcal{C}}(\pi_p))$ is connected.

Let $a_i = \pi_p(i+1) - \pi_p(i)$ for $1 \leq i \leq p-2$. Then, by (pp1), $a_i = a_{p-1-i}$. Let $r = \frac{p-3}{2}$. Then, by (pp1), (pp2), $a_{r+1} = 1$ and, by (pp3), $\{a_1,\ldots,a_{r+1},-a_1,\ldots,-a_{r+1}\} = \mathbb{Z}_p \setminus \{0\}$.

If r is even then the cycles containing i are $(i,u,\ldots,i+a_1)$, $(i,i+a_1,\ldots,i-a_2)$, $(i,i-a_2,\ldots,i+a_3)$, $\ldots$, $(i,i+a_{r-1},\ldots,i-a_r)$, $(i,i-a_r,\ldots,i+1)$, $(i,i+1,i+2,\ldots,i+p-1)$, $(i,i+p-1,\ldots,i+a_{r+2})$, $\ldots$, $(i,i+a_{2r},\ldots,i-a_{2r+1})$, $(i,i-a_{2r+1},\ldots,u)$.

If r is odd then the cycles containing i are $(i,u,\ldots,i+a_1)$, $(i,i+a_1,\ldots,i-a_2)$, $(i,i-a_2,\ldots,i+a_3)$, $\ldots$, $(i,i-a_{r-1},\ldots,i+a_r)$, $(i,i+a_r,\ldots,i+p-1)$, $(i,i+p-1,\ldots,i+2,i+1)$, $(i,i+1,\ldots,i-a_{r+2})$, $\ldots$, $(i,i+a_{2r},\ldots,i-a_{2r+1})$, $(i,i-a_{2r+1},\ldots,u)$.

The cycles containing u are $(u,\pi_p(1),\ldots,\pi_p(p-1))$, $(u,1+\pi_p(1),\ldots,1+\pi_p(p-1))$, $\ldots$, $(u,p-1+\pi_p(1),\ldots,p-1+\pi_p(p-1))$. Since $\{\pi_p(p-1),1+\pi_p(p-1),\ldots,p-1+\pi_p(p-1)\} = \mathbb{Z}_p$, the cycles containing u can be arranged as $(u,\pi_p(i_1),\ldots,\pi_p(j_1)),\ldots,(u,\pi_p(i_p),\ldots,\pi_p(j_p))$, where $j_1 = i_2,\ldots,j_{p-1} = i_p, j_p = i_1$. The lemma now follows by Lemma 1. □

Clearly, π_3 is the identity permutation and $\mathcal{C}(\pi_3)$ is the 4-vertex 2-sphere S^2_4. Also, $\chi(\bar{\mathcal{C}}(\pi_p)) = 2(p+1) - \left(\binom{p+1}{2} + p(p+1)\right) + (p+1)p = (p+1)(4-p)/2$. So, if $p = 4k+1$ for some $k \geq 1$ then $\chi(\bar{\mathcal{C}}(\pi_p))$ is odd and hence $\bar{\mathcal{C}}(\pi_p)$ is non-orientable. Here we prove.

Lemma 6. *$\bar{\mathcal{C}}(\pi_p)$ is non-orientable for $p > 3$.*

Proof. Let $F = (0, 1, \ldots, p-1)$ and $F_i = (u, i+\pi_p(1), \ldots, i+\pi_p(p-1))$ for $1 \leq i \leq p-1$. We can choose a p-gonal disc (not necessarily convex) in the plane for each cycle in $\bar{\mathcal{C}}(\pi_p)$ so that the disc corresponding to F_i is attached with that for F along the common edge $\{i+\pi_p(\frac{p-1}{2}), i+\pi_p(\frac{p+1}{2})\}$ for each i and there are no other intersections. This gives us a $p(p-1)$-gonal disc $D(\pi_p)$. Then there are two edges in $D(\pi_p)$ corresponding to an edge jk $(j,k \in \mathbb{Z}_p,\ -1 \neq j-k \neq 1)$ in some cycle F_i and they appear in the same direction (clockwise or anti-clockwise). Since $|\bar{\mathcal{C}}(\pi_p)|$ is homeomorphic to the space obtained by identifying such pairs of edges (and some more) of $D(\pi_p)$, $|\bar{\mathcal{C}}(\pi_p)|$ is non-orientable. □



$D(\sigma_7)$

Lemma 7. *Let $p > 3$ be a prime.*

(a) *If $p = 4k+3$ for some $k \geq 1$ then the permutation $\sigma_p = (2, 4k+1)(4, 4k-1)\cdots(2k, 2k+3)$ of $\mathbb{Z}_p \setminus \{0\}$ satisfies* (pp1), (pp2) *and* (pp3) *of Lemma* 5.

(b) *If $p = 4l+1$ for some $l \geq 1$ then the permutation $\rho_p = (1, 4l)(3, 4l-2)\cdots(2l-1, 2l+2)$ of $\mathbb{Z}_p \setminus \{0\}$ satisfies* (pp1), (pp2) *and* (pp3) *of Lemma* 5.

Proof. Clearly, σ_p and ρ_p satisfy hypothesis (pp1) and (pp2).

Now, $\{\sigma_p(2) - \sigma_p(1), \ldots, \sigma_p(\frac{p+1}{2}) - \sigma_p(\frac{p-1}{2})\} = \{4k, -(4k-2), 4k-4, \ldots, 4, -2, 1\} = \{-2, 4, -6, \ldots, -(4k-2), 4k, -(4k+2)\}$. Thus σ_p satisfies (pp3).

Again, $\{\rho_p(2) - \rho_p(1), \ldots, \rho_p(\frac{p+1}{2}) - \rho_p(\frac{p-1}{2})\} = \{-(4l-2), 4l-4, -(4l-6), \ldots, 4, -2, 1\} = \{-2, 4, -6, \ldots, (4l-4), -(4l-2), -4l\}$. Thus ρ_p satisfies (pp3). □

Proof of Theorem 1. Let $m \geq 3$ and $n \geq 0$. By Lemma 2, $M_{2m-1, 2(3^{m-1}+2n-1)}$ is a $2(3^{m-1} + 2n - 1)$-vertex polyhedral map and hence a $\{2m-1, 2m-1\}$-equivelar polyhedral map. Again, by Lemma 2, $M_{2m, 3^m+2n-1}$ is a $(3^m + 2n - 1)$-vertex polyhedral map and hence a $\{2m, 2m\}$-equivelar polyhedral map. The theorem now follows from Lemma 3. □

Proof of Theorem 2. Let $p > 3$ be a prime and K_{p+1} be the complete graph on the vertex set $\mathbb{Z}_p \cup \{u\}$. By Lemma 7, there exists a permutation π_p of $\mathbb{Z}_p \setminus \{0\}$ which satisfies (pp1),

(pp2) and (pp3) of Lemma 5. Let $\mathcal{C}(\pi_p)$ be as in Lemma 5. Then, by Lemma 5, $\bar{\mathcal{C}}(\pi_p)$ is a connected combinatorial 2-manifold. So, if $N_p := |\bar{\mathcal{C}}(\pi_p)|$ then (N_p, K_{p+1}) is a non-singular $\{p, p\}$-pattern and the cycles in $\mathcal{C}(\pi_p)$ are the boundaries of the faces of (N_p, K_{p+1}). Finally, the 4-vertex 2-sphere S_4^2 gives a $\{3, 3\}$-pattern. This completes the proof. □

Acknowledgements. The author is thankful to Bhaskar Bagchi and Sunanda Bagchi for useful conversations. The author thanks the anonymous referee for many useful references and comments which helped to improve the presentation of this paper.

References

[1] Brehm, U.: *Polyhedral maps with few edges.* Topics in Comb. and Graph Theory (Ringel-Festschrift) (eds. Bodendiek, R. and Henn, R.), Physica-Verlag, Heidelberg 1990, 153–162. Zbl 0703.57008

[2] Brehm, U.; Datta, B.; Nilakantan, N.: *The edge-minimal polyhedral maps of Euler characteristic* −8. Beitr. Algebra Geom. **43** (2002), 583–596. Zbl 1017.52010

[3] Brehm, U.; Schulte, E.: *Polyhedral maps.* Handbook of Discrete and Computational Geometry (eds. Goodman, J. E. and O'Rourke, J.), CRC Press 1997, 345–358. Zbl 0920.52004

[4] Brehm, U.; Wills, J. M.: *Polyhedral manifolds.* Handbook of Convex Geometry (eds. Gruber, P. M. and Wills, J. M.), Elsevier Publishers 1993, 535–554. Zbl 0823.52014

[5] Coxeter, H. S. M.: *Regular skew polyhedra in three and four dimensions, and their topological analogues.* Proc. London Math. Soc. (2) **43** (1937), 33–62. (Reprinted with amendments in: Twelve Geometric Essays, Southern Illinois Univ. Press, Carbondale 1968.) Zbl 0016.27101

[6] Datta, B.; Nilakantan, N.: *Equivelar polyhedra with few vertices.* Discrete Comput. Geom. **26** (2001), 429–461. Zbl 1004.51028

[7] Edmonds, A. L.; Ewing, J. H.; Kulkarni, R. S.: *Regular tessellations of surfaces and* $(p, q, 2)$*-triangle groups.* Ann. Math. **116** (1982), 113–132. Zbl 0497.57001

[8] Gott III, J. R.: *Pseudopolyhedrons.* Am. Math. Mon. **74** (1967), 497–504. Zbl 0147.39201

[9] Grünbaum, B.: *Regular polytopes – Old and new.* Aeqationes Math. **16** (1977), 1–20. Zbl 0381.51012

[10] McMullen, P.; Schulte, E.: *Abstract Regular Polytopes.* Cambridge Univ. Press, 2002. Zbl 1039.52011

[11] McMullen, P.; Schulz, Ch.; Wills, J. M.: *Equivelar polyhedral manifolds in* E^3. Isr. J. Math. **41** (1982), 331–346. Zbl 0497.52004

[12] McMullen, P.; Schulz, Ch.; Wills, J. M.: *Polyhedral 2-manifolds in* E^3 *with unusually large genus.* Isr. J. Math. **46** (1983), 127–144. Zbl 0525.52008

[13] Nilakantan, N.: *Non-existence of weakly neighbourly polyhedral map of type* $\{6, 6\}$. Experimental Maths **12** (2003), 257–262.

Received May 28, 2004