

Journal URL

<http://www.math.washington.edu/~ejpecp/>

Paper URL

<http://www.math.washington.edu/~ejpecp/EjpVol7/paper12.abs.html>

A STOCHASTIC TWO-POINT BOUNDARY VALUE PROBLEM

S. J. Luo

FinancialCAD Corp., 100-7455-132 St., Surrey, B.C., Canada

John B. Walsh

Department of Mathematics, University of British Columbia, Vancouver, B.C. B6T 1Y4

walsh@math.ubc.ca

Abstract We investigate the two-point stochastic boundary-value problem on $[0, 1]$:

$$\begin{aligned}U'' &= f(U)\dot{W} + g(U, U') \\U(0) &= \xi \\U(1) &= \eta.\end{aligned}\tag{0}$$

where \dot{W} is a white noise on $[0, 1]$, ξ and η are random variables, and f and g are continuous real-valued functions.

This is the stochastic analogue of the deterministic two point boundary-value problem, which is a classical example of bifurcation.

We find that if f and g are affine, there is no bifurcation: for any r.v. ξ and η , (0) has a unique solution a.s. However, as soon as f is non-linear, bifurcation appears. We investigate the question of when there is either no solution whatsoever, a unique solution, or multiple solutions. We give examples to show that all these possibilities can arise. While our results involve conditions on f and g , we conjecture that the only case in which there is no bifurcation is when f is affine.

Keywords Stochastic boundary-value problems, bifurcations

AMS subject classification (2000) Primary. 60H10; Secondary. 34F05

Submitted to EJP on November 27, 2000. Final version accepted on September 14, 2001.

1 Introduction

This problem arose from an attempt to solve the stochastic Dirichlet problem on a nice domain D in \mathbb{R}^2 with multiplicative white noise:

$$\begin{aligned}\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} &= f(U(x, y)) \dot{W}(x, y), \quad (x, y) \in D \\ U(x, y) &= 0, \quad (x, y) \in \partial D.\end{aligned}$$

where f is a smooth function and \dot{W} is a white noise on \mathbb{R}^2 . Early calculations showed that something goes very wrong for most f 's, and we decided to look at a simpler situation to understand why. The simplest non-trivial setting of this problem is in one dimension, where it becomes the two-point boundary value problem:

$$\begin{aligned}\frac{d^2 U}{dt^2} &= f(U_t) \circ \dot{W}_t + g(U_t, U'_t), \quad 0 < t < 1, \\ U_0 &= \xi, \quad U_1 = \eta,\end{aligned}\tag{1.1}$$

where \dot{W} is a white noise on $[0, 1]$, ξ and η are random variables, and the little circle indicates Stratonovich integration.

It was immediately clear why the two-dimensional problem was pathological: even in the simplest non-stochastic setting, this problem exhibits bifurcation, and the two-dimensional problem can be expected to be much worse. Indeed, consider the non-stochastic problem

$$\begin{aligned}u''(t) &= \lambda g(u(t)), \quad 0 < t < 1, \\ u(0) &= u(1) = 0,\end{aligned}\tag{1.2}$$

where $g(0) = 0$ and $g'(0) > 0$. This always has the solution $u \equiv 0$, but as λ increases, it bifurcates, and at certain λ there are multiple nontrivial solutions.

The question is then how the presence of noise affects this bifurcation. It turns out that it regularizes the problem to a certain extent. If f and g are affine, for instance, and f does not vanish, the solution of (1.1) exists and is a.s. unique. However, as soon as the coefficients are non-linear, bifurcation seems to re-appear: there will be a positive probability of multiple solutions. (“Bifurcation” may not be the correct word here: there is no parameter λ in the problem, but its place is essentially taken by the stochastic variable ω .)

The deterministic version of the problem (where $f \equiv 0$) has been extensively studied. See [8] and the references therein. Variations of the stochastic problem have been studied before by several authors. Ocone and Pardoux [7], generalizing and extending work of Krener [2], considered systems of SDE's with general linear boundary conditions and affine functions f and g . Under some non-degeneracy conditions, they showed that there exists a unique solution which satisfies the sharp Markov property.

Dembo and Zeitouni [1] studied the corresponding Neumann problem and showed the existence of a global solution for f and g satisfying certain properties.

Nualart and Pardoux [4] looked at the case where f was constant, and showed that under certain conditions on g there exists a unique global solution. Moreover, under these conditions, the solution has the sharp Markov field property if and only if g is affine.

In this paper we concentrate on the case where f is not constant. It is rather rare to have a unique global solution: basically, that happens only when f is affine. Existence is more common: if f and g are asymptotically linear, for instance, a global solution exists, but it may not be unique unless f and g are actually affine. We give an example in which there is no global solution at all, and others in which there are sets on which there is one solution, exactly two solutions, or many solutions.

The equations (1.1) are shorthand for the integral equations

$$\begin{cases} U_t &= \xi + Yt + \int_0^t \int_0^s f(U_u) \circ dW_u ds + \int_0^t \int_0^s g(U_u, U'_u) du ds, & 0 \leq t \leq 1, \\ U_1 &= \eta, \end{cases}$$

where $\{W_t, t \geq 0\}$ is a standard Brownian motion, the circle denotes Stratonovich integration and Y is the (unknown) value of U'_0 . (White noise is just the derivative—in the distribution sense—of Brownian motion. We use it here to write the differential equation in an intuitive form, but it is the integral equation which has rigorous meaning.)

It is convenient to restate the above as a system by letting $V = U'$:

$$\begin{cases} U_t &= \xi + \int_0^t V_s ds, & 0 < t < 1 \\ V_t &= Y + \int_0^t f(U_s) \circ dW_s + \int_0^t g(U_s, V_s) ds, & 0 \leq t \leq 1 \\ U_1 &= \eta. \end{cases} \quad (1.3)$$

Here, the random variables ξ and η are given, but Y , which is the initial value of U' , is not.

Allowing g to depend on V as well as on U causes no extra difficulty; allowing f to depend on V , however, would fundamentally alter the problem.

All these statements of the problem are equivalent. Thus, if Ω_0 is a measurable set with $P\{\Omega_0\} > 0$, we say that a pair (U, V) of processes is a *local solution* of (1.1) (or of (1.3), or (1.3)) *on* Ω_0 if it satisfies (1.3) a.e. on Ω_0 . We say it is a *global solution* if $P\{\Omega_0\} = 1$. Similarly, we say that a process U is a *local* (resp. *global*) solution if there is a process V such that the pair (U, V) is a local (resp. global) solution. (Notice that we use the words “local” and “global” in a slightly non-standard sense: they refer to the probability space, not the parameter space.)

2 The Main Results

Let (Ω, \mathcal{F}, P) be a probability space and let $(\mathcal{F}_t) \subset \mathcal{F}$ be a right-continuous filtration. Let $\{W_t, t \geq 0\}$ be a standard Brownian motion adapted to (\mathcal{F}_t) .

Here are the main results of the paper. First, in the linear case, we have the following existence and uniqueness result.

Theorem 2.1 *Suppose f and g are affine functions, with, say, $f(x) = \alpha x + \beta$ and $g(x, y) = \gamma x + \kappa y + \lambda$, where $\alpha, \beta, \gamma, \kappa$, and λ are constants. If f is not identically zero, or if f vanishes identically and $\gamma + \frac{\kappa^2}{4} \neq -k^2\pi^2$, $k = 1, 2, \dots$, then for any random variables ξ and η , (1.3) has a unique global solution.*

Remark 2.2 If $f \equiv 0$, the equation is deterministic and the second condition just rules out the well-known case in which uniqueness fails. Notice that once f is non-trivial, the white noise regularizes the problem, i.e. there is always a unique global solution.

Moving on to the non-linear case, we look at the existence of global solutions. They will always exist if f and g are asymptotically linear.

Theorem 2.3 *Suppose f and g are differentiable, and their first (partial) derivatives are bounded and Hölder continuous. Suppose further that there exist constants a , c , and d such that*

$$\lim_{|x| \rightarrow \infty} \frac{f(x) - ax}{x} = \lim_{x^2 + y^2 \rightarrow \infty} \frac{g(x, y) - cx - dy}{\sqrt{x^2 + y^2}} = 0. \quad (2.4)$$

and either $a \neq 0$ or $a = 0$ and $c + \frac{d^2}{4} \neq -k^2\pi^2$, $k = 1, 2, \dots$.

Then for any random variables ξ and η , (1.3) has at least one global solution.

Remark 2.4 This covers a fairly large class of coefficients. If f and g are bounded, or if they just increase more slowly than linearly, then a , c , and d will vanish, and the theorem implies that there is always at least one global solution.

There are no adaptedness restrictions on the random variables ξ and η in Theorem 2.3. In fact there is a trade-off between adaptedness restrictions and regularity conditions on the coefficients. If we assume some adaptedness of ξ and η we can weaken the regularity hypotheses somewhat.

Theorem 2.5 *Suppose f and g are Lipschitz continuous functions satisfying (2.4). Then for any \mathcal{F}_0 -measurable r.v. ξ and any random variable η , (1.3) has at least one global solution.*

Remark 2.6 By time-reversal, the same conclusion holds if ξ is arbitrary and η is independent of \mathcal{F}_1 .

Let us now look at some conditions which ensure local existence and uniqueness. In the deterministic case, the two-point boundary-value problem has a unique solution if the interval in question is small enough. In the stochastic case, this remains true if the interval is short and the Brownian motion is small. Again there is a trade-off, this time between the length of the interval and the Lipschitz constants of f and g . Thus, let us pose the problem on the interval $[0, T]$ for a fixed $T > 0$.

$$\begin{cases} U_t'' &= f(U_t) \circ \dot{W}_t + g(U_t, U_t'), & 0 < t < T, \\ U_0 &= \xi, \quad U_T = \eta. \end{cases} \quad (2.5)$$

The first theorem shows that if f is relatively smooth and g is Lipschitz with a small enough Lipschitz constant, there exists a unique local solution.

Theorem 2.7 *Let $f \in C^{(1)}$ with f' Lipschitz, and suppose g is Lipschitz with Lipschitz constant L_g . If $L_g < \left(\frac{T^2}{8} + \frac{3T}{2}\right)^{-1}$, then there is a set $\Omega_0 \subset \Omega$ of positive probability such that for any random variables ξ and η , (2.5) has a unique solution on Ω_0 . If f and g are fixed, $P\{\Omega_0\} \rightarrow 1$ as $T \rightarrow 0$.*

Remark 2.8 Only the Lipschitz constant of g appears explicitly. The Lipschitz constant of f is implicit: Ω_0 will be a set on which W_t is small. The smaller the Lipschitz constants of f and f' , the larger W_t can be.

If the solution is not unique, it is natural to ask how many solutions there can be. We will restrict ourself to the case where $g = 0$. Consider the problem

$$\begin{cases} U_t'' = f(U_t) \circ \dot{W}_t, & 0 < t < 1, \\ U_0 = U_1 = x_0, \end{cases} \quad (2.6)$$

Theorem 2.9 *Suppose that $f \in C^{(1)}$, f has at least one zero, and f' is Lipschitz continuous.*

(i) *If f is not an affine function, then there exists a point x_0 such that the problem (2.6) has at least two global solutions.*

(ii) *Suppose that f has an isolated zero at x_0 . Let $\rho > 0$ be the distance from x_0 to the nearest other zero of f . If f is not affine on $(x_0 - \rho, x_0 + \rho)$, then for each integer $k \geq 1$ there exists a set $\Omega_k \subset \Omega$ of positive probability such that on Ω_k there are at least k distinct local solutions of (2.6).*

If we put some small conditions on f , we can show that there is global existence and uniqueness for the two-point boundary value problem if and only if f is affine. We conjecture that this is true in general, with no added conditions on f .

Theorem 2.10 *Let f be a differentiable function with bounded derivative. Suppose that f has at least one zero, is not identically zero, and its derivative is Lipschitz continuous.*

Then the equations

$$\begin{cases} U_t'' = f(U_t) \circ \dot{W}_t \\ U_0 = \xi, \\ U_1 = \eta, \end{cases} \quad (2.7)$$

have a unique global solution for all pairs of r.v. ξ and η if and only if f is affine.

Remark 2.11 A process $\{X_t, 0 \leq t \leq 1\}$ satisfies Lévy's sharp Markov property (SMP) if for each subinterval $[a, b] \subset [0, 1]$, $\{X_t, t \in [a, b]\}$ and $\{X_t, t \notin [a, b]\}$ are conditionally independent given X_a and X_b .

We do not know how Nualart and Pardoux' results [4] extend to non-constant f , but Theorems 2.9 and 2.10 strongly suggest that the SMP will fail for the solution(s) U_t of (2.5) if f is not affine. In fact, Theorem 2.9 shows that U_t cannot have a transition probability, i.e. the conditional distribution of $\{U_t, t \in [a, b]\}$ is not determined by U_a and U_b . Indeed, if it were, the solution to

(2.5) would have to be unique in law. (Take $[a, b] = [0, 1]$, so that U_a and U_b are deterministic, and the conditional and unconditional distributions are the same.) But it is not: by Theorem 2.9 there are non-trivial solutions as well as the constant solution $U_t \equiv x_0$.

Here is a class of equations for which there are no global solutions.

Theorem 2.12 *Suppose f and g are Lipschitz, and that $f'(x)$ exists for large enough x . Suppose there exist constants a_{\pm} , b_{\pm} and c_{\pm} such that*

$$\lim_{x \rightarrow -\infty} f'(x) = a_{-}, \quad \lim_{x \rightarrow \infty} f'(x) = a_{+},$$

and

$$\lim_{x \rightarrow -\infty} \frac{g(x, y) - b_{-}x - c_{-}y}{\sqrt{x^2 + y^2}} = \lim_{x \rightarrow \infty} \frac{g(x, y) - b_{+}x - c_{+}y}{\sqrt{x^2 + y^2}} = 0$$

uniformly in y . If either $a_{-} \neq a_{+}$, or else $a_{-} = a_{+}$ and $4b_{-} + c_{-}^2 \neq 4b_{+} + c_{+}^2$, then there exists a set Ω_0 of positive probability and an unbounded interval (α, β) such that (1.3) with the boundary conditions $U_0 = \xi = 0$ and $U_1 = \eta \in (\alpha, \beta)$ has no solution on Ω_0 .

3 The Initial Value Problem

A standard method of solving the two-point boundary-value problem is to pose it as an initial value problem, and then vary the initial derivative to get the desired value at the end point. Accordingly, we will consider the initial-value problem

$$\begin{cases} U_t'' &= f(U_t) \circ \dot{W}_t + g(U_t, U_t'), \quad t > 0, \\ U_0 &= x \\ U_0' &= y \end{cases} \quad (3.8)$$

which is equivalent to the system of integral equations:

$$\begin{cases} U_t &= x + \int_0^t V_s ds \\ V_t &= y + \int_0^t f(U_s) \circ dW_s + \int_0^t g(U_s, V_s) ds. \end{cases} \quad (3.9)$$

Suppose that both f and g are Lipschitz continuous functions. Evidently V is locally integrable—otherwise the equations don't make sense—so that U must be absolutely continuous. Since f is Lipschitz, $t \mapsto f(U_t)$ is again absolutely continuous, hence a.e. differentiable, and it is easy to verify that for a.e. t its derivative is $f'(U_t)V_t$. Thus, following [5], we can integrate by parts in the Stratonovich integral to see that (3.8) is equivalent to

$$\begin{cases} U_t &= x + \int_0^t V_s ds, \\ V_t &= y + f(U_t)W_t - \int_0^t f'(U_s)V_s W_s ds + \int_0^t g(U_s, V_s) ds. \end{cases} \quad (3.10)$$

Proposition 3.1 *The equations (3.9) and (3.10) have the same set of solutions.*

One consequence of this is that the initial-value problem can be solved omega-by-omega: the exceptional null sets involved in the definition of the stochastic integrals do not depend on the initial conditions. Denote the solutions of (3.10) by $U_t(x, y)$ and $V_t(x, y)$. We then see from Kunita's Theorem [3] that

Theorem 3.2 (Kunita) *If f and g are Lipschitz, then for a.e. ω we have*

(a) *the pair $(U_t(x, y, \omega), V_t(x, y, \omega))$ is continuous in (x, y, t) ;*

(b) *for a.e. ω , we have for all t that $(x, y) \mapsto (U_t(x, y, \omega), V_t(x, y, \omega))$ is a homeomorphism of \mathbb{R}^2 onto itself.*

Moreover, if f and g are n -times Lipschitz continuously differentiable, then $U_t(x, y)$ and $V_t(x, y)$ are n -times continuously differentiable in (x, y) , and if p and q are positive integers with $p+q \leq n$, and if $D^{p,q} = \frac{\partial^{p+q}}{\partial x^p \partial y^q}$, $\dot{U} = D^{p,q}U(x, y)$ and $\dot{V} = D^{p,q}V(x, y)$, then \dot{U} and \dot{V} satisfy

$$\begin{cases} d\dot{U}_t &= \dot{V}_t dt \\ d\dot{V}_t &= D^{p,q}(f(U_t(x, y))) \circ dW_t + D^{p,q}(g(V_t(x, y))) dt \\ \dot{U}_0 &= D^{p,q}x, \quad \dot{V}_0 = D^{p,q}y. \end{cases} \quad (3.11)$$

We say that $(U_t(x, y), V_t(x, y))$ is a *flow of homeomorphisms*. This flow of homeomorphisms will be adapted if the initial values are deterministic. Moreover, as the integrands are absolutely continuous, the stochastic integrals in (3.10) can be interpreted as either Stratonovich or Ito integrals—their values will be the same either way. However, if the initial values are random, the flow need not be adapted and the Ito integrals may no longer be defined. Nevertheless, they will still be well-defined as Stratonovich integrals, and we have

Proposition 3.3 *Let $f \in C(\mathbb{R})$ and $g \in C(\mathbb{R}^2)$ be Lipschitz continuous functions, and let X and Y be random variables. Then there exists a solution (U, V) of (3.10), and, moreover, with probability one, $(U_t, V_t) = (U_t(X, Y), V_t(X, Y))$ is a solution, where $\{(U_t(x, y), V_t(x, y)), t \geq 0, (x, y) \in \mathbb{R}^2\}$ is the flow generated by (3.8). If, moreover, f' is Lipschitz continuous, then this solution is unique.*

Remark 3.4 There is a nuance here. We know from Kunita's theorem that the flow is unique among *adapted* flows. However, we need it to be unique among *all*—including non-adapted—flows. Since (3.10) is an ordinary integral equation, no stochastic integrals enter, so that if we choose ω so that $t \rightarrow W_t(\omega)$ is continuous, the theory of (non-stochastic) integral equations guarantees that the solution will be unique if f' and g are Lipschitz. This unique solution necessarily coincides with the solution guaranteed by Kunita's theorem.

4 The linear case: proof of Theorem 2.1

We will prove the existence and uniqueness of the solution of the two-point boundary value problem in the linear case. This has been treated by Ocone and Pardoux [7] in a setting which is more general than ours in some respects, but which requires that the boundary values be regular in a Malliavin calculus sense. These conditions turn out to be unnecessary in our case. As our result is rather simple, we will give a separate proof here.

Let f and g be affine, say, $f(x) = \alpha x + \beta$ and $g(x, y) = \gamma x + \kappa y + \lambda$ where $\alpha, \beta, \gamma, \kappa$ and λ be real numbers. Then the two-point boundary value problem (1.1) becomes

$$\begin{cases} dU_t &= V_t dt \\ dV_t &= (\alpha U_t + \beta) \circ dW_t + (\gamma U_t + \kappa V_t + \lambda) dt \\ U_0 &= \xi, \quad U_1 = \eta. \end{cases}$$

We will use the initial value problem to solve this. Set $U_0 = \xi$. To show the existence of a solution, we must prove that we can choose V_0 to make $U_1 = \eta$. To show uniqueness, we must show there is only one possible value of V_0 .

Let X and Y be random variables, and consider the initial value problem (3.9) with random initial conditions:

$$\begin{cases} U_t &= X + \int_0^t V_s ds \\ V_t &= Y + \int_0^t (\alpha U_s + \beta) \circ dW_s + \int_0^t (\gamma U_s + \kappa V_s + \lambda) ds. \end{cases} \quad (4.12)$$

By Proposition 3.3 there is a solution of this initial-value problem for any X and Y , and it is unique.

Since the problem is linear, we can decompose any solution of (4.12) into the sum of a particular solution plus a linear combination of solutions of the associated homogeneous equation. We will simplify our notation slightly by writing $Z_t = (U_t, V_t)$.

Thus any solution $Z = (U, V)$ of the system (4.12) can be written uniquely in the form

$$Z_t = Z^p + XZ^{10} + YZ^{01} \quad (4.13)$$

where Z^p is a solution of (4.12) with initial value $Z_0^p = (0, 0)$, and Z^{10} and Z^{01} are solutions of the associated homogeneous equation—i.e. (4.12) with $\beta = \lambda = 0$ —with initial values $Z^{10} = (1, 0)$ and $Z^{01} = (0, 1)$ respectively. Since $Z_0 = (X, Y)$, evidently $X = \xi$. Then $\eta = U_1 = U_1^p + \xi U_1^{10} + Y U_1^{01}$, so that as long as $U_1^{01} \neq 0$,

$$Y = \frac{\eta - U_1^p - \xi U_1^{10}}{U_1^{01}}.$$

It follows that we have a unique global solution if and only if $P\{U_1^{01} = 0\} = 0$.

We have two cases. First suppose $\alpha = 0$, which is the deterministic case. The equation for U_t^{01} becomes $U'' = \gamma U + \kappa U'$ with initial conditions $U_0 = 0, U_0' = 1$. We can solve this explicitly, and we find that $U_1 = 0$ if and only if $\gamma + \kappa^2/4 = -k^2\pi^2$ for some integer k .

Otherwise, if $\alpha \neq 0$, the process Z_t is a diffusion process, and by Hörmander's theorem we see that Z_1 has a density in \mathbb{R}^2 , and hence the probability that Z_1 lies on the axis $x = 0$ is zero. This proves Theorem 2.1. \square

5 Existence of global solutions

One way to show the existence of a solution of (1.3) is to consider the vertical line $L_0 \equiv \{(x, y) : x = \xi\}$ in the plane, and follow its mapping by the flow of homeomorphisms

$Z_t = \{(U_t(x, y), V_t(x, y)), x, y \in \mathbb{R}\}$ of Theorem 3.2. For each t , Z_t maps the line L_0 into the curve $L_t \equiv Z_t(L_0) = \{(x, y) : x = U_t(\xi, u), y = V_t(\xi, u), u \in \mathbb{R}\}$. Since $(x, y) \mapsto Z_t(x, y)$ is a homeomorphism, L_t must be a non-self-intersecting continuous curve. Take $t = 1$. If L_1 has a non-empty intersection with the line $x = \eta$, then there exists $u = u(\omega)$ such that $(U_1(\xi, u), V_1(\xi, u)) \in \{x = \eta\}$, i.e. $U_1(\xi, u) = \eta$, so $U_t(\xi, u)$ is a solution of (1.3).

Thus (1.3) has a solution if and only if L_1 intersects the line $\{x = \eta\}$, and it has a unique solution if and only if this intersection is a singleton. In particular, since L_1 is continuous we have

Lemma 5.1 *A sufficient condition for the existence of a global solution of (1.3) is that the projection of L_1 on the x -axis is the whole axis.*

It is worthwhile to see how this works in the linear case. If f and g are affine functions, then by (4.13) the image of the line L_0 is $Z_1^p + \xi Z_1^{10} + y Z_1^{01}$, $-\infty < y < \infty$, which is again a line. Unless it is vertical, it will have a unique intersection with $\{x = \xi\}$, and it can only be vertical if $U_1^{01} = 0$. Except for trivial deterministic cases, this happens with probability zero.

Turning to the non-linear case, let us prove Theorems 2.3 and 2.5. We must compare the solutions of two initial-value problems. Let f, g be Lipschitz functions, let \hat{f} and \hat{g} be linear functions, and consider the solutions $Z_t = (U_t, V_t)$ and $\hat{Z}_t = (\hat{U}_t, \hat{V}_t)$ of

$$\begin{cases} U' &= V \\ V' &= f(U) \circ \dot{W} + g(U, V) \\ (U_0, V_0) &= Z_0, \end{cases} \quad (5.14)$$

and

$$\begin{cases} \hat{U}' &= \hat{V} \\ \hat{V}' &= \hat{f}(\hat{U}) \circ \dot{W} + \hat{g}(\hat{U}, \hat{V}) \\ (\hat{U}_0, \hat{V}_0) &= \hat{Z}_0. \end{cases} \quad (5.15)$$

We will need two lemmas.

Lemma 5.2 *Suppose f and g are Lipschitz continuous, with Lipschitz constants dominated by L , and suppose \hat{f} and \hat{g} are affine. Put $F(x) = f(x) - \hat{f}(x)$ and $G(x, y) = g(x, y) - \hat{g}(x, y)$. Suppose Z_0 and \hat{Z}_0 are \mathcal{F}_0 -measurable. Then for each $T > 0$ there exists a constant $c = c(T, L)$ such that for $0 \leq t \leq T$,*

$$E \left\{ |Z_t - \hat{Z}_t|^2 \right\} \leq 3E \left\{ |Z_0 - \hat{Z}_0|^2 \right\} e^{ct} + 6 \int_0^t e^{c(t-s)} E \left\{ F^2(\hat{U}_s) + tG^2(\hat{Z}_s) \right\} ds. \quad (5.16)$$

PROOF. From (5.14) and the Schwartz inequality,

$$\begin{aligned} E \left\{ (U_t - \hat{U}_t)^2 \right\} &\leq 2E \left\{ (U_0 - \hat{U}_0)^2 + 2t \int_0^t (V_s - \hat{V}_s)^2 ds \right\} \\ &\leq 2E \left\{ (U_0 - \hat{U}_0)^2 \right\} + 2t \int_0^t E \left\{ |Z_s - \hat{Z}_s|^2 \right\} ds. \end{aligned}$$

Similarly

$$E \left\{ (V_t - \hat{V}_t)^2 \right\} \leq E \left\{ \left(V_0 - \hat{V}_0 + \int_0^t (f(U_s) - \hat{f}(\hat{U}_s)) \circ dW_s + \int_0^t (g(Z_s) - \hat{g}(\hat{Z}_s)) ds \right)^2 \right\}.$$

Now the initial conditions are adapted, hence so is the solution, so the stochastic integral is an Ito integral. We compute its expected square as usual to see it is

$$\leq 3E\{(V_0 - \hat{V}_0)^2\} + 3 \int_0^t E\{(f(U_s) - \hat{f}(\hat{U}_s))^2\} ds + 3t \int_0^t E\{(g(Z_s) - \hat{g}(\hat{Z}_s))^2\} ds.$$

Write $|f(U) - \hat{f}(\hat{U})| = |F(\hat{U})| + |f(U) - f(\hat{U})| \leq |F(\hat{U})| + L|Z - \hat{Z}|$, this last since f is Lipschitz (L) and $|U - \hat{U}| \leq |Z - \hat{Z}|$. Similarly, $|g(Z) - \hat{g}(\hat{Z})| \leq |G(\hat{Z})| + L|Z - \hat{Z}|$. Thus the above is

$$\begin{aligned} \leq 3E\{(V_0 - \hat{V}_0)^2\} + 6 \int_0^t E\{F(\hat{U}_s)^2 + tG(\hat{Z}_s)^2\} ds \\ + 6L^2 \int_0^t E\{(Z_s - \hat{Z}_s)^2\} ds + 6L^2t \int_0^t E\{(Z_s - \hat{Z}_s)^2\} ds. \end{aligned}$$

Note that $E\{|Z_t - \hat{Z}_t|^2\} = E\{(U_t - \hat{U}_t)^2\} + E\{(V_t - \hat{V}_t)^2\}$. Thus, for $t \leq T$, if we take $c(T, L) = 2T + 6L^2 + 6TL^2$, we get

$$\begin{aligned} E\{(Z_t - \hat{Z}_t)^2\} \leq 3E\{(Z_0 - \hat{Z}_0)^2\} + 6 \int_0^t E\{F(\hat{U}_s)^2 + tG(\hat{Z}_s)^2\} ds \\ + c(T, L) \int_0^t E\{|Z_s - \hat{Z}_s|^2\} ds. \quad (5.17) \end{aligned}$$

This implies (5.16) by Gronwall's inequality. \square

Lemma 5.3 *Suppose that f and g are differentiable with bounded Lipschitz continuous derivatives. Let $T > 0$ and $M > 0$. Then there exists constants C and D , depending on T , M , and the functions f , g , \hat{f} , and \hat{g} , but not on x or y , such that for $0 \leq t \leq T$*

$$E\left\{ \sup_{|x| \leq M} (Z_t(x, y) - \hat{Z}_t(x, y))^2 \right\} \leq C(1+t) + D \int_0^t \left(F(y\hat{U}_s(0, 1))^2 + G(y\hat{Z}_s(0, 1))^2 \right) ds.$$

PROOF. $|Z_t(x, y) - \hat{Z}_t(x, y)| \leq |Z_t(x, y) - Z_t(0, y)| + |Z_t(0, y) - \hat{Z}_t(0, y)| + |\hat{Z}_t(0, y) - \hat{Z}_t(x, y)|$. Since \hat{f} and \hat{g} are linear, not just affine, (4.13) tells us that $\hat{Z}_t(x, y) = x\hat{Z}_t(1, 0) + y\hat{Z}_t(0, 1)$, so $\hat{Z}_t(x, y) - \hat{Z}_t(0, y) = x\hat{Z}_t(1, 0)$ and

$$E \left\{ \sup_{|x| \leq M} (\hat{Z}_t(x, y) - \hat{Z}_t(0, y))^2 \right\} \leq M^2 E \{ \hat{Z}_t(1, 0)^2 \} \equiv A(t). \quad (5.18)$$

Now $\hat{Z}_t(0, y) = y \hat{Z}_t(0, 1)$, so by Lemma 5.2, there exists a constant c such that

$$E \left\{ (Z_t(0, y) - \hat{Z}_t(0, y))^2 \right\} \leq 6 \int_0^t e^{c(t-s)} E \left\{ F(y \hat{U}_s(0, 1))^2 + tG(y \hat{Z}_s(0, 1))^2 \right\} ds. \quad (5.19)$$

This leaves the term $E \left\{ \sup_{|x| \leq M} (Z_t(x, y) - Z_t(0, y))^2 \right\}$ to estimate. Under the hypotheses of the lemma, $Z_t(x, y)$ is differentiable with respect to its initial conditions, and if we put $\dot{U}_t = \frac{\partial U_t}{\partial x}(x, y)$ and $\dot{V}_t = \frac{\partial V_t}{\partial x}(x, y)$, then by Kunita's theorem (Thm 3.2), \dot{U} and \dot{V} satisfy (3.11), which is, in this case:

$$\begin{cases} d\dot{U}(x, y) &= \dot{V}(x, y) dt \\ d\dot{V}(x, y) &= f'(U_t(x, y)) \dot{U}_t(x, y) \circ dW_t + g_x(U_t(x, y), V_t(x, y)) \dot{U}_t(x, y) dt \\ &\quad + g_y(U_t(x, y), V_t(x, y)) \dot{V}_t(x, y) dt \\ \dot{U}_0(x, y) &= 1, \\ \dot{V}_0(x, y) &= 0. \end{cases} \quad (5.20)$$

Notice that the coefficients of (5.20) are bounded (because the derivatives of f and g are), and the initial conditions are independent of x and y . Thus it is easy to see that there is a function $D(t)$ which is independent of x and y such that

$$E \left\{ \dot{U}_t^2(x, y) + \dot{V}_t^2(x, y) \right\} \leq D(t).$$

Thus, if $|x| \leq M$

$$\begin{aligned} |Z_t(x, y) - Z_t(0, y)|^2 &= \left(\int_0^x \dot{U}_t(\tau, y) d\tau \right)^2 + \left(\int_0^x \dot{V}_t(\tau, y) d\tau \right)^2 \\ &\leq |x| \int_0^x \dot{U}_t^2(\tau, y) + \dot{V}_t^2(\tau, y) d\tau \\ &\leq M \int_{-M}^M \dot{U}_t^2(\tau, y) + \dot{V}_t^2(\tau, y) d\tau. \\ &\leq 2M^2 D(t), \end{aligned} \quad (5.21)$$

and the conclusion follows by combining (5.18), (5.19), and (5.21). \square

5.1 Proof of Theorem 2.3

By Lemma 5.1 it is enough to show that the projection of L_1 is \mathbb{R} , or equivalently, that $\inf_y U_1(\xi, y) = -\infty$ and $\sup_y U_1(\xi, y) = \infty$. We will do this by comparing the solution to the solution \hat{U} of the linear equation, i.e. of (1.3) with f and g replaced by $\hat{f}(x) = ax$, and

$\hat{g}(x, y) = cx + dy$. It is enough to prove this for the case where ξ is bounded, say $|\xi| \leq M$. Consider

$$\begin{aligned} y^{-2}E\left\{\left(U_1(\xi, y) - \hat{U}_1(\xi, y)\right)^2\right\} &\leq y^{-2}E\left\{\sup_{|x|\leq M}\left(U_1(x, y) - \hat{U}_1(x, y)\right)^2\right\} \\ &\leq y^{-2}E\left\{\sup_{|x|\leq M}\left(Z_1(x, y) - \hat{Z}_1(x, y)\right)^2\right\} \\ &\leq c\frac{1+t}{y^2} + \frac{b}{y^2}\int_0^1 E\left\{F(y\hat{U}_s(0, 1))^2 + G(y\hat{Z}_s(0, 1))^2\right\} ds \end{aligned}$$

by Lemma 5.3, where $F(x) = f(x) - \hat{f}(x)$ and $G(x, y) = g(x, y) - \hat{g}(x, y)$. Now $\hat{U}_s(0, 1) \neq 0$, so $|y\hat{U}_s(0, 1)| \rightarrow \infty$ as $|y| \rightarrow \infty$. But

$$\frac{F(y\hat{U}_s(0, 1))}{y} = \hat{U}_s(0, 1) \frac{F(y\hat{U}_s(0, 1))}{y\hat{U}_s(0, 1)}$$

which converges to zero boundedly as $|y| \rightarrow \infty$ by hypothesis. The same is true of the second term. Thus the expectation tends to zero as $|y| \rightarrow \infty$, so we conclude that $y^{-1}(U_1(\xi, y) - \hat{U}_1(\xi, y)) \rightarrow 0$ in L^2 and hence in probability as $|y| \rightarrow \infty$. But now $y^{-1}\hat{U}_1(\xi, y) \rightarrow \hat{U}_1(0, 1)$ as $|y| \rightarrow \infty$, hence so does $y^{-1}U_1(\xi, y)$, in probability, at least. It follows easily that $\inf_y U_1(\xi, y) = -\infty$ and $\sup_y U_1(\xi, y) = \infty$, as claimed. \square

5.2 Proof of Theorem 2.5

Suppose now that the initial condition on U is adapted, i.e. that $\xi \in \mathcal{F}_0$. In this case, if we consider the initial value problem, we can condition on the value of ξ ; given $\xi = x$, $U_t(\xi, y) \equiv U_t(x, y)$, and we can use Lemma 5.2, which has weaker hypotheses than Lemma 5.3. We must show that, conditional on ξ , $\inf_{y \in \mathbb{R}} U_1(\xi, y) = -\infty$ and $\sup_{y \in \mathbb{R}} U_1(\xi, y) = \infty$ with probability one.

By Lemma 5.2,

$$E\left\{\left(\frac{Z_1(\xi, y) - \hat{Z}_1(\xi, y)}{y}\right)^2 \mid \xi\right\} \leq 6 \int_0^1 e^{c(t-s)} E\left\{\frac{F(\hat{U}_s(\xi, y))^2 + G(\hat{Z}_s)^2}{y^2} \mid \xi\right\}$$

But now, given that $\xi = x$, $\hat{U}_s(\xi, y) = x\hat{U}_s(1, 0) + y\hat{U}_s(0, 1)$ and $\hat{Z}_s(\xi, y) = x\hat{Z}_s(1, 0) + y\hat{Z}_s(0, 1)$, so that $F^2(\hat{U}_s(x, y))/y^2 = (F^2(\hat{U}_s(x, y))/\hat{U}_s(x, y))(\hat{U}_s(x, y)^2/y^2)$. As $|y| \rightarrow \infty$, $|\hat{U}_s(x, y)| \rightarrow \infty$ and $F^2(\hat{U}_s(x, y))/\hat{U}_s(x, y)$ tends boundedly to zero by hypothesis. The same is true of $G^2(\hat{Z}_s(x, y))/\hat{Z}_s(x, y)$, so $Z_1(\xi, y) - \hat{Z}_1(\xi, y)/y$ tends to zero in L^2 and hence in probability as $|y| \rightarrow \infty$. Since $\hat{Z}_1(\xi, y)/y \rightarrow \hat{U}_1(0, 1) \neq 0$, we conclude that as $y \rightarrow \pm\infty$, that $Z_1(\xi, y)$ and $\hat{Z}_1(\xi, y)$ have the same (infinite) limits. This implies that with probability one, $\inf_{y \in \mathbb{R}} U_1(\xi, y) = -\infty$ and $\sup_{y \in \mathbb{R}} U_1(\xi, y) = \infty$, as claimed. \square

6 The Green's function

The Green's function for the two-point boundary-value problem on $[0, T]$ is

$$G(s, t) = \begin{cases} T^{-1}s(T-t) & \text{if } 0 \leq s \leq t \leq T \\ T^{-1}t(T-s) & \text{if } 0 \leq t \leq s \leq T. \end{cases}$$

Let us also define

$$H(s, t) = \frac{\partial}{\partial s} G(s, t) = \begin{cases} T^{-1}(T-t) & \text{if } 0 \leq s \leq t \leq T, \\ -T^{-1}t & \text{if } 0 \leq t \leq s \leq T. \end{cases}$$

For an integrable (possibly random) function h on $[0, T]$, define $Gh(t) \equiv \int_0^T G(s, t)h(s) ds$ and $G(h\dot{W})(t) \equiv \int_0^T G(s, t)h(s) \circ dW_s$. Note that $G \geq 0$ and

$$\int_0^T G(s, t) ds = \frac{t(T-t)}{2} \leq \frac{T^2}{8}; \quad (6.22)$$

$$\int_0^T |H(s, t)| ds = \frac{2t(T-t)}{T} \leq \frac{T}{2}. \quad (6.23)$$

Lemma 6.1 *Suppose $h \in L^1(0, T)$ and that $\{Z_t, 0 \leq t \leq T\}$ is a process which is a.s. absolutely continuous in t . Then*

(i) $Gh(t)$ is a differentiable function of t and

$$\frac{d}{dt} Gh(t) = \int_0^t h(s) ds - T^{-1} \int_0^T \int_0^s h(u) du ds;$$

(ii) $G(Z\dot{W})(t)$ is a differentiable function of t and

$$\begin{aligned} \frac{d}{dt} G(Z\dot{W})(t) &= \int_0^t Z_s \circ dW_s - T^{-1} \int_0^T \int_0^s Z_u \circ dW_u ds \\ &= Z_t W_t - \int_0^t W_s Z'_s ds - T^{-1} \int_0^T \int_0^s Z_u \circ dW_u ds. \end{aligned}$$

PROOF. The proof is nearly the same in both cases: we simply integrate by parts. In (i), set $J(s) = \int_0^s h(u) du$ and integrate by parts: $Gh(t) = G(T, t)J(T) - G(0, t)J(0) - \int_0^T H(s, t)J(s) ds$. The first two terms vanish, and we can write the last integral out explicitly and differentiate it to see (i). In (ii), set $K(s) = \int_0^s Z_u \circ dW_u$, and integrate the Stratonovich integral by parts:

$$G(Z\dot{W})(t) = G(T, t)K(T) - G(0, t)K(0) - \int_0^t H(s, t)K(s) ds. \quad (6.24)$$

The first two terms on the right-hand side vanish, and we can again write out the integral explicitly and differentiate it. \square

It is now clear (and quite well-known) that $u(t) \equiv Gh(t)$ satisfies

$$\begin{cases} u'' = h(t) & \text{if } 0 < t < T \\ u(0) = u(T) = 0. \end{cases}$$

This leads to another statement of the stochastic problem.

Proposition 6.2 *Let f and g be Lipschitz continuous functions on \mathbb{R} and \mathbb{R}^2 respectively, and let $\{U(t), 0 \leq t \leq T\}$ be a process which is a.s. absolutely continuous as a function of t . Then (1.3) is equivalent to*

$$U(t) = T^{-1}((T-t)\xi + t\eta) + G(f(U, \dot{W}))(t) + G(g(U, U'))(t), \quad 0 \leq t \leq T, \quad (6.25)$$

in the sense that if Λ is any set of positive probability, then U satisfies (1.3) a.e. on Λ if and only if it satisfies (6.25) a.e. on Λ .

Remark 6.3 *Equation (6.25) is sometimes called the mild form of (1.3).*

PROOF. Suppose U is absolutely continuous and satisfies (1.3) on a set Λ . Then $U'_t = V_t$ exists for a.e. t , so on Λ

$$U(t) = \xi + \int_0^t \left(U'_0 + \int_0^s f(U_u) \circ dW_u + \int_0^s g(U_u, U'_u) du \right) ds \quad (6.26)$$

$$= \xi + U'_0 t + \int_0^t (t-s)f(U_s) \circ dW_s + \int_0^t (t-s)g(U_s, U'_s) ds. \quad (6.27)$$

Set $t = T$. Then $U_T = \eta$ so

$$U'_0 = T^{-1}(\eta - \xi) - T^{-1} \int_0^T (T-s)f(U_s) \circ dW_s - T^{-1} \int_0^T (T-s)g(U_s, U'_s) ds. \quad (6.28)$$

Plug this in (6.27) and combine terms to see that

$$U(t) = T^{-1}((T-t)\xi + t\eta) + \int_0^t G(s, t)f(U_s) \circ dW_s + \int_0^t G(s, t)g(U_s, U'_s) ds, \quad (6.29)$$

which is (6.25). Conversely, if U is absolutely continuous and satisfies (6.25) a.e. on Λ , then Lemma 6.1 shows it must also satisfy (1.3) a.e. on Λ . \square

6.1 Proof of Theorem 2.7

Note first that it is enough to prove this for the case where ξ and η are bounded, say $|\xi| + |\eta| \leq M$, for some $M > 0$. Since f' is bounded and Lipschitz, we let L_f be a common bound for $|f'|$ and the Lipschitz constant for f' . (So L_f is a common Lipschitz constant for both f and f' .)

Let $Z_t = (U_t, V_t)$, set

$$U_t^0 = T^{-1}((T-t)\xi + t\eta), \quad V_t^0 = T^{-1}(\eta - \xi),$$

and define

$$U_t^{n+1} = U_t^0 + G \left(f(U^n) \dot{W} + g(Z^n) \right).$$

Integrate the Stratonovich integral by parts to see that

$$\begin{aligned} U_t^{n+1} &= U_t^0 - \int_0^T H(s, t) f(U_s^n) W_s ds \\ &\quad + \int_0^T G(s, t) (g(Z_s^n) - W_s f'(U_s^n) V_s^n) ds. \end{aligned} \quad (6.30)$$

Now $V_t^{n+1} = \frac{\partial}{\partial t} U_t^{n+1}$, so by Lemma 6.1

$$\begin{aligned} V_t^{n+1} &= V_t^0 + W_t f(U_t^n) - T^{-1} \int_0^T f(U_s^n) W_s ds + \int_0^T (g(Z_s^n) - W_s f'(U_s^n) V_s^n) ds \\ &\quad + T^{-1} \int_0^T \int_0^s (g(Z_u^n) + W_u f'(U_u^n) V_u^n) du ds, \end{aligned} \quad (6.31)$$

while on the other hand

$$U_t^{n+1} = \xi + \int_0^t V_s^{n+1} ds. \quad (6.32)$$

Both f and g are Lipschitz so $|f(x)| \leq f_0 + L_f|x|$, $|f'(x)| \leq L_f$, and $|g(z)| \leq g_0 + L_g|z|$. Denote $\|h\| = \sup_{t \in [0, T]} |h(t)|$. Using (6.22) and (6.23), and the fact that $\|U^n\|$ and $\|V^n\|$ are dominated by $\|Z^n\|$, we see

$$\|U^{n+1}\| \leq \|U^0\| + \frac{T}{2} \|W\| f_0 + \frac{T^2}{8} g_0 + \left(\|W\| L_f \left(\frac{T}{2} + L_g \frac{T^2}{8} \right) + \frac{T^2}{8} \right) \|Z^n\|,$$

$$\|V^{n+1}\| \leq \|V^0\| + 2\|W\| f_0 + \frac{3}{2} T g_0 + \left(\|W\| L_f \left(2 + \frac{3T}{2} \right) + \frac{3T}{2} L_g \right) \|Z^n\|.$$

Now $\|Z\| \leq \|U\| + \|V\|$ so, adding,

$$\begin{aligned} \|Z^{n+1}\| &\leq \|U^0\| + \|V^0\| + \left(2 + \frac{T}{2} \right) \|W\| f_0 + \left(\frac{3T}{2} + \frac{T^2}{8} \right) g_0 \\ &\quad + \left(\|W\| L_f \left(2 + 2T + \frac{T^2}{8} \right) + \left(\frac{3T}{2} + \frac{T^2}{8} \right) L_g \right) \|Z^n\|. \end{aligned}$$

By hypothesis, $\left(\frac{3T}{2} + \frac{T^2}{8} \right) L_g < 1$, so we can choose $\epsilon > 0$ and $\delta_1 > 0$ small enough so that

$$\delta_1 L_f \left(2 + 2T + \frac{T^2}{8} \right) + L_g \left(\frac{3T}{2} + \frac{T^2}{8} \right) < 1 - 2\epsilon. \quad (6.33)$$

If $\|W(\omega)\| \leq \delta_1$, then

$$\|Z^{n+1}\| \leq \|U^0\| + \|V^0\| + \left(2 + \frac{T}{2} \right) |f_0| \delta_1 + \left(\frac{3T}{2} + \frac{T^2}{8} \right) |g_0| + (1 - 2\epsilon) \|Z^n\|.$$

If necessary, replace M by $\max \left(M, \left(\frac{3T}{2} + \frac{T^2}{8} \right) \right)$. Now $|U^0|$ and $|V^0|$ are both bounded by M , so if we choose δ_1 small enough that $(2 + T/2)|f_0|\delta_1 \leq M$, we have $\|Z^{n+1}\| \leq 4M + (1 - 2\epsilon)\|Z^n\|$. It follows that for all n ,

$$\|Z^n\| \leq 4M \|Z^0\| / 2\epsilon \leq 4M^2 / \epsilon. \quad (6.34)$$

Now consider the increments, using the usual notation: $\Delta Z^n \equiv Z^n - Z^{n-1}$.

$$\begin{aligned} \Delta U_t^{n+1} &= - \int_0^T H(s, t) W_s (f(U_s^n) - f(U_s^{n-1})) ds \\ &\quad - \int_0^T G(s, t) W_s (f'(U_s^n) V_s^n - f'(U_s^{n-1}) V_s^{n-1}) ds \\ &\quad + \int_0^T G(s, t) (g(Z_s^n) - g(Z_s^{n-1})) ds. \end{aligned} \quad (6.35)$$

Similarly

$$\begin{aligned} \Delta V_t^{n+1} &= W_t (f(U_t^n) - f(U_t^{n-1})) - \frac{1}{T} \int_0^T W_s (f(U_s^n) - f(U_s^{n-1})) ds \\ &\quad + \int_0^t W_s (f'(U_s^n) V_s^n - f'(U_s^{n-1}) V_s^{n-1}) ds \\ &\quad + \int_0^t (g(Z_s^n) - g(Z_s^{n-1})) ds + \frac{1}{T} \int_0^T \int_0^s (g(Z_u^n) - g(Z_u^{n-1})) du ds \\ &\quad - \frac{1}{T} \int_0^T \int_0^s W_u (f'(U_u^n) V_u^n - f'(U_u^{n-1}) V_u^{n-1}) du ds \end{aligned} \quad (6.36)$$

From the Lipschitz conditions, we see $\|\Delta f(U^n)\| \leq L_f \|\Delta U^n\| \leq L_f \|\Delta Z^n\|$, $\|\Delta g(Z^n)\| \leq L_g \|\Delta Z^n\|$, and $\|\Delta(f'(U)V)\| \leq L_f \|V^n\| \|\Delta U^n\| + L_f \|\Delta V^n\| \leq L_f (\|Z^n\| + 1) \|\Delta Z^n\|$. Thus

$$\begin{aligned} \|\Delta U^{n+1}\| &\leq \left(\|W\| L_f \left(\frac{T}{2} + \frac{T^2}{8} (\|Z^n\| + 1) \right) + \frac{T^2}{8} L_g \right) \|\Delta Z^n\|; \\ \|\Delta V^{n+1}\| &\leq \left(\|W\| L_f \left(2 + \frac{3}{2} T (\|Z^n\| + 1) \right) + \frac{3T}{2} L_g \right) \|\Delta Z^n\|. \end{aligned}$$

Adding these two:

$$\|\Delta Z^{n+1}\| \leq \left(\left(\frac{3T}{2} + \frac{T^2}{8} \right) L_g + \|W\| L_f \left(2 + \frac{T}{2} + \left(\frac{3T}{2} + \frac{T^2}{8} \right) (\|Z^n\| + 1) \right) \right) \|\Delta Z^n\|. \quad (6.37)$$

By (6.33), if $\|W\| \leq \delta_1$, this is

$$\leq \left(1 - 2\epsilon + L_f \delta_1 \left(\left(2 + \frac{T}{2} \right) + \left(\frac{3T}{2} + \frac{T^2}{8} \right) \left(\frac{3M}{\epsilon} + 1 \right) \right) \right) \|\Delta Z^n\|$$

We can choose $\delta_2 > 0$ small enough so that $\delta_2 \leq \delta_1$ and

$$L_f \delta_2 \left(2 + \frac{T}{2} + \left(\frac{3T}{2} + \frac{T^2}{8} \right) \left(\frac{3M}{\epsilon} + 1 \right) \right) < \epsilon. \quad (6.38)$$

Thus

$$\|W\| < \delta_2 \implies \|\Delta Z^{n+1}\| \leq (1 - \epsilon) \|\Delta Z^n\|, \quad n = 1, 2, \dots \quad (6.39)$$

Let $\Omega_0 = \{\omega : \|W(\omega)\| < \delta_2\}$. If $\omega \in \Omega_0$, the sequence $(Z_t^n(\omega))$ converges uniformly for $0 \leq t \leq T$ to a limit $Z_t(\omega)$, and we can go to the limit on both sides of (6.30), (6.31), and (6.32) to see that Z satisfies (6.25) and hence (1.3) on Ω_0 .

To see uniqueness, suppose that $Z^i = (U^i, V^i)$, $i = 1, 2$ are solutions. Then

$$\begin{aligned} U_t^2 - U_t^1 &= - \int_0^T H(s, t) W_s (f(U_s^2) - f(U_s^1)) ds + \int_0^T G(s, t) (g(Z_s^2) - g(Z_s^1)) ds \\ &\quad - \int_0^T G(s, t) W_s (f'(U_s^2) V_s^2 - f'(U_s^1) V_s^1) ds. \end{aligned}$$

This has the same right hand side as (6.35) with $n = 2$. $V^2 - V^1$ satisfies a similar equation, and if we repeat the calculations above, we see that on Ω_0 , $\|Z^2 - Z^1\| \leq (1 - \epsilon) \|Z^2 - Z^1\|$, hence $Z^2 \equiv Z^1$.

Finally, to see what happens for small T , let $T \rightarrow 0$ and note that for any Lipschitz g , we can find $T_0 > 0$ such that for $T \leq T_0$, $\left(\frac{3T}{2} + \frac{T^2}{8} \right) L_g < \frac{1}{4}$. We then choose $\delta_1 > 0$ small enough so that $T < T_0 \implies \delta_1 L_f \left(2 + 2T + \frac{T^2}{8} \right) \leq \frac{1}{4}$.

Thus by (6.34), $T \leq T_0$ and $\|W\| \leq \delta_1$ imply that for all n ,

$$\|Z^{n+1}\| \leq 2 \left(2M + \left(2 + \frac{T^0}{2} \right) f_0 \delta_1 + \left(\frac{3T_0}{2} + \frac{T_0^2}{8} \right) g_0 \right)$$

hence by (6.37)

$$\begin{aligned} \|\Delta Z_{n+1}\| &\leq \left(\frac{1}{2} + 2 \left(\frac{3T_0}{2} + \frac{T_0^2}{8} \right) \right) L_f \|W\| \left(2M + \left(2 + \frac{T_0}{2} \right) f_0 \delta_1 \right. \\ &\quad \left. + \left(\frac{3T_0}{2} + \frac{T_0^2}{8} \right) g_0 \right) \|\Delta Z^n\|. \end{aligned}$$

Choose $\delta_2 > 0$ so that

$$2\delta_2 L_f \left(2M + \left(2 + \frac{T_0}{2} \right) f_0 \delta_1 + \left(\frac{3T_0}{2} + \frac{t_0^2}{8} \right) g_0 \right) < \frac{1}{4}.$$

Then $T \leq T_0$ and $\|W\| \leq \delta_2$ imply that $\|\Delta Z^{n+1}\| \leq \frac{3}{4}\|\Delta Z^n\|$. As we saw, this implies existence and uniqueness, so that if we set $\Omega_{0T} = \{\omega : \sup_{0 \leq t \leq T} |W_t(\omega)| < \delta_2\}$, the two point boundary value problem will have a unique local solution on Ω_{0T} . But $P\{\Omega_{0T}\} = P\{\sup_{0 \leq t \leq T} |W_t| < \delta_2\}$, which tends to one as $T \rightarrow 0$. \square

7 Counterexamples

One way of constructing counterexamples is to work from the deterministic case. With a few regularity conditions on the coefficients, the solutions are continuous as a function of the integrators, and one can use this to construct stochastic examples from deterministic ones.

For any process $\{X_t, t \geq 0\}$, let $X_t^* = \sup_{0 \leq s \leq t} |X_s|$. Let $\{N_t, t \geq 0\}$ be a continuous semimartingale with $N_0 = 0$, and let $Z_t = (U_t, V_t)$, where U and V are solutions of the system

$$\begin{cases} U_t &= U_0 + \int_0^t V_s ds \\ V_t &= V_0 + \int_0^t f(U_s) \circ dN_s + \int_0^t g(U_s, V_s) ds. \end{cases} \quad (7.40)$$

Lemma 7.1 *Let f and g be Lipschitz, and suppose that f is also differentiable with a Lipschitz continuous derivative. Let $T > 0$. Then there exist constants $C = C(T)$ and $D = D(T)$ such that*

$$Z_T^* \leq C(1 + |Z_0|)e^{DN_T^*}. \quad (7.41)$$

PROOF. We can suppose that there exists L such that $|f(x) - f(x')| \leq L|x - x'|$, $|f(x)| \leq L(1 + |x|)$, $|g(z) - g(z')| \leq L|z - z'|$, and $|g(z)| \leq L(1 + |z|)$. Let $t \leq T$ and integrate (7.40) by parts:

$$V_t = V_0 + N_t f(U_t) - \int_0^t N_s f'(U_s) V_s ds + \int_0^t g(Z_s) ds.$$

Now $|f(U_t)| \leq L(1 + |U_t|) \leq L(1 + |U_0| + \int_0^t |V_s| ds)$, so we can write

$$|V_t| \leq |V_0| + L|N_t| \left(1 + |U_0| + \int_0^t |V_s| ds \right) + LN_t^* \int_0^t |V_s| ds + \int_0^t L(1 + |Z_s|) ds$$

Both $|U_s|$ and $|V_s|$ are dominated by $|Z_s| \leq Z_s^*$ and $Z_s^* \leq U_s^* + V_s^*$, so that

$$Z_s^* \leq |U_0| + \int_0^t V_s^* ds + LN_T^* (1 + |U_0|) + LT + (1 + L + 2N_T^*) \int_0^t Z_s^* ds.$$

Gronwall's inequality then yields

$$Z_t^* \leq \left(|Z_0|(\sqrt{2} + LN_T^*) + L(T + N_T^*) \right) e^{(1+L+2LN_T^*)t}.$$

If we use the (very coarse) inequality $LN_T^* \leq e^{LN_T^*}$, we get (7.41). \square

Suppose that M_t is another continuous semimartingale with $M_0 = 0$, and consider the system

$$\begin{cases} U_t &= U_0 + \int_0^t V_s ds \\ V_t &= V_0 + \int_0^t f(U_s) \circ dM_s + \int_0^t g(U_s, V_s) ds \end{cases} \quad (7.42)$$

Proposition 7.2 *Let f and g be Lipschitz, and suppose that f is also differentiable with a Lipschitz continuous derivative. Let $T > 0$. Let $Z^M = (U^M, V^M)$ and $Z^N = (U^N, V^N)$ be solutions of (7.42) and (7.40) respectively, such that $Z_0^M = Z_0^N$. Then if $\bar{M} = \max(N_T^*, M_T^*)$, there exists a function $C(T, M, |Z_0|)$, increasing in each of the variables, such that*

$$(\Delta Z)_T^* \leq C(T, \bar{M}, |Z_0|)(\Delta N)_T^*, \quad (7.43)$$

where $\Delta Z_t = Z_t^N - Z_t^M$ and $\Delta N_t = N_t - M_t$.

PROOF. Let $t \leq T$. Then

$$\begin{aligned} V_t^N - V_t^M &= N_t f(U_t^N) - M_t f(U_t^M) - \int_0^t (N_s f'(U_s^N) V_s^N - M_s f'(U_s^M) V_s^M) ds \\ &\quad + \int_0^t (g(Z_s^N) - g(Z_s^M)) ds, \end{aligned}$$

and

$$U_t^N - U_t^M = \int_0^t (V_s^N - V_s^M) ds.$$

Let L dominate the Lipschitz constants of f , f' , and g , so that $|f'(x)| \leq L$ for all x . Write $\Delta(Nf) = N\Delta f + f\Delta N$, $\Delta(Nf'V) = Nf'\Delta V + NV^M\Delta f' + V^M f'\Delta N$ and note that $|f(U_t^N) - f(U_t^M)| \leq L|\Delta U_t| \leq L \int_0^t |\Delta V_s| ds \leq L \int_0^t \Delta Z_s^* ds$, to see that

$$\begin{aligned} |\Delta V_t| &\leq N_t^* L |\Delta U_t| + L(1 + |U_t^M|) |\Delta N_t| + L \int_0^t (|N_s| |\Delta V_s| + |N_s| |V_s^M| |\Delta U_s| \\ &\quad + V_s^M |\Delta N_s|) ds + \int_0^t L |\Delta Z_s| ds, \end{aligned}$$

and

$$\Delta U_s = \int_0^t \Delta V_s ds.$$

Collect terms, use the obvious bounds $|\Delta U_s| \leq (\Delta Z_s)^* \leq (\Delta U)^* + (\Delta V)^*$, and $N_t^* \leq N_T^*$ for $t \leq T$:

$$(\Delta Z)_t^* \leq L(1 + (1 + T)Z_T^{M*}) (\Delta N)_T^* + L(Z_T^{M*} + 2N_T^*) \int_0^t (\Delta Z)_s^* ds,$$

hence, by Gronwall's inequality,

$$(\Delta Z)_t^* \leq L(1 + (T + 1)Z_T^{M*}) (\Delta N)_T^* e^{L(Z_T^{M*} + 2N_T^*)t},$$

for $t \leq T$. Now use the bound on Z_T^{M*} given in Lemma 7.1 to finish the proof. \square

Remark 7.3 It is easy to modify the above in case $Z_0^M \neq Z_0^N$. Just notice that $|\Delta Z_t| \leq |\Delta Z_0| + |\Delta U_t - \Delta U_0| + |\Delta V_t - \Delta V_0|$. Then the calculations above show that $|\Delta Z_t^*| \leq |\Delta Z_0| + C_1 |\Delta N_t| + C_2 \int_0^t |\Delta Z_s^*| ds$ for some C_1 and C_2 which depend on T , Z_0 and the bounds on N and M . This leads to

$$|\Delta Z_t^*| \leq (|\Delta Z_0| + C_1 |\Delta N_t^*|) e^{C_2 t}.$$

We will usually use this when $M_t = W_t$ and $N_t = \int_0^t h(s) ds$ for some deterministic function h , in which case we get

Corollary 7.4 *Let $f \in C^1(\mathbb{R})$ and suppose f' is Lipschitz. Let $h \in L_{loc}^1$ on $[0, \infty)$ and let $\hat{U}_t(x, y)$ and $U_t(x, y)$ be the respective solutions of*

$$\begin{cases} \hat{U}_t''(x, y) &= f(\hat{U}_t(x, y)) h(t) \\ \hat{U}_0(x, y) &= x \\ \hat{U}'_0(x, y) &= y. \end{cases} \quad (7.44)$$

and

$$\begin{cases} U_t''(x, y) &= f(U_t(x, y)) \circ \dot{W}_t \\ U_0(x, y) &= x \\ U'_0(x, y) &= y. \end{cases} \quad (7.45)$$

Then for each $R > 0$, $T > 0$ and $\varepsilon > 0$ there exists $\Omega_0 \subset \Omega$ with $P\{\Omega_0\} > 0$ such that for $\omega \in \Omega_0$

$$\sup_{|x|, |y| \leq R} \sup_{0 \leq s \leq T} |U_s(x, y; \omega) - \hat{U}_s(x, y; \omega)| < \varepsilon. \quad (7.46)$$

PROOF. By Proposition 7.2. Ω_0 contains the set of ω for which $\sup_{t \leq T} |W_t(\omega) - \int_0^t h(s) ds| < \varepsilon$ for a sufficiently small $\varepsilon > 0$. \square

Let us look at a simple example in which both existence and uniqueness can fail. Let $g(x, y) \equiv 0$ and set

$$f(x) = \begin{cases} |x| & \text{if } x < 0 \\ 0 & \text{if } x \geq 0. \end{cases}$$

Proposition 7.5 *There exists a set Ω_0 with $P\{\Omega_0\} > 0$ such that on Ω_0 , the problem*

$$\begin{cases} U'' = f(U) \circ \dot{W} \\ U(0) = 0 \\ U(1) = x \end{cases} \quad (7.47)$$

has no solution if $x < 0$ and two solutions if $x > 0$.

PROOF. Consider the solution $U_t(y)$ of the initial value problem

$$\begin{cases} U'' = f(U) \circ \dot{W} \\ U(0) = 0 \\ U'(0) = y \end{cases} \quad (7.48)$$

Note that $a > 0 \implies f(ax) = af(x)$, so that if $Z_t(y) = (U_t(y), V_t(y))$, then

$$Z_t(y) = \begin{cases} |y|Z_t(1) & \text{if } y \geq 0 \\ |y|Z_t(-1) & \text{if } y \leq 0 \end{cases} \quad (7.49)$$

It follows that the image L_t of the y -axis under the map $(0, y) \mapsto Z_t(y)$ is a pair of rays, which are the images of the positive and negative y -axes.

The image of the positive y -axis is easy to determine. If $y > 0$, U_t is initially strictly positive, which means that $U_t'' = f(U_t) = 0$, so U_t' is constant, hence $U_t(y) \equiv yt$ for all $t \geq 0$. Thus $Z_t(y) = (yt, y)$, hence the image of the positive y -axis is contained in the positive quadrant for all $t \geq 0$.

Now consider the image of the negative y -axis. Let $\tau = \inf\{t > 0 : U_t = 0\}$. Suppose that $\tau < 1$ and $U_\tau'(1) > 0$. In that case, the Markov property implies that $Z_{\tau+t}(1)$ will be in the first quadrant for all $t > 0$, and hence that the image of the negative y axis at time $t = 1$ is in the first quadrant. Thus, L_1 is in the first quadrant. But this means that (7.48) has no solution if $y < 0$ and two solutions if $y > 0$.

To see that it is possible that $\tau < 1$, consider the deterministic problem

$$\begin{cases} U''(t) = -4\pi^2 U(t) \\ U(0) = 0 \\ U'(0) = -1 \end{cases} \quad (7.50)$$

whose solution is $U(t) = -(1/2\pi) \sin 2\pi t$. At time $\tau = 1/2$, $U = 0$ and $U' = 1$. By Proposition 7.2 the solution of (7.50) will be uniformly close to the solution of

$$\begin{cases} U'' = U \circ \dot{W} \\ U(0) = 0 \\ U'(0) = y \end{cases} \quad (7.51)$$

on the set where $\sup_{0 \leq t \leq 1} |W_t + 4\pi^2 t|$ is small. This set has positive probability, so there is positive probability that the solution of (7.51) will have $\tau < 1$ and $U'_\tau > 0$. But now, by uniqueness, the solution of (7.51) and (7.48) are identical up to time τ , which means that the probability that $\tau < 1$ and $U'_\tau > 0$ is strictly positive for the latter, too, which is what we wanted to show. \square

7.1 Proof of Theorem 2.10

PROOF. The problem has a unique solution if f is affine by Theorem 2.1. On the other hand, we claim that the solution to (2.7) can not be unique if f is not affine. Indeed, if $f(x_0) = 0$, then $U \equiv x_0$ is a global solution of the problem with $\xi = \eta = x_0$.

But by Theorem 2.9, if f is not affine, there exists a set Ω_0 of positive probability and a non-zero function \hat{U} which is a solution on Ω_0 . But then the function \tilde{U} defined by

$$\tilde{U}_t(\omega) = \begin{cases} \hat{U}_t(\omega) & \text{if } \omega \in \Omega_0 \\ 0 & \text{otherwise} \end{cases}$$

is another global solution. \square

8 Non-Uniqueness

In this section, we will use U_t , for the solution of the two-point boundary-value problem

$$\begin{cases} U_t'' = f(U_t) \circ \dot{W}_t \\ U_0 = U_1 = 0, \end{cases} \quad (8.52)$$

we will use $U_t(y)$ for a solution of the initial value problem

$$\begin{cases} U_t''(y) = f(U_t(y)) \circ \dot{W}_t \\ U_0(y) = 0 \\ U_0'(y) = y, \end{cases} \quad (8.53)$$

and we will use $\hat{U}_t(y)$ for a solution of the deterministic initial value problem

$$\begin{cases} \hat{U}_t''(y) = -f(\hat{U}_t(y)) \\ \hat{U}_0(y) = 0 \\ \hat{U}_0'(y) = y. \end{cases} \quad (8.54)$$

Let us consider some properties of the solutions of (8.54). Define

$$T_1(y) \stackrel{\text{def}}{=} \inf\{t > 0 : \hat{U}_t(y) = 0\}.$$

Note that $y > 0 \implies T_1(y) > 0$. Then it is shown in [8] that

Theorem 8.1 (Schaaf) *Suppose that $f \in C^{(1)}$ and that f' is Lipschitz. Then if $f(x) > 0$ in some interval $0 < x < a$, there exists $b > 0$ such that*

- (i) $T_1(y) < \infty$ for $0 \leq y < b$;
- (ii) $y \mapsto T_1(y)$ is differentiable in $(0, b)$;
- (iii) $\hat{U}'_{T_1(y)}(y) = -\hat{U}'_0(y) = -y$, $0 < y < b$.

Lemma 8.2 *Let $\hat{U}(y)$ be a solution of (8.54). Let $\tau > 0$ and define $\tilde{U}_t(y)$ by*

$$\tilde{U}_t(y) = \hat{U}_{\tau t}(y/\tau), \quad t \geq 0.$$

Then $\tilde{U}_t(y)$ is also a solution of (8.54) with f replaced by $\tau^2 f$.

PROOF. Just note that $\tilde{U}_t''(y) = \tau^2 \hat{U}_{\tau t}''(y/\tau) = -\tau^2 f(\hat{U}_{\tau t}(y/\tau)) = -\tau^2 f(\tilde{U}_t(y))$, while $\tilde{U}_0(y) = 0$ and $\tilde{U}'_0(y) = \tau \hat{U}'_0(y/\tau) = y$. \square

Lemma 8.3 *Let f , a , and b be as above. If f is not linear on $(0, a)$, there exists $y \in (0, b)$ such that $T_1'(y) \neq 0$.*

PROOF. For $0 \leq x \leq b$ set

$$g(x) = \frac{1}{\pi} \int_0^x \frac{y T_1(y)}{\sqrt{x^2 - y^2}} dy.$$

By Theorem 4.2.5 of [8], $f(x) = \frac{1}{2} \frac{d}{dx} (g^{-1}(x))^2$ on $(0, z)$. (The theorem is proved under additional hypotheses on f , but this formula holds without them.). If $T_1(\cdot)$ is constant, say $T_1(y) \equiv A$, then $g(x) = A^2 x / 2\pi^2$, so $f(x) = \pi^2 x / A^2$, and f is linear. It follows that if f is not linear, $y \mapsto T_1(y)$ is non-constant on $(0, b)$, and the lemma follows. \square

8.1 Proof of Theorem 2.9

Our strategy is to first find a solution of the deterministic initial value problem which does about what we want, and then approximate it by a stochastic solution.

PROOF. (i) If f is not affine, f' is not constant, and we can find a point x_1 such that f' is not constant in any neighborhood of x_1 and, moreover, $f'(x_1) \neq 0$. We will assume that $x_1 > x_0$, since the case $x_1 < x_0$ is treated almost identically. By replacing f by $-f$ if necessary, we will

assume $f(x_1) > 0$. Let $x_0 = \sup\{x : x < x_1, f(x) = 0\}$. Then $x_0 < x_1$, $f(x_0) = 0$, and $f(x) > 0$ if $x_0 < x < x_1$.

By replacing U_t by $U_t - x_0$ and $f(u)$ by $f(u + x_0)$ we may assume $x_0 = 0$, and $f(0) = 0$. Then by hypothesis, f is not affine on $(0, x_1)$, so that by Lemma 8.3 there exists $b > 0$ and $y \in (0, b)$ such that $T'(y) \neq 0$. We assume $T'(y) > 0$. (The case $T'(y) < 0$ is similar.) As T' is continuous, there is an interval $[c, d] \subset (0, b)$ such that $T'(y) > 0$ for $c \leq y \leq d$. Thus $T_1(c) < T_1(d)$.

The deterministic solution \hat{U}_t satisfies $\hat{U}_t(d) > 0$ if $0 < t < T_1(d)$ while, as $\hat{U}_{T_1(c)}(c) = 0$ and $\hat{U}'_{T_1(c)}(c) = -c < 0$ by Theorem 8.1, we see $\hat{U}_t(c) < 0$ for $T_1(c) < t < T_1(c) + \delta$ for some $\delta > 0$. Thus, if $\tau \in (T_1(c), T_1(d))$ is close enough to $T_1(c)$, there exists $\varepsilon > 0$ such that $\hat{U}_\tau(c) < -\varepsilon < \varepsilon < \hat{U}_\tau(d)$.

Now define $\tilde{U}_t(y) = \hat{U}_{\tau t}(y/\tau)$. By Lemma 8.2, $\tilde{U}(y)$ satisfies $\tilde{U}''(\tau y) = -\tau^2 f(\tilde{U}(\tau y))$, and

$$\tilde{U}_1(\tau c) < -\varepsilon < \varepsilon < \tilde{U}_1(\tau d).$$

We will now approximate \tilde{U} by a solution of the stochastic initial-value problem. Let $\Omega_\delta \subset \Omega = \{\omega : |W_t(\omega) + \tau^2 t| < \delta, 0 \leq t \leq 1\}$. Let $U_t(y)$ be the solution of (8.53). By Corollary 7.4, if δ is small enough, $\omega \in \Omega_\delta \implies |U_t(y) - \tilde{U}_t(y)| < \varepsilon/2, 0 \leq t \leq 1, \tau c \leq y \leq \tau d$. Therefore, if $\omega \in \Omega_\delta$,

$$U_1(\tau c) < -\varepsilon/2 < \varepsilon/2 < U_1(\tau d).$$

But now, $y \mapsto U_1(y)$ is continuous, so there is a random variable $Y \in (\tau c, \tau d)$ such that $U_1(Y) = 0$. Then $U_t \stackrel{\text{def}}{=} U_t(Y)$ is a solution of the two-point boundary value problem (8.52). It is not identically zero, since $U'_0 = Y \neq 0$. Thus we have found a non-constant local solution.

To find a global solution, note that $U \equiv 0$ is always a solution, so that the function defined by

$$\bar{U}_t = \begin{cases} U_t(Y, \omega) & \text{if } \omega \in \Omega_0 \\ 0 & \text{otherwise} \end{cases}$$

is a global solution which is not identically zero. Since $U \equiv 0$ is also a global solution, this finishes the proof of (i).

(ii) As in (i), we begin by constructing solutions of the deterministic initial-value problem. This time, we want the solutions to return repeatedly to the origin. The key observation is that if a solution of the deterministic problem returns to the origin twice, it is actually periodic, and returns infinitely often. By rescaling, we can find solutions that have different numbers of zeros in $[0, 1]$. We then use the approximation theorem to find similar solutions to the stochastic problem.

By translating the problem, we may assume that $x_0 = 0$, so that $f(0) = 0$. By [8] there exists $b > 0$ such that if $|y| < b$, $T_1(y) < \infty$. Now $T_2(y) = T_1(y) + T_1(-y) < \infty$. It follows that the solution $\hat{U}_t(y)$ of the deterministic initial-value problem (8.54) is periodic, so that $T_m(y) < \infty$ for all m .

Now f is not affine on $(-x_1, x_1)$ so that by Lemma 8.3, $T_1(y)$ is not constant on $(-b, b)$. There are two cases. If $T_2(y)$ is not constant, then neither is $T_{2m}(y) = mT_2(y)$. On the other hand, if $T_2(y)$ is constant, then $T_{2m+1} = mT_2(y) + T_1(y)$ must be non-constant. We will do the first case in detail, and just indicate the modifications needed for the second case.

So let us suppose that $T_2(y)$ is differentiable but not constant on $(-b, b)$. Then there is $y \in (-b, b)$ for which $T_2'(y) \neq 0$, and

$$T_{2m}(y) = mT_2(y) \quad (8.55)$$

$$\hat{U}_{T_{2m}(y)} = \hat{U}_0(y) = 0 \quad (8.56)$$

$$\hat{U}'_{T_{2m}(y)}(y) = \hat{U}'_0(y) = y. \quad (8.57)$$

Since $y \mapsto T_2(y)$ is continuously differentiable, if it is not constant, there are $c_1 < c_2$ for which $T_2'(y) \neq 0$ in $[c_1, c_2]$. We will assume that $0 < c_1 < c_2$ and $T_2'(y) > 0$ in $[c_1, c_2]$. (The other cases are handled similarly.) Then $T_2(c_1) < T_2(c_2)$.

Let $k \geq 1$ be an integer and choose $\tau > 0$ such that

$$\frac{\tau}{T_2(c_1)} - \frac{\tau}{T_2(c_2)} > k.$$

Then there exist at least k consecutive integers, $m_1, \dots, m_k = m_1 + k - 1$ such that $\tau/T_2(c_2) < m_j < \tau/T_2(c_1)$, or

$$m_j T_2(c_1) < \tau < m_j T_2(c_2), \quad j = 1, \dots, k.$$

But $m_j T_2 = T_{2m_j}$, so that $T_{2m_j}(c_1) < \tau < T_{2m_j}(c_2)$. Thus by the intermediate value theorem, there is a unique $y_j \in (c_1, c_2)$ for which $T_{2m_j}(y_j) = \tau$, and $c_1 < y_1 < \dots < y_k < c_2$. Thus for small $\delta > 0$,

$$\hat{U}_\tau(y_j + \delta) < 0 < \hat{U}_\tau(y_j - \delta),$$

for $\hat{U}_\tau(y_j) = 0$ and $\hat{U}'_\tau(y_j) = y_j > 0$, while $T_2(y)$ increases with y .

We can choose $\epsilon > 0$ and $\delta > 0$ small enough so that the intervals $[y_j - \delta, y_j + \delta]$ are disjoint and

$$\hat{U}_\tau(y_j + \delta) < -\epsilon < \epsilon < \hat{U}_\tau(y_j - \delta).$$

Let $\hat{U}^{k,j}(y) = \hat{U}_{\tau t}(y/\tau)$. Then by Lemma 8.2 $\hat{U}^{k,j}(\tau y)$ satisfies the initial-value problem

$$\begin{cases} \hat{U}_t'' &= -\tau^2 f(\hat{U}_t) \\ \hat{U}_0 &= 0 \\ \hat{U}'_0 &= \tau y_j, \end{cases}$$

and moreover,

$$\hat{U}_1^{k,j}(\tau(y_j + \delta)) < -\epsilon < \epsilon < \hat{U}_1^{k,j}(\tau(y_j - \delta)). \quad (8.58)$$

We now approximate the $\hat{U}^{k,j}$ by solutions $U_t(y)$ of the stochastic initial-value problem (8.53). For $\rho > 0$ let $\Omega_{\rho,\tau} = \{\omega : |W_t(\omega) + \tau^2 t| < \rho, 0 \leq t \leq 1\}$. Note that $P\{\Omega_{\rho,\tau}\} > 0$ and, by Corollary 7.4 if ρ is small enough, $U_t(y)$ satisfies

$$\sup_{y \in [c_1, c_2]} \sup_{0 \leq t \leq 1} |\hat{U}_t^{k,j}(y) - U_t(\tau y)| < \frac{\varepsilon}{2}.$$

It follows from (8.58) that on $\Omega_{\rho, \tau}$,

$$\hat{U}_1(y_j + \delta) < -\varepsilon/2 < \varepsilon/2 < \hat{U}_1(y_j - \delta),$$

so, by the intermediate value theorem, there exist random variables $Y_j \in (\tau y_j - \tau\delta, \tau y_j + \tau\delta)$ such that $U_1(Y_j) = 0$. The Y_j are distinct because the intervals containing them are disjoint, so that the functions $0, U_t(Y_1), \dots, U_t(Y_k)$ are $k+1$ distinct (because their initial derivatives are unequal) solutions of the two point boundary-value problem (8.52). This shows that there are at least k distinct local solutions of the problem on the set $\Omega_{\rho, \tau}$. It is easy to construct global solutions from these, since $U \equiv 0$ is a global solution: for each j , let $U_t^{j,k}(\omega) = U_t(Y_j, \omega)$ for $\omega \in \Omega_{\rho, \tau}$ and $U_t^{j,k}(\omega) = 0$ otherwise. For each j and k this is a global solution, and there are infinitely many of them.

In the remaining case, where $T_2(y)$ is constant, $T_1(y)$ must be non-constant. It follows that for each m , $T_{2m+1}(y)$ is not constant, so we can simply use the above argument, replacing T_{2m} by T_{2m+1} . \square

8.2 Proof of Theorem 2.12

We will construct a deterministic equation whose coefficients are semi-linear approximations of f and g , and whose solutions have the desired behavior. We then compare the solutions of the stochastic and deterministic equations. Most of the work in the proof comes in handling the solutions for large values of the initial derivative, where we cannot use continuity arguments.

We begin with two lemmas. The first constructs the deterministic equation, and the second shows that when the initial derivative is large, the solution becomes large very quickly, regardless of the coefficients of the equation.

Let us first define semi-linear functions $f_0(x)$ and $g_0(x, y)$. Let

$$f_0(x) = \begin{cases} a_- x & \text{if } x < 0 \\ a_+ x & \text{if } x \geq 0. \end{cases}$$

and

$$g_0(x, y) = \begin{cases} b_- x + c_- y & \text{if } x < 0 \\ b_+ x + c_+ y & \text{if } x \geq 0 \end{cases}$$

and consider the deterministic problem

$$\begin{cases} U'' & = \theta f_0(U) + g_0(U, U') \\ U(0) & = 0 \\ U'(0) & = y \end{cases} \quad (8.59)$$

This is a second order ODE, and we can solve it explicitly: define $A_{\pm}(\theta) = \theta a_{\pm} + b_{\pm} - \frac{1}{4}c_{\pm}^2$. Let $\gamma \equiv \gamma(\theta) = |A_{\pm}(\theta)|^{\frac{1}{2}}$. Notice that if $y > 0$, U_t will be positive for small t , and if $y < 0$, it will be negative for small t . Then if $A_{\pm}(\theta) > 0$, the solution is

$$U_t = \begin{cases} (\gamma_+)^{-1} U'_0 e^{\frac{1}{2}c_+ t} \sinh(\gamma_+ t) & \text{if } y > 0, t \geq 0 \\ -(\gamma_-)^{-1} U'_0 e^{\frac{1}{2}c_- t} \sinh(\gamma_- t) & \text{if } y < 0, t \geq 0, \end{cases} \quad (8.60)$$

and if $A_{\pm} < 0$, the solution will change signs periodically:

$$U_t = \begin{cases} (\gamma_+)^{-1} U'_0 e^{\frac{1}{2}c_+ t} \sin(\gamma_+ t) & \text{if } y > 0, 0 \leq t \leq \pi/\gamma_+ \\ -(\gamma_-)^{-1} U'_0 e^{\frac{1}{2}c_- t} \sin(\gamma_- (t - \pi/\gamma_+)) & \text{if } y > 0, \pi/\gamma_+ \leq t \leq \pi/\gamma_+ + \pi/\gamma_-, \end{cases} \quad (8.61)$$

and the solution is periodic with period $\pi/\gamma_+ + \pi/\gamma_-$. If $y < 0$, U_t is initially negative, so it is given by (8.60) and (8.61) with $+$ and $-$ interchanged.

Remark 8.4 For future reference, let us note that if $y > 0$, U_t will be positive from 0 until $t_1 \equiv \pi/\gamma_+$, then negative until $t_2 \equiv \pi/\gamma_+ + \pi/\gamma_-$, then positive again until $t_3 \equiv t_2 + \pi/\gamma_+$, and so on. If $y < 0$, U_t will be negative until $s_1 \equiv \pi/\gamma_-$, then positive until $s_2 \equiv t_2$, then negative again until $s_3 \equiv t_2 + \pi/\gamma_+$, and so on. In both cases, U_t is periodic with period t_2 .

Note also that by (8.60), if $y > 0$, $U_t(y) = yU_t(1)$, while if $y < 0$, $U_t(y) = -yU_t(-1)$. Thus for the semi-linear equation, we need only know $U_t(1)$ and $U_t(-1)$. Furthermore, if $\bar{U}_t(y) \equiv \frac{1}{y}U_t(y)$, then $\bar{U}_t(y)$ equals $U_t(1)$ if $y > 0$, and it equals $U_t(-1)$ if $y < 0$.

Lemma 8.5 *Suppose either $a_+ \neq a_-$ or $a_+ = a_- \neq 0$ and $4b_- + c_-^2 \neq 4b_+ + c_+^2$. Let $U_t = U_t(y)$ be the solution of (8.59). Then it is possible to choose θ so that either $U_1(y) > 0$ for all $y \neq 0$ or $U_1(y) < 0$ for all $y \neq 0$.*

PROOF. This comes down to showing that we can choose θ so that $U_1(y)$ has the same sign for all y . Equivalently, we will show that $t = 1$ falls in the intersection of some interval of positivity of both $U_t(1)$ and $U_t(-1)$, or in some interval of negativity, i.e. either $1 \in (t_{2k}, t_{2k+1}) \cap (s_{2j-1}, s_{2j})$ or $1 \in (t_{2k-1}, t_{2k}) \cap (s_{2j}, s_{2j+1})$ for some j and k .

Note that the $A_{\pm}(\theta)$ are linear functions of θ , and consider the two lines in the z - θ plane: $z = A_+(\theta)$ and $z = A_-(\theta)$. If $a_+ = a_- \neq 0$ and $4b_- + c_-^2 \neq 4b_+ + c_+^2$, these lines are parallel but unequal, so that there is a θ_0 such that either

$$A_+(\theta_0) < -\pi^2 < A_-(\theta_0) \quad \text{or} \quad A_-(\theta_0) < -\pi^2 < A_+(\theta_0). \quad (8.62)$$

On the other hand, if $a_+ \neq a_-$, the two lines intersect: there is θ_1 such that $A_+(\theta_1) = A_-(\theta_1)$. If $A_+(\theta_1) \neq -\pi^2$, then one can find θ_0 such that (8.62) holds. But if $A_+(\theta_1) = -\pi^2$ and a_+ and a_- have opposite signs, then (8.62) holds for every value of $\theta_0 \neq \theta_1$. Thus, (8.62) holds in every case except when $a_+ \neq a_-$, $A_+(\theta_1) = -\pi^2$, and a_+ and a_- have the same sign.

Thus, let us first suppose that one of the two alternatives of (8.62) holds. Since the proof is virtually the same for the two, we will assume that $A_-(\theta_0) < -\pi^2 < A_+(\theta_0)$ and show that

$U_1(y) > 0$ for all $y \neq 0$. (If the other alternative holds, we simply show that $U_1(y) < 0$ for $y \neq 0$.)

Now if $y > 0$, $U_t^d(y)$ is either a positive constant times $\sinh(\gamma_+ t) > 0$, (if $A_+(\theta_0) > 0$) or a positive constant times $\sin(\gamma_+ t)$, (if $A_+(\theta_0) < 0$) which is again positive for $0 < t \leq 1$ since $\gamma_+ < \pi$. In either case, $U_1(y) > 0$. On the other hand, if $y < 0$, the solution is initially equal to a negative constant times $\sin(\gamma_- t)$. This holds until $t = \pi/\gamma_- < 1$, when it changes sign, and, depending on the sign of $A_+(\theta_0)$, equals a positive constant times either $\sin(\gamma_+(t - \pi/\gamma_-))$ or $\sinh(\gamma_+(t - \pi/\gamma_-))$. In both cases, $U_t(y)$ will still be positive at time $t = 1$.

This leaves the case where $a_+ \neq a_-$, a_+ and a_- have the same sign, and there exists θ_0 such that $A_+(\theta_0) = A_-(\theta_0) = -\pi^2$. The idea is nearly the same, but this time we use the fact that U is periodic, and wait for it to come around a second time. Assume both a_{\pm} are positive (the case where both are negative is similar) and, to be concrete, assume $0 \leq a_+ < a_-$. Then if $\theta < \theta_0$, $A_-(\theta) < A_+(\theta) < -\pi$, so, taking square roots, $\pi < \gamma_+(\theta) < \gamma_-(\theta)$, or $\pi/\gamma_-(\theta) < \pi/\gamma_+(\theta) < 1$. A little thought shows that we can choose $\theta < \theta_0$ such that

$$\frac{2\pi}{\gamma_-(\theta)} + \frac{\pi}{\gamma_+(\theta)} < 1 < \frac{\pi}{\gamma_-(\theta)} + \frac{2\pi}{\gamma_-(\theta)} \quad (8.63)$$

But now, recall the times s_j and t_j defined in Remark 8.4. It is easy to see from (8.63) that $t_2 < s_3 < 1 < t_3 < s_4$, so that $1 \in (t_2, t_3) \cap (s_3, s_4)$, which is an interval of positivity of $U_t(y)$ for both $y > 0$ and $y < 0$. In other words, for this value of θ , $U_1(y) > 0$ for all $y \neq 0$, as claimed. This finishes the proof. \square

Let $\Omega_K = \{\sup_{0 \leq t \leq 1} |W_t| \leq K\}$.

Lemma 8.6 *Let $0 \leq \tau \leq 1$ be a (possibly random) time, and let $\hat{y} = V_\tau(y)$. Given $M > 0$, let $T_M(\hat{y}) = \inf\{t > \tau : |U_t - U_\tau| > M\}$. Then for $K > 0$ and $\rho > 0$, there exists $N = N(\rho, M, K)$ such that on Ω_K , if $|\hat{y}| \geq N$, and $|U_\tau| \leq M$,*

(i) $(1 - \rho)|\hat{y}| \leq |V_t(y) - V_\tau(y)| \leq (1 + \rho)|\hat{y}|$ for $\tau \leq t \leq T_M(y)$;

(ii) $\frac{M}{(1+\rho)\hat{y}} \leq T_M(y) - \tau \leq \frac{M}{(1-\rho)\hat{y}}$

The same is true of the solution to (8.59) if $|\theta| < K$.

PROOF. It is intuitively evident that if the initial velocity is large, the velocity will remain large for a certain time, regardless of f and g , so that T_M will be small. The work comes in verifying the uniformity in \hat{y} . We rescale: let $\bar{U}_t = U_t/\hat{y}$, and $\bar{V}_t = V_t/\hat{y}$.

Notice that (ii) follows from (i), since if we integrate (i), we see that $(1 - \rho)(t - \tau) \leq \bar{U}_t - \bar{U}_\tau \leq (1 + \rho)(t - \tau)$. Set $\bar{U}_t - \bar{U}_\tau = M/\hat{y}$ to see (ii).

It remains to show (i). To do this, integrate (3.9) by parts, setting $x = 0$:

$$V_t = \hat{y} + W_t f(U_\tau) + \int_\tau^t (W_t - W_s) f'(U_s) V_s ds + \int_\tau^t g(U_s, V_s) ds. \quad (8.64)$$

Now $|W_t| \leq K$ on Ω_K , so $|W_t - W_s| \leq 2K$, and $|f'(U_s)| \leq L$ since f is Lipschitz with constant L . Similarly, $|g(U_s, V_s)| \leq L(1 + |U_s| + |V_s|)$. But since $|U_\tau| \leq M$ and $t \leq T_M(y)$, $|U_s - U_\tau| \leq M$ so in fact $|g(U_s, V_s)| \leq L(2M + 1 + |V_s|)$. Let $f^* \equiv \sup\{|f(x)| : |x| \leq M\}$. Divide (8.64) by \hat{y} :

$$|\bar{V}_t| \leq 1 + Kf^*/\hat{y} + 2KL \int_0^t |\bar{V}_s| ds + (2M+1)Lt/\hat{y} + L \int_0^t |\bar{V}_s| ds.$$

Now $t \leq 1$, so, collecting terms,

$$|\bar{V}_t| \leq 1 + \frac{Kf^* + (M+1)L}{\hat{y}} + (2K+1)L \int_0^t |\bar{V}_s| ds. \quad (8.65)$$

By Gronwall,

$$|\bar{V}_t| \leq \left(1 + \frac{Kf^* + (2M+1)L}{\hat{y}}\right) e^{2(K+1)Lt}.$$

This is bounded uniformly in \hat{y} on $0 \leq t \leq 1$, $|\hat{y}| > 1$, say $|\bar{V}_t| \leq C_2$.

Return to (8.64) and repeat the above steps to see that, as $t \leq 1$,

$$|\bar{V}_t - 1| \leq \frac{Kf^* + (2M+1)L}{\hat{y}} + (2K+1)L \int_0^t |\bar{V}_s| ds \leq \frac{C}{\hat{y}} + D(t - \tau)$$

for constants C and D . Integrating, we see that

$$\left(1 - \frac{C}{\hat{y}}\right)(t - \tau) - D(t - \tau)^2 \leq \bar{U}_t - \bar{U}_\tau \leq \left(1 + \frac{C}{\hat{y}}\right)(t - \tau) + D(t - \tau)^2$$

But $T_M(y)$ is the first time $\bar{U}_t(y) - \bar{U}_\tau(y)$ hits M/\hat{y} , and it clearly decreases to τ as $\hat{y} \rightarrow \infty$. Thus there is some $N > N_1$ such that if $\hat{y} > N$, $C/\hat{y} + D(T_M(\hat{y}) - \tau) < \rho$, and (i) follows. This finishes the proof for the solution of the stochastic equation. Since the only thing we used about Brownian motion above was that $|W_t| \leq K$, it also applies to the solution of (8.59) as soon as $|\theta| \leq K$. \square

PROOF. (Of Theorem 2.12.) We will show that there is a set Ω_δ and a r.v. Z such that the solution of the initial value problem $U_1(y)$ is either strictly greater than Z for all y , or strictly less than Z for all y . This implies that there is no global solution of the two-point problem

$$\begin{cases} U'' &= f(U) \circ \dot{W} + g(U, V) \\ U_0 &= 0, \quad U_1 = Z. \end{cases}$$

In order to handle the case of large initial derivatives y , we note that if y is large, the solution will spend almost all its time in the region $|U_t| > M$, and in this region, f and f_0 are close.

The key step is to show that for δ small enough and N large enough, if $\omega \in \Omega_\delta$, then either

$$U_1(y) > 0, \quad \forall |y| > N \text{ or } U_1(y) < 0, \quad \forall |y| > N. \quad (8.66)$$

Once we have this, we simply note that $y \mapsto U_1(y)$ is continuous, so $\max_{|y| \leq N} |U_1(y)|$ is finite, hence $U_1(y)$ is bounded either below or above, and we can take

$$Z \equiv \min_y U_1(y) - 1 \text{ or } Z \equiv \max_y U_1(y) + 1$$

to prove the theorem.

It remains to prove (8.66). Let $(U_t^d(y), V_t^d(y))$ be the solution of the deterministic initial-value problem (8.59), given explicitly by (8.60), and let $(U_t(y), V_t(y))$ be the solution of the stochastic initial-value problem

$$\begin{cases} U'' &= f(U) \circ \dot{W} + g(U, V) \\ U_0 &= 0, \quad U'_0 = y. \end{cases} \quad (8.67)$$

By Lemma 8.5, we can choose $\theta = \theta_0$ in order to have $U_1^d(y) > 0$ for all $y > 0$. Recall that $U_t^d(y) = |y|U_t^d(\text{sgn } y)$, so that if we set $\rho_1 = \frac{1}{2} \min\{U_1^d(1), U_1^d(-1)\}$, we have

$$U_1^d(y) \geq 2|y|\rho_1 > 0.$$

We will go through the proof in detail in the case where $A_+(\theta_0) < -\pi^2 < A_-(\theta_0) < 0$, where A_\pm is defined above. The other cases can be done by similar reasoning. In that case, Lemma 8.5 tells us that $y > 0 \implies U_t^d(y)$ is positive for $t \leq 1$, while $y < 0 \implies U_t^d(y)$ is negative if $0 < t < s_1$ and positive if $s_1 < t \leq 1$ for $s_1 = \pi/\sqrt{A_-(\theta_0)}$.

Suppose $y > 0$. Let $\epsilon > 0$ and choose M large enough so that $|x| > M/2$ implies that

$$\begin{aligned} |f'(x) - f'_0(x)| &< \epsilon \\ |f(x) - f_0(x)| &< \epsilon|x| \\ |g(x, y) - g_0(x, y)| &< \epsilon(|x| + |y|). \end{aligned} \quad (8.68)$$

Let $T_M(y) = \inf\{t : U_t(y) = M\}$, and $T_M^d(y) = \inf\{t : U_t^d(y) = M\}$. Let $\tau_1(y) = \max\{T_M(y), T_M^d(y)\}$, and $\tau_2(y) = \sup\{t < 1 : U_t(y) > M, U_t^d(y) > M\}$, so both $U_t^d(y)$ and $U_t(y)$ are greater than M on (τ_1, τ_2) . Define $\Omega_\delta = \{\omega : |W_t(\omega) - \theta_0 t| < \delta, 0 \leq t \leq 1\}$.

We rescale the functions: let

$$\bar{U}_t = \frac{1}{|y|}U_t, \quad \bar{V}_t = \frac{1}{|y|}V_t, \quad \bar{U}_t^d = \frac{1}{|y|}U_t^d \text{ and } \bar{V}_t^d = \frac{1}{|y|}V_t^d.$$

By Lemma 8.6, if $\rho > 0$, we can choose N large enough so that if $y > N$ and $\omega \in \Omega_\delta$

$$|\bar{V}_t(y) - 1| \leq \rho, \quad 0 \leq t \leq T_M(y), \quad |\bar{V}_t^d(y) - 1| \leq \rho, \quad 0 \leq t \leq T_M^d(y), \quad (8.69)$$

and

$$|T_M(y) - M/y| < 2\rho M/y, \quad |T_M^d(y) - M/y| < 2\rho M/y. \quad (8.70)$$

Thus $|T_M(y) - T_M^d(y)| < 2M\rho/y$. Let $U_{\tau_1}(y) = x_1$, $U_{\tau_1}^d(y) = x_d$. Now $U_{T_m} = U_{T_M^d}^d = M$. Since $V_t(y)$ is bounded by $(1 + \rho)y$ on $0 \leq t \leq T_M(y)$, as is V^d , we see that $|x_1 - x_d| \leq 4M\rho(1 - \rho)$.

Consider V and V^d on the interval (τ_1, τ_2) . Both $U_t(y)$ and $U_t^d(y)$ are greater than M there. Integrating (3.9) by parts

$$V_t = x_1 + (W_t - W_{\tau_1})f(x_1) + \int_{\tau_1}^t (W_t - W_s)f'(U_s)V_s ds + \int_{\tau_1}^t g(U_s, V_s) ds \quad (8.71)$$

and

$$V_t^d = x_d + \theta_0(t - \tau_1)f_0(x_d) + \int_{\tau_1}^t \theta_0(t - s)f'_0(U_s)V_s ds + \int_{\tau_1}^t g_0(U_s, V_s) ds \quad (8.72)$$

Divide both equations by y and subtract. Notice that on Ω_δ , $|W_t - W_{\tau_1} - \theta_0(t - \tau_1)| \leq 2\delta$ and, because $U_s \geq M$, $|f'(U_s) - f'_0(U_s^d)| < \epsilon$, so

$$\begin{aligned} |(W_t - W_s)f'(U_s)\bar{V}_s - \theta_0(t - s)f'_0(U_s^d)\bar{V}_s^d| & \leq |W_t - W_s||f'(U_s)||\bar{V}_s - \bar{V}_s^d| + |W_t - W_s||f'(U_s) - f'_0(U_s^d)||\bar{V}_s^d| \\ & \quad + |W_t - W_{\tau_1} - \theta_0(t - \tau_1)||f'_0(U_s^d)||\bar{V}_s^d| \\ & \leq L(\theta_0 + 2\delta)|\bar{V}_s - \bar{V}_s^d| + (\theta_0 + 2\delta)\epsilon + 2\delta L\|\bar{V}^d\|_\infty. \end{aligned}$$

Now \bar{V}_t^d is bounded, independent of y , so $\|V^d\|_\infty < \infty$, so the above is

$$\leq a_0|\bar{V}_s - \bar{V}_s^d| ds + a_1\epsilon + a_2\delta.$$

for some constants a_0 , a_1 , and a_2 . Similarly, g is Lipschitz and $U_s^d > M$ if $\tau_1 < t < \tau_2$, so

$$\begin{aligned} \frac{1}{y}|g(U_s, V_s) - g_0(U_s^d, V_s^d)| & \leq \frac{1}{y}|g(U_s, V_s) - g(U_s^d, V_s^d)| + |g(U_s^d, V_s^d) - g_0(U_s^d, V_s^d)| \\ & \leq L(|\bar{U}_s - \bar{U}_s^d| + |\bar{V}_s - \bar{V}_s^d|) + \epsilon(\|U^d\|_\infty + \|V^d\|_\infty) \end{aligned}$$

Now $t - \tau < 1$ so $\int_\tau^t |\bar{U}_s - \bar{U}_s^d| ds \leq \int_0^t \frac{|x_1 - x_d|}{y} + \int_\tau^s |\bar{V}_u - V_u^d| du ds \leq \frac{|x_1 - x_d|}{y} + \int_\tau^t |\bar{V}_s - V_s^d| ds$ so that we can collect terms above to see that for some constants A , B , C , and D ,

$$|\bar{V}_t - \bar{V}_t^d| \leq A \frac{|x_1 - x_d|}{y} + B\epsilon + C\delta + D \int_\tau^t |\bar{V}_s - \bar{V}_s^d| ds.$$

Let $\epsilon_1 > 0$. We can choose ϵ and δ small enough and $|y|$ large enough so that the right-hand side is bounded by $\epsilon_1 + D \int_\tau^t |\bar{V}_s - \bar{V}_s^d| ds$. Thus by Gronwall,

$$|\bar{V}_t - \bar{V}_t^d| \leq \epsilon_1 e^{D(t-\tau)} \leq \epsilon_1 e^D$$

Thus, given $\rho > 0$, we can choose N such that if ϵ and δ are small enough and $|y| > N$, $|\bar{V}_t - \bar{V}_t^d| \leq \rho$ for all $\tau \leq t \leq 1$. It follows that $|\bar{U}_t - \bar{U}_t^d| < \rho$ as well. Since $\rho < (1/2)U_1^d(1)$, then

$$|\bar{U}_1 - \bar{U}_1^d| \leq \frac{|x_1 - x_d|}{y} + |\bar{V}_1 - \bar{V}_1^d| \leq \rho + \rho < U_1^d(1).$$

It follows that $\bar{U}_1(y) > 0$ for all $|y| > N$. But this means that $U_1(y) > 0$ for all $|y| > N$, as we claimed.

Now consider the case $y < 0$. The proof is quite similar, with one extra step. Indeed, the idea is that during the time that the solutions are greater than M in absolute value, the coefficients of the deterministic and stochastic equations are very nearly the same, so the solutions are close. On the other hand, if y is large, the times when they are smaller than M in absolute value are so short that the solutions have to behave in approximately the same way.

The solution will initially be negative. The solution to the deterministic equation will return to zero at a time $s_1 < 1$, and will then stay positive from s_1 to 1. We want to show that the stochastic equation does the same thing. Let $T_{-M}(y)$ and $T_{-M}^d(y)$ be the first times that $U_t(y)$ and $U_t^d(y)$ respectively hit $-M$, and let $T'_{-M}(y)$ and $T_{-M}^{d'}(y)$ be the next time they (respectively) hit $-M$. Let τ_1 be the maximum of $T_{-M}(y)$ and $T_{-M}^d(y)$, and let τ_2 be the minimum of $T'_{-M}(y)$ and $T_{-M}^{d'}(y)$. By Lemma 8.6, τ_1 decreases to zero as y tends to $-\infty$, so for large enough y , $\tau_1 < \tau_2 < 1$. Moreover, the deterministic function is symmetric about its minimum, so that $V_{T_{-M}^d(y)}^d = -V_{T_{-M}^d(y)}^d$. If we apply the same argument we used for $y > 0$, but this time on $(0, \tau_2)$ instead of $(0, 1)$, we see that both $|\bar{U}_{\tau_2}(y) - \bar{U}_{\tau_2}^d| < \rho$ and $|\bar{V}_{\tau_2}(y) - \bar{V}_{\tau_2}^d| < \rho$. As $V_{T_{-M}^d(y)}^d = -V_{T_{-M}^d(y)}^d > (1 - \rho)|y|$, (again by Lemma 8.5), it follows that $V_{T_{-M}(y)}(y) < (1 - 2\rho)|y|$. Now apply Lemma 8.5 again with $\tau = T_{-M}(y)$ and $\hat{y} = V_\tau(y)$ to see that $U_t(y)$ hits zero again before time $t = 1$, and that at that time, $V \geq (1 - 3\rho)|y|$. The process then re-starts from $U = 0$, $V \geq (1 - 3\rho)|y|$, so the argument for $y > 0$ again applies to show that U_t will still be positive at time $t = 1$. Thus, if we choose N large enough, if $\omega \in \Omega_\delta$ and $|y| > N$, $U_1(y) \geq 0$ for all y . This finishes the proof. \square

References

- [1] Dembo, A. and Zeitouni O., A change of variables formula for Stratonovich integrals and existence of solutions for two-point stochastic boundary value problems, *Prob. Th. Rel Fields* 84 (1990), 411–425.
- [2] Krener, A., Reciprocal processes and the stochastic realization problem of a causal system, in Byrnes, C.Z., Lindquist, E. (Eds) *Modelling, Identification, and Robust Control*. Elsevier/Noble, Holland, 1986
- [3] Kunita, H., *Stochastic differential equations and stochastic flow of diffeomorphisms*, *Lecture Notes in Math.*, 1047, Springer-Verlag 1984, 144–300
- [4] Nualart, D. and Pardoux, E., Second order stochastic differential equations with Dirichet boundary conditions, *Stochastic Proc. Appl.*, 34 (1991) 1–24
- [5] Nualart, D. and Pardoux, E., Stochastic calculus with anticipative integrals, *Prob. Th. Rel. Fields*, 78 (1988) 535–581.
- [6] Nualart, D. and Zakai, M., Generalized stochastic integrals and the Malliavin calculus, *Prob. Th. Rel. Fields* 73, (1986) 255–280.
- [7] Ocone, D. and Pardoux, E., (1989) Linear stochastic differential equations with boundary conditions, *Prob. Th. Rel. Fields* 82 (1989) 489–526.
- [8] Schaaf, R., Global solution branches of two-point boundary value problems, *Lecture Notes in Math.* 1458, 1990.
- [9] Watanabe, S., *Lectures on stochastic differential equations and Malliavin calculus*, Springer Verlag, Berlin/Heidelberg/New York/Tokyo, 1984.