

Research Article

Iterative Algorithms with Variable Coefficients for Asymptotically Strict Pseudocontractions

Ci-Shui Ge,¹ Jin Liang,² and Ti-Jun Xiao³

¹ Department of Mathematics and Physics, Anhui University of Architecture, Hefei, Anhui 230022, China

² Department of Mathematics, Shanghai Jiao Tong University, Shanghai 200240, China

³ School of Mathematical Sciences, Fudan University, Shanghai 200433, China

Correspondence should be addressed to Jin Liang, jliang@ustc.edu.cn

Received 8 October 2009; Revised 29 November 2009; Accepted 22 January 2010

Academic Editor: Anthony To Ming Lau

Copyright © 2010 Ci-Shui Ge et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce and study some new CQ-type iterative algorithms with variable coefficients for asymptotically strict pseudocontractions in real Hilbert spaces. General results for asymptotically strict pseudocontractions are established. The main result extends the previous results.

1. Introduction

Let H be a real Hilbert space, C a nonempty closed convex subset of H , $T : C \rightarrow C$ a self-mapping of C and $\text{Fix}(T) := \{x \in C : Tx = x\}$.

Recall that a mapping $T : C \rightarrow C$ is called to be nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in C. \quad (1.1)$$

T is called to be asymptotically nonexpansive [1] if there exists a sequence $\{k_n\}$ with $k_n \geq 1$ and $\lim_{n \rightarrow \infty} k_n = 1$ such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\|, \quad \forall x, y \in C, \text{ and all integers } n \geq 1. \quad (1.2)$$

T is called to be an asymptotically κ -strict pseudocontraction, if there exist $0 \leq \kappa < 1$ and $0 \leq \gamma_n \rightarrow 0$ ($n \rightarrow \infty$) such that

$$\|T^n x - T^n y\|^2 \leq (1 + \gamma_n) \|x - y\|^2 + \kappa \|(I - T^n)x - (I - T^n)y\|^2 \quad (1.3)$$

for all $x, y \in C$ and all integers $n \geq 1$.

As $\kappa = 0$, asymptotically κ -strict pseudocontraction T is asymptotically nonexpansive.

In [2], Nakajo and Takahashi studied the iterative approximation of fixed points of nonexpansive mappings and proved the following strong convergence theorem.

Theorem A. *Let C be a nonempty closed convex subset of a Hilbert space H and let T be a nonexpansive mapping of C into itself such that $\text{Fix}(T) \neq \emptyset$. Suppose $\{x_n\}$ is given by*

$$\begin{aligned} x_0 &\in C \text{ chosen arbitrarily,} \\ y_n &= \alpha_n x_n + (1 - \alpha_n) T x_n, \\ C_n &= \{z \in C : \|y_n - z\| \leq \|x_n - z\|\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0, \quad n \in \mathbb{N}, \end{aligned} \tag{1.4}$$

where $P_{C_n \cap Q_n}$ is the metric projection from C onto $C_n \cap Q_n$ and α_n is chosen so that $0 \leq \alpha_n \leq a < 1$. Then, $\{x_n\}$ converges strongly to $P_{\text{Fix}(T)} x_0$, where $P_{\text{Fix}(T)}$ is the metric projection from C onto $\text{Fix}(T)$.

Such algorithm in (1.4) is referred to be the (CQ) algorithm in [3], due to the fact that each iterate x_{n+1} is obtained by projecting x_0 onto the intersection of the suitably constructed closed convex sets C_n and Q_n . It is known that the (CQ) algorithm in (1.4) is of independent interest, and the (CQ) algorithm has been extended to various mappings by many authors (cf., e.g., [3–11]).

Very recently, by extending the (CQ) algorithm, Takahashi et al. [9] studied a family of nonexpansive mappings and gave some good strong convergence theorems. Kim and Xu [5] extended the (CQ) algorithm to study asymptotically κ -strict pseudocontractions and established the following interesting result with the help of some boundedness conditions.

Theorem B. *Let C be a closed convex subset of a Hilbert space H and let $T : C \rightarrow C$ be an asymptotically κ -strict pseudocontractions for some $0 \leq \kappa < 1$. Assume that the fixed point set $\text{Fix}(T)$ of T is nonempty and bounded. Let $\{x_n\}_{n=0}^\infty$ be the sequence generated by the following (CQ) algorithm:*

$$\begin{aligned} x_0 &\in C, \text{ chosen arbitrarily,} \\ y_n &= \alpha_n x_n + (1 - \alpha_n) T^n x_n, \\ C_n &= \left\{ z \in C : \|y_n - z\|^2 \leq \|x_n - z\|^2 + [\kappa - \alpha_n(1 - \alpha_n)] \|x_n - T x_n\|^2 + \theta_n \right\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0, \end{aligned} \tag{1.5}$$

where

$$\theta_n = \Delta_n^2 (1 - \alpha_n) \gamma_n \longrightarrow 0 \quad (n \longrightarrow \infty), \quad \Delta_n = \sup \{\|x_n - z\| : z \in \text{Fix}(T)\} < \infty. \tag{1.6}$$

Assume that control sequence $\{\alpha_n\}_{n=0}^\infty$ is chosen so that $\limsup_{n \rightarrow \infty} \alpha_n < 1 - \kappa$. Then $\{x_n\}$ converges strongly to $P_{\text{Fix}(T)} x_0$.

It is our purpose in this paper to try to obtain some new fixed point theorems for asymptotically strict pseudocontractions without the boundedness conditions as in Theorem B. Motivated by Nakajo and Takahashi [2], Takahashi et al. [9], and Kim and Xu [5], we introduce and study certain new CQ-type iterative algorithms with variable coefficients for asymptotically strict pseudocontractions in real Hilbert spaces. Our results improve essentially the corresponding results of [5].

2. Results and Proofs

Throughout this paper,

- (i) $x_n \rightharpoonup x$ means that $\{x_n\}$ converges weakly to x .
- (ii) $x_n \rightarrow x$ means that $\{x_n\}$ converges strongly to x .
- (iii) $\omega_w(x_n) := \{x : \exists x_{n_j} \rightharpoonup x\}$, that is, the weak ω -limit set of $\{x_n\}$.
- (iv) $B_r(x_0) := \{x \in H : \|x - x_0\| \leq r\}$.
- (v) \mathbb{N} is the set of nonnegative integers.

The following lemmas are basic (cf., e.g., [6] for Lemma 2.1, and [5] for Lemmas 2.2-2.3).

Lemma 2.1. *Let K be a closed convex subset of a real Hilbert space H . Given $x \in H, z \in K$. Then $z = P_K x$ if and only if*

$$\langle x - z, y - z \rangle \leq 0, \quad \forall y \in K, \quad (2.1)$$

where $P_K x$ is the unique point in K with the property

$$\|x - P_K x\| \leq \|x - y\|, \quad \forall y \in K. \quad (2.2)$$

Lemma 2.2. *Let K be a closed convex subset of a real Hilbert space H , $\{x_n\} \subset H, u \in H$, and $q = P_K u$. Suppose that $\{x_n\}$ satisfies*

$$\|x_n - u\| \leq \|u - q\|, \quad \forall n \in \mathbb{N}, \quad (2.3)$$

and $\omega_w(x_n) \subset K$. Then $x_n \rightarrow q$.

Lemma 2.3. *Let C be a closed convex subset of a Hilbert space H and $T : C \rightarrow C$ an asymptotically κ -strict pseudocontraction. Then*

(I) *for each $n \geq 1, T^n$ satisfies the Lipschitz condition:*

$$\|T^n x - T^n y\| \leq L_n \|x - y\|, \quad \forall x, y \in C, \quad (2.4)$$

where

$$L_n = \frac{\kappa + \sqrt{1 + \gamma_n(1 - \kappa)}}{1 - \kappa}, \quad \{\gamma_n\} \text{ is as in (1.3);} \quad (2.5)$$

(II) if $\{x_n\}$ is a sequence in C such that $x_n \rightarrow \tilde{x}$ and

$$\limsup_{m \rightarrow \infty} \limsup_{n \rightarrow \infty} \|x_n - T^m x_n\| = 0, \quad (2.6)$$

then

$$(I - T)x_n \rightarrow 0 \implies (I - T)\tilde{x} = 0. \quad (2.7)$$

In particular,

$$x_n \rightarrow \tilde{x}, \quad (I - T)x_n \rightarrow 0 \implies (I - T)\tilde{x} = 0. \quad (2.8)$$

(III) $\text{Fix}(T)$ is closed and convex so that the projection $P_{\text{Fix}(T)}$ is well defined.

Theorem 2.4. Let C be a closed convex subset of a Hilbert space H , $T : C \rightarrow C$ an asymptotically κ -strict pseudocontraction for some $0 \leq \kappa < 1$, and $\text{Fix}(T) \neq \emptyset$. Let $\{x_n\}$ be the sequence generated by the following CQ-type algorithm with variable coefficients:

$$\begin{aligned} x_0 &\in C \text{ chosen arbitrarily,} \\ y_n &= (1 - \hat{\beta}_n)x_n + \hat{\beta}_n T^n x_n, \\ C_n &= \{z \in C : \|y_n - z\|^2 \leq \|x_n - z\|^2 + \hat{\beta}_n(\kappa + \hat{\beta}_n - 1)\|x_n - T^n x_n\|^2 + \theta_n\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0, \quad n \in \mathbb{N}, \end{aligned} \quad (2.9)$$

where

$$\hat{\beta}_n = \frac{\beta_n}{1 + \|x_n - x_0\|^2}, \quad \beta_n \in \left[\frac{1}{2}, 1\right], \quad \theta_n = 2(1 + r_0^2)\beta_n\gamma_n, \quad (2.10)$$

the sequence $\{\beta_n\}$ is chosen so that $\beta_n \rightarrow 1$ ($n \rightarrow \infty$), the positive real number r_0 is chosen so that $B_{r_0}(x_0) \cap \text{Fix}(T) \neq \emptyset$, and $\{\gamma_n\}$ is as in (1.3). Then $\{x_n\}$ converges strongly to $P_{\text{Fix}(T)}x_0$.

Proof. We divide the proof into five steps.

Step 1. We prove that $C_n \cap Q_n$ is nonempty, convex and closed.

Clearly, both Q_n and C_n are convex and closed, so is $C_n \cap Q_n$. Since $T : C \rightarrow C$ is an asymptotically κ -strict pseudocontraction, we have by (1.3),

$$\begin{aligned} \|T^n x - p\|^2 &\leq (1 + \gamma_n)\|x - p\|^2 + \kappa\|(I - T^n)x - (I - T^n)p\|^2 \\ &\leq (1 + \gamma_n)\|x - p\|^2 + \kappa\|x - T^n x\|^2, \end{aligned} \quad (2.11)$$

for all $x \in C$, $p \in \text{Fix}(T)$, and all integers $n \geq 1$.

By (2.9) and (2.11), we deduce that for each $p \in B_{r_0}(x_0) \cap \text{Fix}(T)$, $n \in \mathbb{N}$,

$$\begin{aligned}
\|y_n - p\|^2 &= \left\| (1 - \widehat{\beta}_n)(x_n - p) + \widehat{\beta}_n(T^n x_n - p) \right\|^2 \\
&= (1 - \widehat{\beta}_n)\|x_n - p\|^2 + \widehat{\beta}_n\|T^n x_n - p\|^2 - \widehat{\beta}_n(1 - \widehat{\beta}_n)\|x_n - T^n x_n\|^2 \\
&= (1 - \widehat{\beta}_n)\|x_n - p\|^2 + \widehat{\beta}_n \left[(1 + \gamma_n)\|x_n - p\|^2 + \kappa\|x_n - T^n x_n\|^2 \right] \\
&\quad - \widehat{\beta}_n(1 - \widehat{\beta}_n)\|x_n - T^n x_n\|^2 \\
&\leq \|x_n - p\|^2 + \widehat{\beta}_n(\kappa + \widehat{\beta}_n - 1)\|x_n - T^n x_n\|^2 + \beta_n \gamma_n \frac{2(\|x_n - x_0\|^2 + \|x_0 - p\|^2)}{1 + \|x_n - x_0\|^2} \\
&\leq \|x_n - p\|^2 + \widehat{\beta}_n(\kappa + \widehat{\beta}_n - 1)\|x_n - T^n x_n\|^2 + 2(1 + r_0^2)\beta_n \gamma_n \\
&= \|x_n - p\|^2 + \widehat{\beta}_n(\kappa + \widehat{\beta}_n - 1)\|x_n - T^n x_n\|^2 + \theta_n.
\end{aligned} \tag{2.12}$$

Therefore,

$$B_r(x_0) \cap \text{Fix}(T) \subset C_n, \quad \forall n \in \mathbb{N}. \tag{2.13}$$

Next, we prove by induction that

$$B_{r_0}(x_0) \cap \text{Fix}(T) \subset Q_n, \quad \forall n \in \mathbb{N}. \tag{2.14}$$

Obviously, $B_{r_0}(x_0) \cap \text{Fix}(T) \subset C = Q_0$, that is, (2.14) holds for $n = 0$. Assume that $B_{r_0}(x_0) \cap \text{Fix}(T) \subset Q_n$ for some $n \in \mathbb{N}$. Then, (2.13) implies that $B_{r_0}(x_0) \cap \text{Fix}(T) \subset C_n \cap Q_n \neq \emptyset$ and $x_{n+1} = P_{C_n \cap Q_n} x_0$ is well defined.

By Lemma 2.1, we get $\langle x_{n+1} - z, x_0 - x_{n+1} \rangle \geq 0, \forall z \in C_n \cap Q_n$. In particular, for each $z \in B_{r_0}(x_0) \cap \text{Fix}(T)$, we have $\langle x_{n+1} - z, x_0 - x_{n+1} \rangle \geq 0$. This together with the definition of Q_{n+1} , the inequality (2.14) holds for $n + 1$. So (2.14) is true.

Step 2. We prove that $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$.

By the definition of Q_n and Lemma 2.1, we get $x_n = P_{Q_n} x_0$. Hence,

$$\|x_n - x_0\| \leq \|p - x_0\|, \quad \forall p \in B_{r_0}(x_0) \cap \text{Fix}(T). \tag{2.15}$$

Denoting $M := \|x_0\| + \|p - x_0\|$, we have $\|x_n\| \leq M$, for all $n \in \mathbb{N}$, and

$$\|x_n - x_0\| \leq \|q - x_0\|, \quad \forall n \in \mathbb{N}, \tag{2.16}$$

where $q = P_{\text{Fix}(T)}x_0 \in B_{r_0}(x_0) \cap \text{Fix}(T)$. The definition of x_{n+1} shows that $x_{n+1} \in Q_n$, that is, $\langle x_{n+1} - x_n, x_n - x_0 \rangle \geq 0$. This implies that

$$\begin{aligned} \|x_{n+1} - x_n\|^2 &= \|x_{n+1} - x_0\|^2 - \|x_n - x_0\|^2 - 2\langle x_{n+1} - x_n, x_n - x_0 \rangle \\ &\leq \|x_{n+1} - x_0\|^2 - \|x_n - x_0\|^2. \end{aligned} \quad (2.17)$$

Thus $\{\|x_n - x_0\|\}$ is increasing. Since $\{x_n\}$ is bounded, $\lim_{n \rightarrow \infty} \|x_n - x_0\|$ exists and

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (2.18)$$

Step 3. We prove that $\lim_{n \rightarrow \infty} \|x_n - T^n x_n\| = 0$.

The definition of x_{n+1} shows that $x_{n+1} \in C_n$, that is,

$$\|y_n - x_{n+1}\|^2 \leq \|x_n - x_{n+1}\|^2 + \hat{\beta}_n (\kappa + \hat{\beta}_n - 1) \|x_n - T^n x_n\|^2 + \theta_n. \quad (2.19)$$

By (2.19) and the definition of y_n in (2.9), we deduce that

$$\begin{aligned} \hat{\beta}_n^2 \|x_n - T^n x_n\|^2 &= \|y_n - x_n\|^2 \\ &\leq \|y_n - x_{n+1}\|^2 + \|x_{n+1} - x_n\|^2 + 2\|y_n - x_{n+1}\| \cdot \|x_{n+1} - x_n\| \\ &\leq \hat{\beta}_n (\kappa + \hat{\beta}_n - 1) \|x_n - T^n x_n\|^2 + \theta_n + 2\|x_{n+1} - x_n\|^2 + 2\|y_n - x_{n+1}\| \cdot \|x_{n+1} - x_n\|. \end{aligned} \quad (2.20)$$

Further, we have

$$(1 - \kappa) \hat{\beta}_n \|x_n - T^n x_n\|^2 \leq 2\|x_{n+1} - x_n\|^2 + 2\|x_{n+1} - x_n\| \cdot \|y_n - x_{n+1}\| + \theta_n. \quad (2.21)$$

Thus, (2.19) and (2.21) imply that

$$\begin{aligned} (1 - \kappa) \hat{\beta}_n \|x_n - T^n x_n\|^2 &\leq 4\|x_{n+1} - x_n\|^2 + 2\|x_{n+1} - x_n\| \cdot \|x_n - T^n x_n\| \sqrt{\hat{\beta}_n |\kappa + \hat{\beta}_n - 1|} \\ &\quad + 2\|x_{n+1} - x_n\| \sqrt{\theta_n} + \theta_n. \end{aligned} \quad (2.22)$$

Noticing $\|x_n\| \leq M, \beta_n \in [1/2, 1]$, we get

$$\hat{\beta}_n = \frac{\beta_n}{1 + \|x_n\|^2} \geq \frac{1}{2(1 + M^2)} > 0. \quad (2.23)$$

From $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$, $\lim_{n \rightarrow \infty} \theta_n = 0$, and (2.22), it follows that

$$\lim_{n \rightarrow \infty} \|x_n - T^n x_n\| = 0. \quad (2.24)$$

Step 4. We prove that

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0. \quad (2.25)$$

By Lemma 2.3 and the definition of T , we obtain

$$\begin{aligned} \|x_n - Tx_n\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - T^{n+1}x_{n+1}\| + \|T^{n+1}x_{n+1} - T^{n+1}x_n\| + \|T^{n+1}x_n - Tx_n\| \\ &\leq (1 + L_{n+1})\|x_{n+1} - x_n\| + \|x_{n+1} - T^{n+1}x_{n+1}\| + L_1\|x_n - T^n x_n\|, \end{aligned} \quad (2.26)$$

where

$$L_n = \frac{\kappa + \sqrt{1 + \gamma_n(1 - \kappa)}}{1 - \kappa}, \quad \{\gamma_n\} \text{ is as in (1.3)}. \quad (2.27)$$

By (2.18), (2.24), and (2.26), we know that (2.25) holds.

Step 5. Finally, by Lemma 2.3 and (2.25), we have $\omega_w(x_n) \subset \text{Fix}(T)$. Furthermore, it follows from (2.16) and Lemma 2.2 that the sequence $\{x_n\}$ converges strongly to $q = P_{\text{Fix}(T)}x_0$. \square

Remark 2.5. Theorem 2.4 improves [5, Theorem 4.1] since the condition that $\theta_n \rightarrow 0$ is satisfied and the boundedness of $\text{Fix}(T)$ is dropped off.

Theorem 2.6. *Let C be a closed convex subset of a Hilbert space H , $T : C \rightarrow C$ an asymptotically κ -strict pseudocontraction for some $0 \leq \kappa < 1$, and $\text{Fix}(T)$ be nonempty and bounded. Let $\{x_n\}$ the sequence generated by the following CQ-type algorithm with variable coefficients:*

$$\begin{aligned} x_0 &\in C \text{ chosen arbitrarily,} \\ y_n &= (1 - \hat{\beta}_n)x_n + \hat{\beta}_n T^n x_n, \\ C_n &= \left\{ z \in C : \|y_n - z\|^2 \leq \|x_n - z\|^2 + \hat{\beta}_n (\kappa + \hat{\beta}_n - 1) \|x_n - T^n x_n\|^2 + \theta_n \right\}, \\ Q_n &= \{z \in C : \langle x_n - z, x_0 - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n} x_0, \quad n \in \mathbb{N}, \end{aligned} \quad (2.28)$$

where

$$\hat{\beta}_n = \frac{\beta_n}{1 + \|x_n - x_0\|^2}, \quad \beta_n \in \left[\frac{1}{2}, 1 \right], \quad \theta_n = \left\{ \sup_{z \in \text{Fix}(T)} \|x_n - z\| \right\}^2 \hat{\beta}_n \gamma_n, \quad (2.29)$$

the sequence $\{\beta_n\}$ is chosen so that $\beta_n \rightarrow 1$ ($n \rightarrow \infty$), and $\{\gamma_n\}$ is as in (1.3). Then $\{x_n\}$ converges strongly to $P_{\text{Fix}(T)}x_0$.

Proof. It is easy to see that $\theta_n \rightarrow 0$ in Theorem 2.6. Following the reasoning in the proof of Theorem 2.4 and using $\text{Fix}(T)$ instead of $B_{r_0}(x_0) \cap \text{Fix}(T)$, we deduce the conclusion of Theorem 2.6. \square

Acknowledgments

The authors are very grateful to the referee for his/her valuable suggestions and comments. The work was supported partly by the NSF of China (10771202), the Research Fund for Shanghai Key Laboratory for Contemporary Applied Mathematics (08DZ2271900), and the Specialized Research Fund for the Doctoral Program of Higher Education of China (2007035805). This work is dedicated to W. Takahashi.

References

- [1] K. Goebel and W. A. Kirk, "A fixed point theorem for asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 35, no. 1, pp. 171–174, 1972.
- [2] K. Nakajo and W. Takahashi, "Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups," *Journal of Mathematical Analysis and Applications*, vol. 279, no. 2, pp. 372–379, 2003.
- [3] C. Martinez-Yanes and H.-K. Xu, "Strong convergence of the CQ method for fixed point iteration processes," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 11, pp. 2400–2411, 2006.
- [4] C. S. Ge and J. Liang, "Convergence theorems of new Ishikawa iterative procedures with errors for multi-valued Φ -hemicontractive mappings," *Communications in Mathematical Analysis*, vol. 7, no. 1, pp. 12–20, 2009.
- [5] T.-H. Kim and H.-K. Xu, "Convergence of the modified Mann's iteration method for asymptotically strict pseudocontractions," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 68, no. 9, pp. 2828–2836, 2008.
- [6] G. Marino and H.-K. Xu, "Weak and strong convergence theorems for strict pseudocontractions in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 336–346, 2007.
- [7] M. O. Osilike, A. Udomene, D. I. Igbokwe, and B. G. Akuchu, "Demiclosedness principle and convergence theorems for K -strictly asymptotically pseudocontractive maps," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 1334–1345, 2007.
- [8] X. L. Qin, Y. J. Cho, S. M. Kang, and M. J. Shang, "A hybrid iterative scheme for asymptotically K -strict pseudocontractions in Hilbert spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 5, pp. 1902–1911, 2009.
- [9] W. Takahashi, Y. Takeuchi, and R. Kubota, "Strong convergence theorems by hybrid methods for families of nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 276–286, 2008.
- [10] W. Takahashi and K. Zembayashi, "Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 1, pp. 45–57, 2009.
- [11] H. Zegeye and N. Shahzad, "Strong convergence theorems for a finite family of asymptotically nonexpansive mappings and semigroups," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 12, pp. 4496–4503, 2008.