

## ON $n$ -NORMED SPACES

HENDRA GUNAWAN and M. MASHADI

(Received 6 August 2000 and in revised form 12 October 2000)

**ABSTRACT.** Given an  $n$ -normed space with  $n \geq 2$ , we offer a simple way to derive an  $(n-1)$ -norm from the  $n$ -norm and realize that any  $n$ -normed space is an  $(n-1)$ -normed space. We also show that, in certain cases, the  $(n-1)$ -norm can be derived from the  $n$ -norm in such a way that the convergence and completeness in the  $n$ -norm is equivalent to those in the derived  $(n-1)$ -norm. Using this fact, we prove a fixed point theorem for some  $n$ -Banach spaces.

2000 Mathematics Subject Classification. 46B20, 46B99, 46A19, 46A99, 47H10.

**1. Introduction.** Let  $n \in \mathbb{N}$  and  $X$  be a real vector space of dimension  $d \geq n$ . (Here we allow  $d$  to be infinite.) A real-valued function  $\|\cdot, \dots, \cdot\|$  on  $X^n$  satisfying the following four properties

- (1)  $\|x_1, \dots, x_n\| = 0$  if and only if  $x_1, \dots, x_n$  are linearly dependent;
- (2)  $\|x_1, \dots, x_n\|$  is invariant under permutation;
- (3)  $\|x_1, \dots, x_{n-1}, \alpha x_n\| = |\alpha| \|x_1, \dots, x_{n-1}, x_n\|$  for any  $\alpha \in \mathbb{R}$ ;
- (4)  $\|x_1, \dots, x_{n-1}, y + z\| \leq \|x_1, \dots, x_{n-1}, y\| + \|x_1, \dots, x_{n-1}, z\|$ ,

is called an  $n$ -norm on  $X$  and the pair  $(X, \|\cdot, \dots, \cdot\|)$  is called an  $n$ -normed space.

A trivial example of an  $n$ -normed space is  $X = \mathbb{R}^n$  equipped with the following  $n$ -norm:

$$\|x_1, \dots, x_n\|_E := \text{abs} \left( \begin{vmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & \ddots & \vdots \\ x_{n1} & \cdots & x_{nn} \end{vmatrix} \right), \quad (1.1)$$

where  $x_i = (x_{i1}, \dots, x_{in}) \in \mathbb{R}^n$  for each  $i = 1, \dots, n$ . (The subscript  $E$  is for Euclidean.)

Note that in an  $n$ -normed space  $(X, \|\cdot, \dots, \cdot\|)$ , we have, for instance,  $\|x_1, \dots, x_n\| \geq 0$  and  $\|x_1, \dots, x_{n-1}, x_n\| = \|x_1, \dots, x_{n-1}, x_n + \alpha_1 x_1 + \cdots + \alpha_{n-1} x_{n-1}\|$  for all  $x_1, \dots, x_n \in X$  and  $\alpha_1, \dots, \alpha_{n-1} \in \mathbb{R}$ .

The theory of 2-normed spaces was first developed by Gähler [3] in the mid 1960's, while that of  $n$ -normed spaces can be found in [11]. Recent results can be found, for example, in [9, 10]. Related works on  $n$ -metric spaces and  $n$ -inner product spaces may be found, for example, in [1, 2, 4, 5, 7, 6, 12].

In this note, we will show that every  $n$ -normed space with  $n \geq 2$  is an  $(n-1)$ -normed space and hence, by induction, an  $(n-r)$ -normed space for all  $r = 1, \dots, n-1$ . In particular, given an  $n$ -normed space, we offer a simple way to derive an  $(n-1)$ -norm from the  $n$ -norm, different from that in [5].

We will also apply our result to study convergence and completeness in  $n$ -normed spaces, which will be defined later. This enables us to prove a fixed point theorem for some  $n$ -normed spaces.

The case  $n = 2$  was previously studied in [8].

**2. Preliminary results.** Suppose hereafter that  $n \geq 2$  and  $(X, \|\cdot, \dots, \cdot\|)$  is an  $n$ -normed space of dimension  $d \geq n$ . Take a linearly independent set  $\{a_1, \dots, a_n\}$  in  $X$ . With respect to  $\{a_1, \dots, a_n\}$ , define the following function  $\|\cdot, \dots, \cdot\|_\infty$  on  $X^{n-1}$  by

$$\|x_1, \dots, x_{n-1}\|_\infty := \max \{\|x_1, \dots, x_{n-1}, a_i\| : i = 1, \dots, n\}. \tag{2.1}$$

Then we have the following result.

**THEOREM 2.1.** *The function  $\|\cdot, \dots, \cdot\|_\infty$  defines an  $(n - 1)$ -norm on  $X$ .*

**PROOF.** We will verify that  $\|\cdot, \dots, \cdot\|_\infty$  satisfies the four properties of an  $(n - 1)$ -norm.

(1) If  $x_1, \dots, x_{n-1}$  are linearly dependent, then  $\|x_1, \dots, x_{n-1}\| = 0$  for each  $i = 1, \dots, n$ , and hence  $\|x_1, \dots, x_{n-1}\|_\infty = 0$ . Conversely, if  $\|x_1, \dots, x_{n-1}\|_\infty = 0$ , then  $\|x_1, \dots, x_{n-1}, a_i\| = 0$  and accordingly  $x_1, \dots, x_{n-1}, a_i$  are linearly dependent for each  $i = 1, \dots, n$ . But this can only happen when  $x_1, \dots, x_{n-1}$  are linearly dependent.

(2) Since  $\|x_1, \dots, x_{n-1}, a_i\|$  is invariant under any permutation of  $\{x_1, \dots, x_{n-1}\}$ , we find that  $\|x_1, \dots, x_{n-1}\|_\infty$  is also invariant under any permutation.

(3) Observe that

$$\begin{aligned} \|x_1, \dots, x_{n-2}, \alpha x_{n-1}\|_\infty &= \max \{\|x_1, \dots, x_{n-2}, \alpha x_{n-1}, a_i\| : i = 1, \dots, n\} \\ &= |\alpha| \max \{\|x_1, \dots, x_{n-2}, x_{n-1}, a_i\| : i = 1, \dots, n\} \\ &= |\alpha| \|x_1, \dots, x_{n-2}, x_{n-1}\|_\infty. \end{aligned} \tag{2.2}$$

(4) Observe that

$$\begin{aligned} \|x_1, \dots, x_{n-2}, y + z\|_\infty &= \max \{\|x_1, \dots, x_{n-2}, y + z, a_i\| : i = 1, \dots, n\} \\ &\leq \max \{\|x_1, \dots, x_{n-2}, y, a_i\| : i = 1, \dots, n\} \\ &\quad + \max \{\|x_1, \dots, x_{n-2}, z, a_i\| : i = 1, \dots, n\} \\ &= \|x_1, \dots, x_{n-2}, y\|_\infty + \|x_1, \dots, x_{n-2}, z\|_\infty. \end{aligned} \tag{2.3}$$

Therefore  $\|\cdot, \dots, \cdot\|_\infty$  defines an  $(n - 1)$ -norm on  $X$ . □

**COROLLARY 2.2.** *Every  $n$ -normed space is an  $(n - r)$ -normed space for all  $r = 1, \dots, n - 1$ . In particular, every  $n$ -normed space is a normed space.*

**REMARK 2.3.** Note that in general the function  $\|x_1, \dots, x_{n-1}\|_p := \{\sum_{i=1}^n \|x_1, \dots, x_{n-1}, a_i\|^p\}^{1/p}$ , where  $1 \leq p \leq \infty$ , also defines an  $(n - 1)$ -norm on  $X$ . These  $(n - 1)$ -norms, however, are equivalent to  $\|\cdot, \dots, \cdot\|_\infty$ , as long as we use the same set of  $n$  vectors  $a_1, \dots, a_n$ . In certain cases, it is possible to get equivalent  $(n - 1)$ -norms even if we use different sets of  $n$  vectors.

**2.1. The standard case.** Take a look at a standard example. Let  $X$  be a real inner product space of dimension  $d \geq n$ . Equip  $X$  with the standard  $n$ -norm

$$\|x_1, \dots, x_n\|_S := \left| \begin{array}{ccc} \langle x_1, x_1 \rangle & \cdots & \langle x_1, x_n \rangle \\ \vdots & \ddots & \vdots \\ \langle x_n, x_1 \rangle & \cdots & \langle x_n, x_n \rangle \end{array} \right|^{1/2}, \tag{2.4}$$

where  $\langle \cdot, \cdot \rangle$  denotes the inner product on  $X$ . (If  $X = \mathbb{R}^n$ , then this  $n$ -norm is exactly the same as the Euclidean  $n$ -norm  $\|\cdot, \dots, \cdot\|_E$  mentioned earlier.)

Notice that for  $n = 1$ , the above  $n$ -norm is the usual norm  $\|x_1\|_S = \langle x_1, x_1 \rangle^{1/2}$ , which gives the length of  $x_1$ , while for  $n = 2$ , it defines the standard 2-norm  $\|x_1, x_2\|_S = \{\|x_1\|_S^2 \|x_2\|_S^2 - \langle x_1, x_2 \rangle^2\}^{1/2}$ , which represents the area of the parallelogram spanned by  $x_1$  and  $x_2$ . Further, if  $X = \mathbb{R}^3$ , then  $\|x_1, x_2, x_3\|_S = \|x_1, x_2, x_3\|_E$  is nothing but the volume of the parallelepipeds spanned by  $x_1, x_2$ , and  $x_3$ . In general,  $\|x_1, \dots, x_n\|_S$  represents the volume of the  $n$ -dimensional parallelepiped spanned by  $x_1, \dots, x_n$  in  $X$ .

Now let  $\{e_1, \dots, e_n\}$  be an orthonormal set in  $X$ . Then, by [Theorem 2.1](#), the following function

$$\|x_1, \dots, x_{n-1}\|_\infty := \max \{\|x_1, \dots, x_{n-1}, e_i\|_S : i = 1, \dots, n\} \tag{2.5}$$

defines an  $(n - 1)$ -norm on  $X$ . Further, we have the following fact.

**FACT 2.4.** *On a standard  $n$ -normed space  $X$ , the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$ , defined with respect to  $\{e_1, \dots, e_n\}$ , is equivalent to the standard  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_S$ . Precisely, we have*

$$\|x_1, \dots, x_{n-1}\|_\infty \leq \|x_1, \dots, x_{n-1}\|_S \leq \sqrt{n} \|x_1, \dots, x_{n-1}\|_\infty \tag{2.6}$$

for all  $x_1, \dots, x_{n-1} \in X$ .

**PROOF.** Assume that  $x_1, \dots, x_{n-1}$  are linearly independent. For each  $i = 1, \dots, n$ , write  $e_i = e_i^\circ + e_i^\perp$  where  $e_i^\circ \in \text{span}\{x_1, \dots, x_{n-1}\}$  and  $e_i^\perp \perp \text{span}\{x_1, \dots, x_{n-1}\}$ . Then we have

$$\begin{aligned} \|x_1, \dots, x_{n-1}, e_i\|_S &= \|x_1, \dots, x_{n-1}, e_i^\perp\|_S \\ &= \left| \begin{array}{cccc} \langle x_1, x_1 \rangle & \cdots & \langle x_1, x_{n-1} \rangle & 0 \\ \vdots & \ddots & \vdots & \vdots \\ \langle x_{n-1}, x_1 \rangle & \cdots & \langle x_{n-1}, x_{n-1} \rangle & 0 \\ 0 & \cdots & 0 & \langle e_i^\perp, e_i^\perp \rangle \end{array} \right|^{1/2} \\ &\leq \left| \begin{array}{ccc} \langle x_1, x_1 \rangle & \cdots & \langle x_1, x_{n-1} \rangle \\ \vdots & \ddots & \vdots \\ \langle x_{n-1}, x_1 \rangle & \cdots & \langle x_{n-1}, x_{n-1} \rangle \end{array} \right|^{1/2} \\ &= \|x_1, \dots, x_{n-1}\|_S. \end{aligned} \tag{2.7}$$

Hence we get  $\|x_1, \dots, x_{n-1}\|_\infty \leq \|x_1, \dots, x_{n-1}\|_S$ .

Next, take a unit vector  $e = \alpha_1 e_1 + \dots + \alpha_n e_n$  such that  $e \perp \text{span}\{x_1, \dots, x_{n-1}\}$ . (Here we are still assuming that  $x_1, \dots, x_{n-1}$  are linearly independent.) Then, by properties (3) and (4) of the  $n$ -norm, we have

$$\begin{aligned} \|x_1, \dots, x_{n-1}\|_S &= \|x_1, \dots, x_{n-1}, e\|_S \\ &\leq |\alpha_1| \|x_1, \dots, x_{n-1}, e_1\|_S + \dots + |\alpha_n| \|x_1, \dots, x_{n-1}, e_n\|_S \\ &\leq (|\alpha_1| + \dots + |\alpha_n|) \|x_1, \dots, x_{n-1}\|_\infty. \end{aligned} \tag{2.8}$$

But, by the Cauchy-Schwarz inequality, we have

$$\sum_{i=1}^n |\alpha_i| \leq \left\{ \sum_{i=1}^n 1^2 \right\}^{1/2} \left\{ \sum_{i=1}^n |\alpha_i|^2 \right\}^{1/2} = \sqrt{n}. \tag{2.9}$$

Hence we obtain

$$\|x_1, \dots, x_{n-1}\|_S \leq \sqrt{n} \|x_1, \dots, x_{n-1}\|_\infty, \tag{2.10}$$

and this completes the proof. □

**2.2. The finite-dimensional case.** For finite-dimensional  $n$ -normed space  $(X, \|\cdot, \dots, \cdot\|)$ , we can in general derive an  $(n - 1)$ -norm from the  $n$ -norm in the following way. Take a linearly independent set  $\{a_1, \dots, a_m\}$  in  $X$ , with  $n \leq m \leq d$ . With respect to  $\{a_1, \dots, a_m\}$ , define the following function  $\|\cdot, \dots, \cdot\|_\infty$  on  $X^{n-1}$  by

$$\|x_1, \dots, x_{n-1}\|_\infty := \max \{ \|x_1, \dots, x_{n-1}, a_i\| : i = 1, \dots, m \}. \tag{2.11}$$

Then, as in [Theorem 2.1](#), the function  $\|\cdot, \dots, \cdot\|_\infty$  defines an  $(n - 1)$ -norm on  $X$ .

As we will see later, we can obtain a better  $(n - 1)$ -norm by using a set of  $d$ , rather than just  $n$ , linearly independent vectors in  $X$  (that is, by using a basis for  $X$ ).

**3. Applications and further results.** Recall that a sequence  $x(k)$  in an  $n$ -normed space  $(X, \|\cdot, \dots, \cdot\|)$  is said to *converge* to an  $x \in X$  (in the  $n$ -norm) whenever

$$\lim_{k \rightarrow \infty} \|x_1, \dots, x_{n-1}, x(k) - x\| = 0 \tag{3.1}$$

for every  $x_1, \dots, x_{n-1} \in X$ .

The following proposition says that the convergence in the  $n$ -norm implies the convergence in the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$ , defined with respect to an arbitrary linearly independent set  $\{a_1, \dots, a_n\}$  in  $X$ .

**PROPOSITION 3.1.** *If  $x(k)$  converges to an  $x \in X$  in the  $n$ -norm, then  $x(k)$  also converges to  $x$  in the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$ , that is,*

$$\lim_{k \rightarrow \infty} \|x_1, \dots, x_{n-2}, x(k) - x\|_\infty = 0 \tag{3.2}$$

for every  $x_1, \dots, x_{n-2} \in X$ .

**PROOF.** If  $x(k)$  converges to an  $x \in X$  in the  $n$ -norm, then

$$\lim_{k \rightarrow \infty} \|x_1, \dots, x_{n-2}, x(k) - x, a_i\| = 0 \tag{3.3}$$

for every  $x_1, \dots, x_{n-2} \in X$  and  $i = 1, \dots, n$ , and hence

$$\lim_{k \rightarrow \infty} \|x_1, \dots, x_{n-2}, x(k) - x\|_\infty = 0 \tag{3.4}$$

for every  $x_1, \dots, x_{n-2} \in X$ , that is,  $x(k)$  converges to  $x$  in the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$ .  $\square$

**3.1. The standard case.** In a standard  $n$ -normed space  $(X, \|\cdot, \dots, \cdot\|_S)$ , the converse of Proposition 3.1 is also true, especially when the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$  is defined with respect to an orthonormal set  $\{e_1, \dots, e_n\}$  in  $X$  as in Section 2.1.

**FACT 3.2.** *A sequence in a standard  $n$ -normed space  $X$  is convergent in the  $n$ -norm if and only if it is convergent in the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$ .*

**PROOF.** Suppose that  $x(k)$  converges to an  $x \in X$  in the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\infty$ . We want to show that  $x(k)$  also converges to  $x$  in the  $n$ -norm. Take  $x_1, \dots, x_{n-1} \in X$ . Then one may observe that

$$\|x_1, \dots, x_{n-2}, x_{n-1}, x(k) - x\|_S \leq \|x_1, \dots, x_{n-2}, x(k) - x\|_S \|x_{n-1}\|_S, \tag{3.5}$$

where  $\|\cdot, \dots, \cdot\|_S$  and  $\|\cdot\|_S$  on the right-hand side denote the standard  $(n - 1)$ -norm and the usual norm on  $X$ , respectively. By Fact 2.4, we have

$$\|x_1, \dots, x_{n-2}, x_{n-1}, x(k) - x\|_S \leq \sqrt{n} \|x_1, \dots, x_{n-2}, x(k) - x\|_\infty \|x_{n-1}\|_S. \tag{3.6}$$

But  $\lim_{k \rightarrow \infty} \|x_1, \dots, x_{n-2}, x(k) - x\|_\infty = 0$ , and so we conclude that

$$\lim_{k \rightarrow \infty} \|x_1, \dots, x_{n-1}, x(k) - x\|_S = 0, \tag{3.7}$$

that is,  $x(k)$  converges to  $x$  in the  $n$ -norm.  $\square$

**COROLLARY 3.3.** *A sequence in a standard  $n$ -normed space is convergent in the  $n$ -norm if and only if it is convergent in the standard  $(n - 1)$ -norm and, by induction, in the standard  $(n - r)$ -norm for all  $r = 1, \dots, n - 1$ . In particular, a sequence in a standard  $n$ -normed space is convergent in the  $n$ -norm if and only if it is convergent in the usual norm  $\|\cdot\|_S := \langle \cdot, \cdot \rangle^{1/2}$ .*

**3.2. The finite-dimensional case.** We also have a similar result for finite-dimensional  $n$ -normed space  $(X, \|\cdot, \dots, \cdot\|)$ . Let  $\{b_1, \dots, b_d\}$  be a basis for  $X$ . With respect to  $\{b_1, \dots, b_d\}$ , define the following function  $\|\cdot, \dots, \cdot\|_\boxtimes$  on  $X^{n-1}$  by

$$\|x_1, \dots, x_{n-1}\|_\boxtimes := \max \{ \|x_1, \dots, x_{n-1}, b_i\| : i = 1, \dots, d \}. \tag{3.8}$$

Then, as mentioned before, the function  $\|\cdot, \dots, \cdot\|_\boxtimes$  defines an  $(n - 1)$ -norm on  $X$ .

With this derived  $(n - 1)$ -norm, we have the following result.

**PROPOSITION 3.4.** *A sequence in the finite-dimensional  $n$ -normed space  $X$  is convergent in the  $n$ -norm if and only if it is convergent in the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\boxtimes$ .*

**PROOF.** If a sequence in  $X$  is convergent in the  $n$ -norm, then it will certainly be convergent in the  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_{\boxtimes}$ . Conversely, suppose that  $x(k)$  converges to an  $x \in X$  in  $\|\cdot, \dots, \cdot\|_{\boxtimes}$ . Take  $x_1, \dots, x_{n-1} \in X$ . Writing  $x_{n-1} = \alpha_1 b_1 + \dots + \alpha_d b_d$ , we get

$$\begin{aligned} \|\|x_1, \dots, x_{n-2}, x_{n-1}, x(k) - x\| &\leq |\alpha_1| \|\|x_1, \dots, x_{n-2}, x(k) - x, b_1\| \\ &\quad + \dots + |\alpha_d| \|\|x_1, \dots, x_{n-2}, x(k) - x, b_d\| \\ &\leq (|\alpha_1| + \dots + |\alpha_d|) \|\|x_1, \dots, x_{n-2}, x(k) - x\|_{\boxtimes}. \end{aligned} \tag{3.9}$$

But  $\lim_{k \rightarrow \infty} \|\|x_1, \dots, x_{n-2}, x(k) - x\|_{\boxtimes} = 0$ , and so we obtain

$$\lim_{k \rightarrow \infty} \|\|x_1, \dots, x_{n-1}, x(k) - x\| = 0, \tag{3.10}$$

that is,  $x(k)$  converges to  $x$  in the  $n$ -norm. □

**3.3. The standard, separable case.** We go back to the standard case, where  $X$  is a real inner product space of dimension  $d \geq n$  equipped with the standard  $n$ -norm  $\|\cdot, \dots, \cdot\|_S$  as in Section 2.1. But suppose now that  $X$  is separable and that  $\{e_i : i \in I_d\}$ , where  $I_d := \{1, \dots, d\}$  (if  $d < \infty$ ) or  $\mathbb{N}$  (if  $d = \infty$ ), is an orthonormal basis for  $X$ . For every  $x_1, \dots, x_{n-1} \in X$  and every basis vector  $e_i$  ( $i \in I_d$ ), we have

$$\|\|x_1, \dots, x_{n-1}, e_i\|_S \leq \|\|x_1, \dots, x_{n-1}\|_S, \tag{3.11}$$

where  $\|\cdot, \dots, \cdot\|_S$  on the right-hand side denotes the standard  $(n - 1)$ -norm on  $X$ . Hence, with respect to  $\{e_i : i \in I_d\}$ , we may define the function  $\|\cdot, \dots, \cdot\|_{\boxtimes}$  on  $X^{n-1}$  by

$$\|\|x_1, \dots, x_{n-1}\|_{\boxtimes} := \sup \{\|\|x_1, \dots, x_{n-1}, e_i\|_S : i \in I_d\} \tag{3.12}$$

and check that it also defines an  $(n - 1)$ -norm on  $X$ . Moreover, we have the following relation between the two derived  $(n - 1)$ -norms  $\|\cdot, \dots, \cdot\|_{\boxtimes}$  and  $\|\cdot, \dots, \cdot\|_{\infty}$  (the latter being defined with respect to  $\{e_1, \dots, e_n\}$  only):

$$\|\|x_1, \dots, x_{n-1}\|_{\infty} \leq \|\|x_1, \dots, x_{n-1}\|_{\boxtimes} \leq \|\|x_1, \dots, x_{n-1}\|_S \leq \sqrt{n} \|\|x_1, \dots, x_{n-1}\|_{\infty} \tag{3.13}$$

for every  $x_1, \dots, x_{n-1} \in X$ . Hence we conclude the following fact.

**FACT 3.5.** *On a standard  $n$ -normed space  $X$ , the two derived  $(n - 1)$ -norms  $\|\cdot, \dots, \cdot\|_{\infty}$  and  $\|\cdot, \dots, \cdot\|_{\boxtimes}$  and the standard  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_S$  are equivalent. Accordingly, a sequence in a standard  $n$ -normed space  $X$  is convergent in the  $n$ -norm if and only if it is convergent in one of the three  $(n - 1)$ -norms.*

**3.4. Cauchy sequences, completeness and fixed point theorem.** Recall that a sequence  $x(k)$  in an  $n$ -normed space  $(X, \|\cdot, \dots, \cdot\|)$  is called *Cauchy* (with respect to the  $n$ -norm) if

$$\lim_{k, l \rightarrow \infty} \|\|x_1, \dots, x_{n-1}, x(k) - x(l)\| = 0 \tag{3.14}$$

for every  $x_1, \dots, x_{n-1} \in X$ . If every Cauchy sequence in  $X$  converges to an  $x \in X$ , then  $X$  is said to be *complete* (with respect to the  $n$ -norm). A complete  $n$ -normed space is then called an  *$n$ -Banach space*.

By replacing the phrases “ $x(k)$  converges to  $x$ ” with “ $x(k)$  is Cauchy” and “ $x(k) - x$ ” with “ $x(k) - x(l)$ ,” we see that the analogues of [Proposition 3.1](#), [Fact 3.2](#), [Corollary 3.3](#), [Proposition 3.4](#), and [Fact 3.5](#) hold for Cauchy sequences.

Hence, for the standard or finite-dimensional case, we have the following result.

**PROPOSITION 3.6.** (a) *A standard  $n$ -normed space is complete if and only if it is complete with respect to one of the three  $(n - 1)$ -norms  $\|\cdot, \dots, \cdot\|_\infty$ ,  $\|\cdot, \dots, \cdot\|_\boxtimes$ , or  $\|\cdot, \dots, \cdot\|_S$ . By induction, a standard  $n$ -normed space is complete if and only if it is complete with respect to the usual norm  $\|\cdot\|_S := \langle \cdot, \cdot \rangle^{1/2}$ .*

(b) *A finite-dimensional  $n$ -normed space is complete if and only if it is complete with respect to the derived  $(n - 1)$ -norm  $\|\cdot, \dots, \cdot\|_\boxtimes$ .*

Consequently, we have the following result.

**COROLLARY 3.7** (fixed point theorem). *Let  $(X, \|\cdot, \dots, \cdot\|)$  be a standard or finite-dimensional  $n$ -Banach space, and  $T$  a contractive mapping of  $X$  into itself, that is, there exists a constant  $C \in (0, 1)$  such that*

$$\|x_1, \dots, x_{n-1}, Ty - Tz\| \leq C \|x_1, \dots, x_{n-1}, y - z\| \tag{3.15}$$

for all  $x_1, \dots, x_{n-1}, y, z$  in  $X$ . Then  $T$  has a unique fixed point in  $X$ .

**PROOF.** First consider the case  $n = 2$  (see [8]). By [Proposition 3.6](#), we know that  $X$  is a Banach space with respect to the derived norm  $\|\cdot\|_\infty$  (for standard case) or  $\|\cdot\|_\boxtimes$  (for finite-dimensional case). Since the mapping  $T$  is also contractive with respect to  $\|\cdot\|_\infty$  or  $\|\cdot\|_\boxtimes$ , we conclude by the fixed point theorem for Banach spaces that  $T$  has a unique fixed point in  $X$ . For  $n > 2$ , the result follows by induction.  $\square$

**REMARK 3.8.** In the finite-dimensional case, it is actually enough to assume that  $X$  is an  $n$ -normed space because we know that all finite-dimensional normed spaces are complete and, by [Proposition 3.6\(b\)](#), so are all finite-dimensional  $n$ -normed spaces.

**4. Concluding remark.** We have shown that an  $n$ -normed space with  $n \geq 2$  is an  $(n - 1)$ -normed space and that, for the standard or finite-dimensional case, the  $(n - 1)$ -norm can be derived from the  $n$ -norm in such a way that the convergence and completeness in the  $n$ -norm is equivalent to those in the derived  $(n - 1)$ -norm.

Below is an example of a non-standard, infinite-dimensional 2-normed space for which we can derive a norm from the 2-norm such that the convergence and completeness in the 2-norm is equivalent to those in the derived norm.

Let  $X = l^\infty$ , the space of bounded sequences of real numbers. Equip  $X$  with the following 2-norm

$$\|x, y\| := \sup_{i \in \mathbb{N}} \sup_{j \in \mathbb{N}} |x_i y_j - x_j y_i|, \tag{4.1}$$

where  $x = (x_1, x_2, x_3, \dots)$  and  $y = (y_1, y_2, y_3, \dots)$ . Let  $a_1 = (1, 0, 0, \dots)$  and  $a_2 = (0, 1, 0, \dots)$ .

With respect to  $\{a_1, a_2\}$ , we derive the norm  $\|\cdot\|_\infty$  via

$$\|x\|_\infty := \max\{\|x, a_1\|, \|x, a_2\|\}. \quad (4.2)$$

But  $\|x, a_1\| = \sup_{i \in \mathbb{N} \setminus \{1\}} |x_i|$  and  $\|x, a_2\| = \sup_{i \in \mathbb{N} \setminus \{2\}} |x_i|$ , and so we obtain

$$\|x\|_\infty = \sup_{i \in \mathbb{N}} |x_i|, \quad (4.3)$$

the usual norm on  $l^\infty$ .

Now suppose that  $x(k)$  is a sequence in  $X$  that converges to  $x$  in the derived norm  $\|\cdot\|_\infty$ . For every  $y \in X$ , we have

$$\begin{aligned} \|x(k) - x, y\| &= \sup_{i \in \mathbb{N}} \sup_{j \in \mathbb{N}} |(x_i(k) - x_i)y_j - (x_j(k) - x_j)y_i| \\ &\leq \sup_{i \in \mathbb{N}} \sup_{j \in \mathbb{N}} |x_i(k) - x_i| |y_j| + |x_j(k) - x_j| |y_i| \\ &\leq 2\|x(k) - x\|_\infty \|y\|_\infty, \end{aligned} \quad (4.4)$$

whence  $\lim_{k \rightarrow \infty} \|x(k) - x, y\| = 0$ . Hence  $x(k)$  converges to  $x$  in the 2-norm  $\|\cdot, \cdot\|$ .

Thus, for this particular example, we see that the convergence in the 2-norm is equivalent to that in the derived norm. By similar arguments, we can also verify that the completeness in the 2-norm is equivalent to that in the derived norm.

For general non-standard, infinite-dimensional  $n$ -normed spaces, however, it is unknown whether we can always derive an  $(n-1)$ -norm from the  $n$ -norm such that the convergence and completeness in the  $n$ -norm is equivalent to those in the derived  $(n-1)$ -norm.

**ACKNOWLEDGEMENTS.** This paper was revised during Gunawan's visit to the School of Mathematics, UNSW, Sydney, in 2000/2001, under an Australia-Indonesia Merdeka Fellowship, funded by the Australian Government through the Department of Education, Training and Youth Affairs and promoted through Australia Education International.

Both authors would like to thank the anonymous referees for their useful comments and suggestions.

## REFERENCES

- [1] C. Diminnie, S. Gähler, and A. White, *2-inner product spaces*, Demonstratio Math. **6** (1973), 525–536. [MR 51#1352](#). [Zbl 296.46022](#).
- [2] ———, *2-inner product spaces. II*, Demonstratio Math. **10** (1977), no. 1, 169–188. [MR 56#3623](#). [Zbl 377.46011](#).
- [3] S. Gähler, *Lineare 2-normierte Räume*, Math. Nachr. **28** (1964), 1–43 (German). [MR 29#6276](#). [Zbl 142.39803](#).
- [4] ———, *Untersuchungen über verallgemeinerte  $m$ -metrische Räume. I*, Math. Nachr. **40** (1969), 165–189 (German). [MR 40#1989](#). [Zbl 182.56404](#).
- [5] ———, *Untersuchungen über verallgemeinerte  $m$ -metrische Räume. II*, Math. Nachr. **40** (1969), 229–264 (German). [MR 40#1989](#). [Zbl 182.56501](#).
- [6] ———, *Untersuchungen über verallgemeinerte  $m$ -metrische Räume. III*, Math. Nachr. **41** (1969), 23–36 (German). [MR 40#1989](#). [Zbl 182.56601](#).
- [7] H. Gunawan, *On  $n$ -inner products,  $n$ -norms, and the Cauchy-Schwarz inequality*, to appear in Sci. Math. Japon.

- [8] H. Gunawan and Mashadi, *On finite-dimensional 2-normed spaces*, to appear in *Soochow J. Math.*
- [9] S. S. Kim and Y. J. Cho, *Strict convexity in linear  $n$ -normed spaces*, *Demonstratio Math.* **29** (1996), no. 4, 739–744. [MR 98a:46011](#). [Zbl 894.46004](#).
- [10] R. Malčeski, *Strong  $n$ -convex  $n$ -normed spaces*, *Mat. Bilten* (1997), no. 21, 81–102. [MR 99m:46059](#).
- [11] A. Misiak,  *$n$ -inner product spaces*, *Math. Nachr.* **140** (1989), 299–319. [MR 91a:46021](#). [Zbl 673.46012](#).
- [12] ———, *Orthogonality and orthonormality in  $n$ -inner product spaces*, *Math. Nachr.* **143** (1989), 249–261. [MR 91a:46022](#). [Zbl 708.46025](#).

HENDRA GUNAWAN: DEPARTMENT OF MATHEMATICS, BANDUNG INSTITUTE OF TECHNOLOGY, BANDUNG 40132, INDONESIA

*E-mail address:* [hgunawan@dns.math.itb.ac.id](mailto:hgunawan@dns.math.itb.ac.id)

MASHADI: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF RIAU, PEKANBARU 28293, INDONESIA