

Research Article

Approximation of Fixed Points of Nonexpansive Mappings and Solutions of Variational Inequalities

C. E. Chidume,¹ C. O. Chidume,² and Bashir Ali³

¹The Abdus Salam International Centre for Theoretical Physics, 34014 Trieste, Italy

²Department of Mathematics and Statistics, College of Sciences and Mathematics, Auburn University, Auburn, AL 36849, USA

³Department of Mathematical Sciences, Bayero University, 3011 Kano, Nigeria

Correspondence should be addressed to C. E. Chidume, chidume@ictp.it

Received 3 July 2007; Accepted 17 October 2007

Recommended by Siegfried Carl

Let E be a real q -uniformly smooth Banach space with constant d_q , $q \geq 2$. Let $T : E \rightarrow E$ and $G : E \rightarrow E$ be a nonexpansive map and an η -strongly accretive map which is also κ -Lipschitzian, respectively. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ that satisfies the following condition: C1: $\lim \lambda_n = 0$ and $\sum \lambda_n = \infty$. For $\delta \in (0, (q\eta/d_q k^q)^{1/(q-1)})$ and $\sigma \in (0, 1)$, define a sequence $\{x_n\}$ iteratively in E by $x_0 \in E$, $x_{n+1} = T^{\lambda_{n+1}} x_n = (1 - \sigma)x_n + \sigma[Tx_n - \delta\lambda_{n+1}G(Tx_n)]$, $n \geq 0$. Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality problem $VI(G, K)$ (search for $x^* \in K$ such that $\langle Gx^*, j_q(y - x^*) \rangle \geq 0$ for all $y \in K$), where $K := \text{Fix}(T) = \{x \in E : Tx = x\} \neq \emptyset$. A convergence theorem related to finite family of nonexpansive maps is also proved.

Copyright © 2008 C. E. Chidume et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let E be a real-normed space and let E^* be its dual space. For some real number q ($1 < q < \infty$), the generalized duality mapping $J_q : E \rightarrow 2^{E^*}$ is defined by

$$J_q(x) = \{f^* \in E^* : \langle x, f^* \rangle = \|x\|^q, \|f^*\| = \|x\|^{q-1}\}, \quad (1.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the pairing between elements of E and elements of E^* .

Let K be a nonempty closed convex subset of E , and let $S : E \rightarrow E$ be a nonlinear operator. The variational inequality problem is formulated as follows. Find a point $x^* \in K$ such that

$$VI(S, K) : \langle Sx^*, j_q(y - x^*) \rangle \geq 0 \quad \forall y \in K. \quad (1.2)$$

If $E = H$, a real Hilbert space, the variational inequality problem reduces to the following. Find a point $x^* \in K$ such that

$$VI(S, K) : \langle Sx^*, y - x^* \rangle \geq 0 \quad \forall y \in K. \quad (1.3)$$

A mapping $G : D(G) \subset E \rightarrow E$ is said to be *accretive* if for all $x, y \in D(G)$, there exists $j_q(x - y) \in J_q(x - y)$ such that

$$\langle Gx - Gy, j_q(x - y) \rangle \geq 0, \quad (1.4)$$

where $D(G)$ denotes the domain of G . For some real number $\eta > 0$, G is called η -*strongly accretive* if for all $x, y \in D(G)$, there exists $j_q(x - y) \in J_q(x - y)$ such that

$$\langle Gx - Gy, j_q(x - y) \rangle \geq \eta \|x - y\|^q. \quad (1.5)$$

G is κ -*Lipschitzian* if for some $\kappa > 0$, $\|G(x) - G(y)\| \leq \kappa \|x - y\|$ for all $x, y \in D(G)$ and G is called *nonexpansive* if $\kappa = 1$.

In Hilbert spaces, accretive operators are called *monotone* where inequalities (1.4) and (1.5) hold with j_q replaced by the identity map of H .

It is known that if S is Lipschitz and *strongly accretive*, then $VI(S, K)$ has a unique solution. An important problem is how to find a solution of $VI(S, K)$ whenever it exists. Considerable efforts have been devoted to this problem (see, e.g., [1, 2] and the references contained therein).

It is known that in a real Hilbert space, the $VI(S, K)$ is equivalent to the following fixed-point equation:

$$x^* = P_K(x^* - \delta Sx^*), \quad (1.6)$$

where $\delta > 0$ is an arbitrary fixed constant and P_K is the *nearest point projection map* from H onto K , that is, $P_K x = y$, where $\|x - y\| = \inf_{u \in K} \|x - u\|$ for $x \in H$. Consequently, under appropriate conditions on S and δ , fixed-point methods can be used to find or approximate a solution of $VI(S, K)$. For instance, if S is strongly monotone and Lipschitz, then a mapping $G : H \rightarrow H$, defined by $Gx = P_K(x - \delta Sx)$, $x \in H$ with $\delta > 0$ sufficiently small, is a strict contraction. Hence, the *Picard iteration*, $x_0 \in H$, $x_{n+1} = Gx_n$, $n \geq 0$ of the classical Banach contraction mapping principle, converges to the unique solution of the $VI(K, S)$.

It has been observed that the projection operator P_K in the fixed-point formulation (1.6) may make the computation of the iterates difficult due to possible complexity of the convex set K . In order to reduce the possible difficulty with the use of P_K , Yamada [2] recently introduced a hybrid descent method for solving the $VI(K, S)$. Let $T : H \rightarrow H$ be a map and let $K := \{x \in H : Tx = x\} \neq \emptyset$. Let S be η -strongly monotone and κ -Lipschitz on H . Let $\delta \in (0, 2\eta/\kappa^2)$ be arbitrary but fixed real number and let a sequence $\{\lambda_n\}$ in $(0, 1)$ satisfy the following conditions:

$$C1: \lim \lambda_n = 0; \quad C2: \sum \lambda_n = \infty; \quad C3: \lim \frac{\lambda_n - \lambda_{n+1}}{\lambda_n^2} = 0. \quad (1.7)$$

Starting with an arbitrary initial guess $x_0 \in H$, let a sequence $\{x_n\}$ be generated by the following algorithm:

$$x_{n+1} = Tx_n - \lambda_{n+1} \delta S(Tx_n), \quad n \geq 0. \quad (1.8)$$

Then, Yamada [2] proved that $\{x_n\}$ converges strongly to the unique solution of $VI(K, S)$.

In the case that $K = \bigcap_{i=1}^r F(T_i) \neq \emptyset$, where $\{T_i\}_{i=1}^r$ is a finite family of nonexpansive mappings, Yamada [2] studied the following algorithm:

$$x_{n+1} = T_{[n+1]}x_{n+1} - \lambda_{n+1}\delta S(T_{[n+1]}x_n), \quad n \geq 0, \quad (1.9)$$

where $T_{[k]} = T_{k \bmod r}$ for $k \geq 1$, with the mod *function* taking values in the set $\{1, 2, \dots, r\}$, where the sequence $\{\lambda_n\}$ satisfies the conditions C1, C2, and C4: $\sum |\lambda_n - \lambda_{n+N}| < \infty$. Under these conditions, he proved the strong convergence of $\{x_n\}$ to the unique solution of the VI(K, S).

Recently, Xu and Kim [1] studied the convergence of the algorithms (1.8) and (1.9), still in the framework of Hilbert spaces, and proved strong convergence with condition C3 replaced by C5: $\lim((\lambda_n - \lambda_{n+1})/\lambda_{n+1}) = 0$ and with condition C4 replaced by C6: $\lim((\lambda_n - \lambda_{n+r})/\lambda_{n+r}) = 0$. These are improvements on the results of Yamada. In particular, the canonical choice $\lambda_n := 1/(n+1)$ is applicable in the results of Xu and Kim but is not in the result of Yamada [2]. For further recent results on the schemes (1.8) and (1.9), still in the framework of Hilbert spaces, the reader may consult Wang [3], Zeng and Yao [4], and the references contained in them.

Recently, the present authors [5] extended the results of Xu and Kim [1] to q -uniformly smooth Banach spaces, $q \geq 2$. In particular, they proved theorems which are applicable in L_p spaces, $2 \leq p < \infty$ under conditions C1, C2, and C5 or C6 as in the result of Xu and Kim.

It is our purpose in this paper to modify the schemes (1.8) and (1.9) and prove strong convergence theorems for the unique solution of the variational inequality VI(K, S). Furthermore, in the case $T_i : E \rightarrow E$, $i = 1, 2, \dots, r$, is a family of nonexpansive mappings with $K = \bigcap_{i=1}^r F(T_i) \neq \emptyset$, we prove a convergence theorem where condition C6 is replaced by $\lim_{n \rightarrow \infty} \|T_{n+1}x_n - T_nx_n\| = 0$. An example satisfying this condition is given see, for example, [6]. All our theorems are proved in q -uniformly smooth spaces, $q \geq 2$. In particular, our theorems are applicable in L_p spaces, $2 \leq p < \infty$.

2. Preliminaries

Let E be a real Banach space and let K be a nonempty, closed, and convex subset of E . Let P be a mapping of E onto K . Then, P is said to be *sunny* if $P(Px + t(x - Px)) = Px$ for all $x \in E$ and $t \geq 0$. A mapping P of E into E is said to be a *retraction* if $P^2 = P$. A subset K is said to be *sunny nonexpansive retract* of E if there exists a sunny nonexpansive retraction of E onto K . A retraction P is said to be *orthogonal* if for each x , $x - P(x)$ is normal to K in the sense of James [7].

It is well known (see [8]) that if E is uniformly smooth and there exists a nonexpansive retraction of E onto K , then there exists a nonexpansive projection of E onto K . If E is a real smooth Banach space, then P is an orthogonal retraction of E onto K if and only if $P(x) \in K$ and $\langle P(x) - x, j_q(P(x) - y) \rangle \leq 0$ for all $y \in K$. It is also known (see, e.g., [9]) that if K is a convex subset of a uniformly convex Banach space whose norm is uniformly Gâteaux differentiable and $T : K \rightarrow K$ is nonexpansive with $F(T) \neq \emptyset$, then $F(T)$ is a nonexpansive retract of K .

Let K be a nonempty closed convex and bounded subset of a Banach space E and let the diameter of K be defined by $d(K) := \sup\{\|x - y\| : x, y \in K\}$. For each $x \in K$, let $r(x, K) := \sup\{\|x - y\| : y \in K\}$ and let $r(K) := \inf\{r(x, K) : x \in K\}$ denote the Chebyshev radius of K relative to itself. The *normal structure coefficient* $N(E)$ of E (see, e.g., [10]) is defined by $N(E) := \inf\{d(K)/r(K) : K \text{ is a closed convex and bounded subset of } E \text{ with } d(K) > 0\}$. A space E such that $N(E) > 1$ is said to have uniform normal structure. It is known that all

uniformly convex and uniformly smooth Banach spaces have uniform normal structure (see, e.g., [11, 12]).

We will denote a Banach limit by μ . Recall that μ is an element of $(l^\infty)^*$ such that $\|\mu\| = 1$, $\liminf_{n \rightarrow \infty} a_n \leq \mu_n a_n \leq \limsup_{n \rightarrow \infty} a_n$ and $\mu_n a_n = \mu_{n+1} a_n$ for all $\{a_n\}_{n \geq 0} \in l^\infty$ (see, e.g., [11, 13]).

Let E be a normed space with $\dim E \geq 2$. The *modulus of smoothness* of E is the function $\rho_E : [0, \infty) \rightarrow [0, \infty)$ defined by

$$\rho_E(\tau) := \sup \left\{ \frac{\|x + y\| + \|x - y\|}{2} - 1 : \|x\| = 1; \|y\| = \tau \right\}. \quad (2.1)$$

The space E is called *uniformly smooth* if and only if $\lim_{t \rightarrow 0^+} (\rho_E(t)/t) = 0$. For some positive constant q , E is called *q -uniformly smooth* if there exists a constant $c > 0$ such that $\rho_E(t) \leq ct^q$, $t > 0$. It is known that

$$L_p \text{ or } (l_p) \text{ spaces are } \begin{cases} 2\text{-uniformly smooth} & \text{if } 2 \leq p < \infty, \\ p\text{-uniformly smooth} & \text{if } 1 < p \leq 2 \end{cases} \quad (2.2)$$

(see, e.g., [13]). It is well known that if E is smooth, then the duality mapping is singled-valued, and if E is uniformly smooth, then the duality mapping is norm-to-norm uniformly continuous on bounded subset of E .

We will make use of the following well-known results.

Lemma 2.1. *Let E be a real-normed linear space. Then, the following inequality holds:*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle \quad \forall x, y \in E, \quad \forall j(x + y) \in J(x + y). \quad (2.3)$$

In the sequel, we will also make use of the following lemmas.

Lemma 2.2 (see [14]). *Let $(a_0, a_1, \dots) \in l^\infty$ such that $\mu_n(a_n) \leq 0$ for all Banach limit μ and $\limsup_{n \rightarrow \infty} (a_{n+1} - a_n) \leq 0$. Then, $\limsup_{n \rightarrow \infty} a_n \leq 0$.*

Lemma 2.3 (see [15]). *Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space E and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf \beta_n \leq \limsup \beta_n < 1$. Suppose $x_{n+1} = \beta_n y_n + (1 - \beta_n)x_n$ for all integers $n \geq 0$ and $\limsup (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$. Then, $\lim \|y_n - x_n\| = 0$.*

Lemma 2.4 (see [16]). *Let $\{a_n\}$ be a sequence of nonnegative real numbers satisfying the following relation:*

$$a_{n+1} \leq (1 - \alpha_n)a_n + \alpha_n \sigma_n + \gamma_n, \quad n \geq 0, \quad (2.4)$$

where (i) $\{\alpha_n\} \subset [0, 1]$, $\sum \alpha_n = \infty$; (ii) $\limsup \sigma_n \leq 0$; (iii) $\gamma_n \geq 0$; ($n \geq 0$), $\sum \gamma_n < \infty$. Then, $a_n \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 2.5 (see [17]). *Let E be a real q -uniformly smooth Banach space for some $q > 1$, then there exists some positive constant d_q such that*

$$\|x + y\|^q \leq \|x\|^q + q\langle y, j_q(x) \rangle + d_q \|y\|^q \quad \forall x, y \in E, \quad j_q(x) \in J_q(x). \quad (2.5)$$

Lemma 2.6 (see [12, Theorem 1]). *Suppose E is a Banach space with uniformly normal structure, K is a nonempty bounded subset of E , and $T : K \rightarrow K$ is uniformly k -Lipschitzian mapping with $k < N(E)^{1/2}$. Suppose also that there exists a nonempty bounded closed convex subset C of K with the following property (P):*

$$x \in C \text{ implies } \omega_w(x) \subset C, \quad (\text{P})$$

where $\omega_w(x)$ is the ω -limi set of T at x , that is, the set

$$\left\{ y \in E : y = \text{weak-lim}_j T^{n_j} x \text{ for some } n_j \rightarrow \infty \right\}. \quad (2.6)$$

Then, T has a fixed point in C .

3. Main results

We first prove the following lemma which will be central in the sequel.

Lemma 3.1. *Let E be a real q -uniformly smooth Banach space with constant d_q , $q \geq 2$. Let $T : E \rightarrow E$ and $G : E \rightarrow E$ be a nonexpansive map and an η -strongly accretive map which is also κ -Lipschitzian, respectively. For $\delta \in (0, (q\eta/d_q\kappa^q)^{1/(q-1)})$, $\sigma \in (0, 1)$, and $\lambda \in (0, 2/p(p-1))$, define a map $T^\lambda : E \rightarrow E$ by $T^\lambda x = (1 - \sigma)x + \sigma[Tx - \lambda\delta G(Tx)]$, $x \in E$. Then, T^λ is a strict contraction. Furthermore,*

$$\|T^\lambda x - T^\lambda y\| \leq (1 - \lambda\alpha)\|x - y\|, \quad x, y \in E, \quad (3.1)$$

where $\alpha = q/2 - \sqrt{q^2/4 - \sigma\delta(q\eta - \delta^{q-1}d_q\kappa^q)} \in (0, 1)$.

Proof. For $x, y \in E$,

$$\begin{aligned} \|T^\lambda x - T^\lambda y\|^q &= \|(1 - \sigma)(x - y) + \sigma[Tx - Ty - \lambda\delta(G(Tx) - G(Ty))]\|^q \\ &\leq (1 - \sigma)\|x - y\|^q + \sigma[\|Tx - Ty\|^q - q\lambda\delta\langle G(Tx) - G(Ty), j_q(Tx - Ty) \rangle \\ &\quad + d_q\lambda^q\delta^q\|G(Tx) - G(Ty)\|^q] \\ &\leq (1 - \sigma)\|x - y\|^q + \sigma[\|Tx - Ty\|^q - q\lambda\delta\eta\|Tx - Ty\|^q + d_q\lambda^q\delta^q\kappa^q\|Tx - Ty\|^q] \\ &\leq [1 - \sigma\lambda\delta(q\eta - d_q\lambda^{q-1}\delta^{q-1}\kappa^q)]\|x - y\|^q \\ &\leq [1 - \sigma\lambda\delta(q\eta - d_q\delta^{q-1}\kappa^q)]\|x - y\|^q. \end{aligned} \quad (3.2)$$

Define

$$f(\lambda) := 1 - \sigma\lambda\delta(q\eta - d_q\delta^{q-1}\kappa^q) = (1 - \lambda\tau)^q \quad \text{for some } \tau \in (0, 1) \text{ say.} \quad (3.3)$$

Then, there exists $\xi \in (0, \lambda)$ such that

$$1 - \sigma\lambda\delta(q\eta - d_q\delta^{q-1}\kappa^q) = 1 - q\tau\lambda + \frac{1}{2}q(q-1)(1 - \xi\tau)^{q-2}\lambda^2\tau^2. \quad (3.4)$$

This implies that

$$1 - \sigma\lambda\delta(q\eta - d_q\delta^{q-1}\kappa^q) \leq 1 - q\tau\lambda + \frac{1}{2}q(q-1)\lambda^2\tau^2. \quad (3.5)$$

Then, we have $\tau \leq q/2 - \sqrt{q^2/4 - \sigma\delta(q\eta - d_q\delta^{q-1}\kappa^q)}$.

Set

$$\alpha := \frac{q}{2} - \sqrt{\frac{q^2}{4} - \sigma\delta(q\eta - d_q\delta^{q-1}\kappa^q)}, \quad (3.6)$$

and the proof is complete. \square

We note that in L_p spaces, $2 \leq p < \infty$, the following inequality holds (see, e.g., [13]). For each $x, y \in L_p$, $2 \leq p < \infty$,

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x) \rangle + (p-1)\|y\|^2. \quad (3.7)$$

Using this inequality and following the method of proof of Lemma 3.1, the following corollary is easily proved.

Corollary 3.2. *Let $E = L_p$, $2 \leq p < \infty$. Let $T : E \rightarrow E$, $G : E \rightarrow E$ be a nonexpansive map, an η -strongly monotone, and κ -Lipschitzian map, respectively. For $\lambda, \sigma \in (0, 1)$ and $\delta \in (0, 2\eta/(p-1)\kappa^2)$, define a map $T^\lambda : E \rightarrow E$ by $T^\lambda x = (1-\sigma)x + \sigma[Tx - \lambda\delta G(Tx)]$, $x \in E$. Then, T^λ is a contraction. In particular,*

$$\|T^\lambda x - T^\lambda y\| \leq (1-\lambda\alpha)\|x - y\|, \quad x, y \in H, \quad (3.8)$$

where $\alpha = 1 - \sqrt{1 - \sigma\delta(2\eta - (p-1)\delta\kappa^2)} \in (0, 1)$.

Corollary 3.3. *Let H be a real Hilbert space, $T : H \rightarrow H$, $G : H \rightarrow H$ a nonexpansive map and an η -strongly monotone map which is also κ -Lipschitzian, respectively. For $\lambda, \sigma \in (0, 1)$ and $\delta \in (0, 2\eta/\kappa^2)$, define a map $T^\lambda : H \rightarrow H$ by $T^\lambda x = (1-\sigma)x + \sigma[Tx - \lambda\delta G(Tx)]$, $x \in H$. Then, T^λ is a contraction. In particular,*

$$\|T^\lambda x - T^\lambda y\| \leq (1-\lambda\alpha)\|x - y\|, \quad x, y \in H, \quad (3.9)$$

where $\alpha = 1 - \sqrt{1 - \sigma\delta(2\eta - \delta\kappa^2)} \in (0, 1)$.

Proof. Set $p = 2$ in Corollary 3.2 and the result follows. \square

Corollary 3.3 is a result of Yamada [2] and is the main tool used in [1–4].

We now prove our main theorems.

Theorem 3.4. *Let E be a real q -uniformly smooth Banach space with constant d_q , $q \geq 2$. Let $T : E \rightarrow E$ and $G : E \rightarrow E$ be a nonexpansive map and an η -strongly accretive map which is also κ -Lipschitzian, respectively. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ satisfying*

$$\text{C1: } \lim \lambda_n = 0; \quad \text{C2: } \sum \lambda_n = \infty. \quad (3.10)$$

For $\delta \in (0, (q\eta/d_q\kappa^q)^{1/(q-1)})$ and $\sigma \in (0, 1)$, define a sequence $\{x_n\}$ iteratively in E by $x_0 \in E$,

$$x_{n+1} = T^{\lambda_{n+1}} x_n = (1-\sigma)x_n + \sigma[Tx_n - \delta\lambda_{n+1}G(Tx_n)], \quad n \geq 0. \quad (3.11)$$

Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality $\text{VI}(G, K)$.

Proof. Let $x^* \in K := \text{Fix}(T)$, then the sequence $\{x_n\}$ satisfies

$$\|x_n - x^*\| \leq \max \left\{ \|x_0 - x^*\|, \frac{\delta}{\alpha} \|G(x^*)\| \right\}, \quad n \geq 0. \quad (3.12)$$

It is obvious that this is true for $n = 0$. Assume that it is true for $n = k$ for some $k \in \mathbb{N}$.

From the recursion formula (3.11), we have

$$\begin{aligned} \|x_{k+1} - x^*\| &= \|T^{\lambda_{k+1}} x_k - x^*\| \\ &\leq \|T^{\lambda_{k+1}} x_k - T^{\lambda_{k+1}} x^*\| + \|T^{\lambda_{k+1}} x^* - x^*\| \\ &\leq (1 - \lambda_{k+1} \alpha) \|x_k - x^*\| + \lambda_{k+1} \delta \|G(x^*)\| \\ &\leq \max \left\{ \|x_0 - x^*\|, \frac{\delta}{\alpha} \|G(x^*)\| \right\}, \end{aligned} \quad (3.13)$$

and the claim follows by induction. Thus, the sequence $\{x_n\}$ is bounded and so are $\{Tx_n\}$ and $\{G(Tx_n)\}$.

Define two sequences $\{\beta_n\}$ and $\{y_n\}$ by $\beta_n := (1 - \sigma)\lambda_{n+1} + \sigma$ and $y_n := (x_{n+1} - x_n + \beta_n x_n) / \beta_n$. Then,

$$y_n = \frac{(1 - \sigma)\lambda_{n+1}x_n + \sigma[Tx_n - \lambda_{n+1}\delta G(Tx_n)]}{\beta_n}. \quad (3.14)$$

Observe that $\{y_n\}$ is bounded and that

$$\begin{aligned} &\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\| \\ &\leq \left| \frac{\sigma}{\beta_{n+1}} - 1 \right| \|x_{n+1} - x_n\| + \left| \frac{\sigma}{\beta_{n+1}} - \frac{\sigma}{\beta_n} \right| \|Tx_n\| + \frac{\lambda_{n+2}(1 - \sigma)}{\beta_{n+1}} \|x_{n+1} - x_n\| \\ &\quad + (1 - \sigma) \left| \frac{\lambda_{n+2}}{\beta_{n+1}} - \frac{\lambda_{n+1}}{\beta_n} \right| \|x_n\| + \frac{\lambda_{n+1}\sigma\delta}{\beta_n} \|G(Tx_n) - G(Tx_{n+1})\| + \sigma\delta \left| \frac{\lambda_{n+1}}{\beta_n} - \frac{\lambda_{n+2}}{\beta_{n+1}} \right| \|G(Tx_{n+1})\|. \end{aligned} \quad (3.15)$$

This implies that $\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$, and by Lemma 2.3,

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \quad (3.16)$$

Hence,

$$\|x_{n+1} - x_n\| = \beta_n \|y_n - x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.17)$$

From the recursion formula (3.11), we have that

$$\sigma \|x_{n+1} - Tx_n\| \leq (1 - \sigma) \|x_{n+1} - x_n\| + \lambda_{n+1} \sigma \delta \|G(Tx_n)\| \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (3.18)$$

which implies that

$$\|x_{n+1} - Tx_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.19)$$

From (3.17) and (3.19), we have

$$\|x_n - Tx_n\| \leq \|x_n - x_{n+1}\| + \|x_{n+1} - Tx_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.20)$$

We now prove that $\limsup_{n \rightarrow \infty} \langle -G(x^*), j(x_{n+1} - x^*) \rangle \leq 0$.

Define a map $\phi : E \rightarrow \mathbb{R}$ by

$$\phi(x) = \mu_n \|x_n - x\|^2 \quad \forall x \in E. \quad (3.21)$$

Then, $\phi(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$, ϕ is continuous and convex, so as E is reflexive, there exists $y^* \in E$ such that $\phi(y^*) = \min_{u \in E} \phi(u)$. Hence, the set

$$K^* := \left\{ x \in E : \phi(x) = \min_{u \in E} \phi(u) \right\} \neq \emptyset. \quad (3.22)$$

By Lemma 2.6, $K^* \cap K \neq \emptyset$. Without loss of generality, assume that $y^* = x^* \in K^* \cap K$. Let $t \in (0, 1)$. Then, it follows that $\phi(x^*) \leq \phi(x^* - tG(x^*))$ and using Lemma 2.1, we obtain that

$$\|x_n - x^* + tG(x^*)\|^2 \leq \|x_n - x^*\|^2 + 2t \langle G(x^*), j(x_n - x^* + tG(x^*)) \rangle \quad (3.23)$$

which implies that

$$\mu_n \langle -G(x^*), j(x_n - x^* + tG(x^*)) \rangle \leq 0. \quad (3.24)$$

Moreover,

$$\begin{aligned} \mu_n \langle -G(x^*), j(x_n - x^*) \rangle &= \mu_n \langle -G(x^*), j(x_n - x^*) - j(x_n - x^* + tG(x^*)) \rangle \\ &\quad + \mu_n \langle -G(x^*), j(x_n - x^* + tG(x^*)) \rangle \\ &\leq \mu_n \langle -G(x^*), j(x_n - x^*) - j(x_n - x^* + tG(x^*)) \rangle. \end{aligned} \quad (3.25)$$

Since j is norm-to-norm uniformly continuous on bounded subsets of E , we have that

$$\mu_n \langle -G(x^*), j(x_n - x^*) \rangle \leq 0. \quad (3.26)$$

Furthermore, since $\|x_{n+1} - x_n\| \rightarrow 0$, as $n \rightarrow \infty$, we also have

$$\limsup_{n \rightarrow \infty} (\langle -G(x^*), j(x_n - x^*) \rangle - \langle -G(x^*), j(x_{n+1} - x^*) \rangle) \leq 0, \quad (3.27)$$

and so we obtain by Lemma 2.2 that $\limsup_{n \rightarrow \infty} \langle -G(x^*), j(x_n - x^*) \rangle \leq 0$.

From the recursion formula (3.11) and Lemma 2.1, we have

$$\begin{aligned} \|x_{n+1} - x^*\|^2 &= \|T^{\lambda_{n+1}} x_n - T^{\lambda_{n+1}} x^* + T^{\lambda_{n+1}} x^* - x^*\|^2 \\ &\leq \|T^{\lambda_{n+1}} x_n - T^{\lambda_{n+1}} x^*\|^2 + 2\lambda_{n+1} \delta \langle -G(x^*), j(x_{n+1} - x^*) \rangle \\ &\leq (1 - \lambda_{n+1} \alpha) \|x_n - x^*\|^2 + 2\lambda_{n+1} \delta \langle -G(x^*), j(x_{n+1} - x^*) \rangle, \end{aligned} \quad (3.28)$$

and by Lemma 2.4, we have that $x_n \rightarrow x^*$ as $n \rightarrow \infty$. This completes the proof. \square

The following corollaries follow from Theorem 3.4.

Corollary 3.5. *Let $E = L_p$, $2 \leq p < \infty$. Let $T : E \rightarrow E$ and $G : E \rightarrow E$ be a nonexpansive map and an η -strongly accretive map which is also κ -Lipschitzian, respectively. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ that satisfies conditions C1 and C2 as in Theorem 3.4. For $\delta \in (0, 2\eta/(p-1)\kappa^2)$ and $\sigma \in (0, 1)$, define a sequence $\{x_n\}$ iteratively in E by (3.11). Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality $\text{VI}(G, K)$.*

Corollary 3.6. *Let $E = H$ be a real Hilbert space. Let $T : H \rightarrow H$ and $G : H \rightarrow H$ be a nonexpansive map and an η -strongly monotone map which is also κ -Lipschitzian, respectively. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ that satisfies conditions C1 and C2 as in Theorem 3.4. For $\delta \in (0, 2\eta/\kappa^2)$ and $\sigma \in (0, 1)$, define a sequence $\{x_n\}$ iteratively in H by (3.11). Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality $\text{VI}(G, K)$.*

Finally, we prove the following more general theorem.

Theorem 3.7. *Let E be a real q -uniformly smooth Banach space with constant d_q , $q \geq 2$. Let $T_i : E \rightarrow E$, $i = 1, 2, \dots, r$, be a finite family of nonexpansive mappings with $K := \bigcap_{i=1}^r \text{Fix}(T_i) \neq \emptyset$. Let $G : E \rightarrow E$ be an η -strongly accretive map which is also κ -Lipschitzian. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ satisfying*

$$\text{C1: } \lim \lambda_n = 0; \quad \text{C2: } \sum \lambda_n = \infty. \quad (3.29)$$

For a fixed real number $\delta \in (0, (q\eta/d_q\kappa^q)^{1/(q-1)})$, define a sequence $\{x_n\}$ iteratively in E by $x_0 \in E$:

$$x_{n+1} = T_{[n+1]}^{\lambda_{n+1}} x_n = (1 - \sigma)x_n + \sigma [T_{[n+1]} x_n - \delta \lambda_n G(T_{[n+1]} x_n)], \quad n \geq 0, \quad (3.30)$$

where $T_{[n]} = T_{n \bmod r}$. Assume also that

$$K = \text{Fix}(T_r T_{r-1} \cdots T_1) = \text{Fix}(T_1 T_r \cdots T_2) = \cdots = \text{Fix}(T_{r-1} T_{r-2} \cdots T_r) \quad (3.31)$$

and $\lim_{n \rightarrow \infty} \|T_{n+1} x_n - T_n x_n\| = 0$. Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality $\text{VI}(G, K)$.

Proof. Let $x^* \in K$, then the sequence $\{x_n\}$ satisfies that

$$\|x_n - x^*\| \leq \max \left\{ \|x_0 - x^*\|, \frac{\delta}{\alpha} \|G(x^*)\| \right\}, \quad n \geq 0. \quad (3.32)$$

It is obvious that this is true for $n = 0$. Assume it is true for $n = k$ for some $k \in \mathbb{N}$.

From the recursion formula (3.30), we have

$$\begin{aligned} \|x_{k+1} - x^*\| &= \|T_{[k+1]}^{\lambda_{k+1}} x_k - x^*\| \\ &\leq \|T_{[k+1]}^{\lambda_{k+1}} x_k - T_{[k+1]}^{\lambda_{k+1}} x^*\| + \|T_{[k+1]}^{\lambda_{k+1}} x^* - x^*\| \\ &\leq (1 - \lambda_{k+1} \alpha) \|x_k - x^*\| + \lambda_{k+1} \delta \|G(x^*)\| \\ &\leq \max \left\{ \|x_0 - x^*\|, \frac{\delta}{\alpha} \|G(x^*)\| \right\}, \end{aligned} \quad (3.33)$$

and the claim follows by induction. Thus, the sequence $\{x_n\}$ is bounded and so are $\{T_{[n]}x_n\}$ and $\{G(T_{[n]}x_n)\}$.

Define two sequences $\{\beta_n\}$ and $\{y_n\}$ by $\beta_n := (1-\sigma)\lambda_{n+1} + \sigma$ and $y_n := (x_{n+1} - x_n + \beta_n x_n) / \beta_n$. Then,

$$y_n = \frac{(1-\sigma)\lambda_{n+1}x_n + \sigma[T_{[n+1]}x_n - \lambda_{n+1}\delta G(T_{[n+1]}x_n)]}{\beta_n}. \quad (3.34)$$

Observe that $\{y_n\}$ is bounded and that

$$\begin{aligned} \|y_{n+1} - y_n\| - \|x_{n+1} - x_n\| &\leq \left| \frac{\sigma}{\beta_{n+1}} - 1 \right| \|x_{n+1} - x_n\| \\ &\quad + \frac{\sigma}{\beta_{n+1}} \|T_{[n+2]}x_n - T_{[n+1]}x_n\| + \left| \frac{\sigma}{\beta_{n+1}} - \frac{\sigma}{\beta_n} \right| \|T_{[n+1]}x_n\| \\ &\quad + \frac{\lambda_{n+2}(1-\sigma)}{\beta_{n+1}} \|x_{n+1} - x_n\| + (1-\sigma) \left| \frac{\lambda_{n+2}}{\beta_{n+1}} - \frac{\lambda_{n+1}}{\beta_n} \right| \|x_n\| \\ &\quad + \frac{\lambda_{n+1}\sigma\delta}{\beta_n} \|G(T_{[n+1]}x_n) - G(T_{[n+2]}x_{n+1})\| \\ &\quad + \sigma\delta \left| \frac{\lambda_{n+1}}{\beta_n} - \frac{\lambda_{n+2}}{\beta_{n+1}} \right| \|G(T_{[n+2]}x_{n+1})\|. \end{aligned} \quad (3.35)$$

This implies that $\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$, and by Lemma 2.3,

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0. \quad (3.36)$$

Hence,

$$\|x_{n+1} - x_n\| = \beta_n \|y_n - x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.37)$$

From the recursion formula (3.30), we have that

$$\sigma \|x_{n+1} - T_{[n+1]}x_n\| \leq (1-\sigma) \|x_{n+1} - x_n\| + \lambda_{n+1}\sigma\delta \|G(T_{[n+1]}x_n)\| \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (3.38)$$

which implies that

$$\|x_{n+1} - T_{[n+1]}x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.39)$$

From (3.37) and (3.39), we have

$$\|x_n - T_{[n+1]}x_n\| \leq \|x_n - x_{n+1}\| + \|x_{n+1} - T_{[n+1]}x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.40)$$

Also,

$$\|x_{n+r} - x_n\| \leq \|x_{n+r} - x_{n+r-1}\| + \|x_{n+r-1} - x_{n+r-2}\| + \cdots + \|x_{n+1} - x_n\|, \quad (3.41)$$

and so

$$\|x_{n+r} - x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.42)$$

Using the fact that T_i is nonexpansive for each i , we obtain the following finite table:

$$\begin{aligned}
x_{n+r} - T_{n+r}x_{n+r-1} &\longrightarrow 0 \quad \text{as } n \longrightarrow \infty; \\
T_{n+r}x_{n+r-1} - T_{n+r}T_{n+r-1}x_{n+r-2} &\longrightarrow 0 \quad \text{as } n \longrightarrow \infty; \\
&\vdots \\
T_{n+r}T_{n+r-1} \cdots T_{n+2}x_{n+1} - T_{n+r}T_{n+r-1} \cdots T_{n+2}T_{n+1}x_n &\longrightarrow 0 \quad \text{as } n \longrightarrow \infty;
\end{aligned} \tag{3.43}$$

and adding up the table yields

$$x_{n+r} - T_{n+r}T_{n+r-1} \cdots T_{n+1}x_n \longrightarrow 0 \quad \text{as } n \longrightarrow \infty. \tag{3.44}$$

Using this and (3.42), we get that $\lim_{n \rightarrow \infty} \|x_n - T_{n+r}T_{n+r-1} \cdots T_{n+1}x_n\| = 0$.

Carrying out similar arguments as in the proof of Theorem 3.4, we easily get that

$$\limsup_{n \rightarrow \infty} \langle -G(x^*), j(x_{n+1} - x^*) \rangle \leq 0. \tag{3.45}$$

From the recursion formula (3.30), and Lemma 2.1, we have

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \|T_{[n+1]}^{\lambda_{n+1}}x_n - T_{[n]}^{\lambda_{n+1}}x^* + T_{[n+1]}^{\lambda_{n+1}}x^* - x^*\|^2 \\
&\leq \|T_{[n+1]}^{\lambda_{n+1}}x_n - T_{[n+1]}^{\lambda_{n+1}}x^*\|^2 + 2\lambda_{n+1}\sigma\delta \langle -G(x^*), j(x_{n+1} - x^*) \rangle \\
&\leq (1 - \lambda_{n+1}\alpha) \|x_n - x^*\|^2 + 2\lambda_{n+1}\sigma\delta \langle -G(x^*), j(x_{n+1} - x^*) \rangle,
\end{aligned} \tag{3.46}$$

and by Lemma 2.4, we have that $x_n \rightarrow x^*$ as $n \rightarrow \infty$. This completes the proof. \square

The following corollaries follow from Theorem 3.7.

Corollary 3.8. *Let $E = L_p$, $2 \leq p < \infty$. Let $T_i : E \rightarrow E$, $i = 1, 2, \dots, r$, be a finite family of nonexpansive mappings with $K = \bigcap_{i=1}^r \text{Fix}(T_i) \neq \emptyset$. Let $G : E \rightarrow E$ be an η -strongly accretive map which is also κ -Lipschitzian. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ that satisfies conditions C1 and C2 as in Theorem 3.7 and also $\lim_{n \rightarrow \infty} \|T_{n+1}x_n - T_nx_n\| = 0$. For $\delta \in (0, 2\eta/(p-1)\kappa^2)$, define a sequence $\{x_n\}$ iteratively in E by (3.30). Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality VI(G, K).*

Corollary 3.9. *Let $E = H$ be a real Hilbert space. Let $T_i : H \rightarrow H$, $i = 1, 2, \dots, r$, be a finite family of nonexpansive mappings with $K = \bigcap_{i=1}^r \text{Fix}(T_i) \neq \emptyset$. Let $G : H \rightarrow H$ be an η -strongly monotone map which is also κ -Lipschitzian. Let $\{\lambda_n\}$ be a real sequence in $[0, 1]$ that satisfies conditions C1 and C2 as in Theorem 3.7 and also $\lim_{n \rightarrow \infty} \|T_{n+1}x_n - T_nx_n\| = 0$. For $\delta \in (0, 2\eta/\kappa^2)$, define a sequence $\{x_n\}$ iteratively in H by (3.30). Then, $\{x_n\}$ converges strongly to the unique solution x^* of the variational inequality VI(G, K).*

Remark 3.10. Observe that condition C6 in Theorem 3.2 of [1] is dropped in Corollary 3.9, being replaced by condition $\lim_{n \rightarrow \infty} \|T_{n+1}x_n - T_nx_n\| = 0$ on the mappings $\{T_i\}_{i=1}^r$.

Acknowledgment

This research is supported by the Japanese Mori Fellowship of UNESCO at The Abdus Salam International Center for Theoretical Physics (Trieste, Italy).

References

- [1] H. K. Xu and T. H. Kim, "Convergence of hybrid steepest-descent methods for variational inequalities," *Journal of Optimization Theory and Applications*, vol. 119, no. 1, pp. 185–201, 2003.
- [2] I. Yamada, "The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings," in *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications*, D. Butnariu, Y. Censor, and S. Reich, Eds., vol. 8, pp. 473–504, North-Holland, Amsterdam, The Netherlands, 2001.
- [3] L. Wang, "An iteration method for nonexpansive mappings in Hilbert spaces," *Fixed Point Theory and Applications*, vol. 2007, Article ID 28619, 8 pages, 2007.
- [4] L.-C. Zeng and J.-C. Yao, "Implicit iteration scheme with perturbed mapping for common fixed points of a finite family of nonexpansive mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 11, pp. 2507–2515, 2006.
- [5] C. E. Chidume, C. O. Chidume, and B. Ali, "Convergence of hybrid steepest descent method for variational inequalities in Banach spaces," to appear in *Proceedings of the American Mathematical Society*.
- [6] C. E. Chidume and B. Ali, "Convergence theorems for common fixed points for finite families of nonexpansive mappings in reflexive Banach spaces," to appear in *Nonlinear Analysis: Theory, Methods & Applications*.
- [7] R. C. James, "Orthogonality and linear functionals in normed linear spaces," *Transactions of the American Mathematical Society*, vol. 61, pp. 265–292, 1947.
- [8] R. E. Bruck Jr., "Nonexpansive projections on subsets of Banach spaces," *Pacific Journal of Mathematics*, vol. 47, pp. 341–355, 1973.
- [9] N. Shioji and W. Takahashi, "Strong convergence of averaged approximants for asymptotically nonexpansive mappings in Banach spaces," *Journal of Approximation Theory*, vol. 97, no. 1, pp. 53–64, 1999.
- [10] W. L. Bynum, "Normal structure coefficients for Banach spaces," *Pacific Journal of Mathematics*, vol. 86, no. 2, pp. 427–436, 1980.
- [11] C. E. Chidume, J. Li, and A. Udomene, "Convergence of paths and approximation of fixed points of asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 133, no. 2, pp. 473–480, 2005.
- [12] T.-C. Lim and H. K. Xu, "Fixed point theorems for asymptotically nonexpansive mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 22, no. 11, pp. 1345–1355, 1994.
- [13] C. E. Chidume, "Geometric properties of Banach spaces and nonlinear iterations," Research Monograph, International Centre for Theoretical Physics, Trieste, Italy, in print.
- [14] N. Shioji and W. Takahashi, "Strong convergence of approximated sequences for nonexpansive mappings in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 125, no. 12, pp. 3641–3645, 1997.
- [15] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [16] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.
- [17] H. K. Xu, "Inequalities in Banach spaces with applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 16, no. 12, pp. 1127–1138, 1991.