

STRONGLY ELLIPTIC OPERATORS FOR A PLANE WAVE DIFFRACTION PROBLEM IN BESSEL POTENTIAL SPACES

LUIS P. CASTRO

Department of Mathematics
University of Aveiro,
3810 - 193 Aveiro, Portugal.
EMail: lcastro@mat.ua.pt



volume 3, issue 2, article 25,
2002.

*Received 08 January, 2001;
accepted 23 January, 2002.*

Communicated by: S.S. Dragomir

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


We consider a plane wave diffraction problem by a union of several infinite strips. The problem is formulated as a boundary-transmission one for the Helmholtz equation in a Bessel potential space setting and where Neumann conditions are assumed on the strips. Using arguments of strong ellipticity and different kinds of operator relations between convolution type operators, it is shown the well-posedness of the problem in a smoothness neighborhood of the Bessel potential space with finite energy norm.

2000 Mathematics Subject Classification: 78A45, 35J25, 47B35, 47A50, 47A20, 47A68

Key words: Diffraction problem, Strong ellipticity, Convolution type operator, Wiener-Hopf operator, Equivalence after extension

Contents

1	Formulation of the Problem	3
2	The Problem from an Operator Point of View	5
3	The Extension to a Half-line Setting	8
4	Strong Ellipticity and Corresponding Well-posedness of the Problem	11
	References	



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

1. Formulation of the Problem

This paper deals with the problem of diffraction of an electromagnetic wave by a union of n infinite magnetic strips from an operator point of view.

We will present a formulation that results from the investigation of plane waves which propagate in a direction orthogonal to the edge $x = y = 0, z \in \mathbb{R}$. Thus, the problem will be posed as a boundary-transmission one for the two-dimensional Helmholtz equation where the dependence on one dimension is dropped already. Moreover, also due to perpendicular wave incidence, the union of n infinite strips will be represented by

$$\Omega =]\gamma_1, \gamma_2[\cup \dots \cup]\gamma_{2n-1}, \gamma_{2n}[,$$

with $0 = \gamma_1 < \dots < \gamma_{2n}$ and $n \in \mathbb{N}$.

We will use the Bessel potential spaces $H^\sigma(\mathbb{R})$, with $\sigma \in \mathbb{R}$, formed by the tempered distributions φ such that $\|\varphi\|_{H^\sigma(\mathbb{R})} = \|\mathcal{F}^{-1}(1 + \xi^2)^{\sigma/2} \cdot \mathcal{F}\varphi\|_{L^2(\mathbb{R})}$ is finite (here \mathcal{F} denotes the Fourier transformation). In addition, we denote by $\tilde{H}^\sigma(\Omega)$ [17, §2.10.3] the closed subspace of $H^\sigma(\mathbb{R})$ defined by the distributions with support contained in $\overline{\Omega}$ and $H^\sigma(\Omega)$ will denote the space of generalized functions on Ω which have extensions into \mathbb{R} that belong to $H^\sigma(\mathbb{R})$. The space $\tilde{H}^\sigma(\Omega)$ is endowed with the subspace topology, and on $H^\sigma(\Omega)$ we put the norm of the quotient space $H^\sigma(\mathbb{R})/\tilde{H}^\sigma(\mathbb{R} \setminus \overline{\Omega})$. In particular, we shall denote by $L^2(\mathbb{R}_+)$ and $L_+^2(\mathbb{R})$, the spaces $H^0(\mathbb{R}_+)$ and $\tilde{H}^0(\mathbb{R}_+)$, respectively. All those definitions can be extended to the multi-index case $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_m) \in \mathbb{R}^m$ by taking the product topology.

The problem is inspired by the classical Sommerfeld type problems considered, for instance in [7, 9, 10, 12, 13, 15, 16], for the half-line case instead of



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 3 of 20

the present Ω . In fact, we present here a generalization of the problem treated in [4] where a corresponding problem was taken into consideration for only one strip. Several changes take place here, in particular, we notice the necessity of different constructions of operator relations that can be found in the next sections.

More concretely, we are interested in studying well-posedness of the problem to find $u \in L^2(\mathbb{R}^2)$, with $u|_{\mathbb{R}^2_{\pm}} \in H^{\epsilon}(\mathbb{R}^2_{\pm})$, $\epsilon \in]1/2, 3/2[$, so that

$$(1.1) \quad (\Delta + k^2) u = 0 \quad \text{in} \quad \mathbb{R}^2_{\pm},$$


$$(1.2) \quad u_1^{\pm} = h \quad \text{on} \quad \Omega,$$

$$(1.3) \quad \begin{cases} u_0^+ - u_0^- = 0 \\ u_1^+ - u_1^- = 0 \end{cases} \quad \text{on} \quad \mathbb{R} \setminus \overline{\Omega},$$

where \mathbb{R}^2_{\pm} represents the upper/lower half-plane, $\Delta = \partial^2/\partial x^2 + \partial^2/\partial y^2$ stands for the Laplace operator, $k \in \mathbb{C}$ is the wave number, which, due to the assumption of a lossy medium, is assumed to fulfill

$$\Im mk > 0,$$

$u_0^{\pm} = u|_{y=\pm 0}$, $u_1^{\pm} = (\partial u/\partial y)|_{y=\pm 0}$ and the element $h \in H^{\epsilon-3/2}(\Omega)$ is arbitrarily given.



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 4 of 20

2. The Problem from an Operator Point of View

In order to study the existence and uniqueness of the solution of the problem, as well as continuous dependence on the data, we will construct several operators that are shown to be, in a sense, connected with the problem.

In the first stage, the problem can be described by the use of a linear operator

$$\mathcal{P} : D(\mathcal{P}) \rightarrow H^{\epsilon-3/2}(\Omega),$$

if we define $D(\mathcal{P})$ as the subspace of $H^\epsilon(\mathbb{R}_+^2) \times H^\epsilon(\mathbb{R}_-^2)$ whose functions fulfill the Helmholtz equation (1.1) and all the remaining homogeneous transmission conditions that appear from (1.2) – (1.3) whereas the action $\mathcal{P}u = h$ results from the non-homogeneous conditions (1.2).

In this sense, we will say that the operator \mathcal{P} is *associated* to the problem and our aim is to prove that \mathcal{P} is bounded and invertible for suitable orders of smoothness ϵ . This goal will be achieved by the construction of several operator relations that will allow us to understand better the structure of \mathcal{P} .

To this end, we begin by introducing some notation. Let $r_{\mathbb{R} \rightarrow \Omega} : H^\sigma(\mathbb{R}) \rightarrow H^\sigma(\Omega)$ be the restriction operator and let

$$t(\xi) = (\xi^2 - k^2)^{1/2}, \quad \xi \in \mathbb{R},$$

denote the branch of the square root that tends to $+\infty$ as $\xi \rightarrow +\infty$ with branch cuts along $\pm k \pm i\eta$, $\eta \geq 0$.

Theorem 2.1. *The operator \mathcal{P} is equivalent to the convolution type operator*

$$(2.1) \quad W_{t,\Omega} = r_{\mathbb{R} \rightarrow \Omega} \mathcal{F}^{-1} t \cdot \mathcal{F} : \tilde{H}^{\epsilon-1/2}(\Omega) \rightarrow H^{\epsilon-3/2}(\Omega),$$

i.e. there are bounded invertible linear operators E and F so that $\mathcal{P} = EW_{t,\Omega}F$.



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 5 of 20

Proof. Analogously to what happens in the half-line case [10], a function $u \in L^2(\mathbb{R}^2)$, with $u|_{\mathbb{R}^2_{\pm}} \in H^\epsilon(\mathbb{R}^2_{\pm})$, satisfies the Helmholtz equation (1.1) if and only if it is representable by

$$(2.2) \quad u(x, y) = \mathcal{F}_{\xi \mapsto x}^{-1} e^{-t(\xi)y} \mathcal{F}_{x \rightarrow \xi} u_0^+(x) \chi_+(y) + \mathcal{F}_{\xi \mapsto x}^{-1} e^{t(\xi)y} \mathcal{F}_{x \rightarrow \xi} u_0^-(x) \chi_-(y)$$

for $(x, y) \in \mathbb{R}^2$, where $\mathcal{F}_{x \rightarrow \xi} u(x, y) = \int_{\mathbb{R}} u(x, y) e^{i\xi x} dx$ and χ_+, χ_- denote the characteristic functions of the positive and negative half-line, respectively.

Let

$$Z = \left\{ (\phi, \psi) \in [H^{\epsilon-1/2}(\mathbb{R})]^2 : \phi - \psi \in \tilde{H}^{\epsilon-1/2}(\Omega), \mathcal{F}^{-1}t \cdot \mathcal{F}(\phi + \psi) = 0 \right\}.$$

Taking into account the representation formula (2.2), we have that the trace operator,

$$T_0 : D(\mathcal{P}) \rightarrow Z \\ u \mapsto u_0 = \begin{bmatrix} u_0^+ \\ u_0^- \end{bmatrix},$$

is continuously invertible by the Poisson operator $K : u_0 \mapsto u$ defined by (2.2).

On the other hand, a direct computation leads us to

$$(2.3) \quad \mathcal{P} = -\frac{1}{2} W_{t,\Omega} R_1 \mathcal{C} T_0,$$

where R_1 is the restriction operator to the first component and \mathcal{C} is the convolution operator (on the full line)

$$\mathcal{C} = \mathcal{F}^{-1} \begin{bmatrix} 1 & -1 \\ -t & -t \end{bmatrix} \cdot \mathcal{F} : Z \rightarrow \tilde{H}^{\epsilon-1/2}(\Omega) \times \{0\}.$$



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents




Go Back

Close

Quit

Page 6 of 20

Thus, (2.3) exhibits an operator equivalence between \mathcal{P} and the convolution type operator $W_{t,\Omega}$, defined in (2.1), because $R_1 \mathcal{C} T_0$ is continuously invertible by $K \mathcal{C}^{-1} [I \ 0]^T$ (please note that, for $u \in D(\mathcal{P})$, we have $u_1^+ = u_1^-$ due to (1.2) and (1.3)). \square



**Strongly Elliptic Operators for a
Plane Wave Diffraction Problem
in Bessel Potential Spaces**

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 7 of 20

with $\alpha = (\epsilon - 1, \dots, \epsilon - 1)$ and $s = 1/2$.

Proof. Let us consider the operators

$$E_2 = r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \lambda_-^{-\sigma} \cdot \mathcal{F} l_0 : [L^2(\mathbb{R}_+)]^{2n} \rightarrow H^\sigma(\mathbb{R}_+)$$

$$F_2 = l_0 r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \lambda_+^\sigma \cdot \mathcal{F} : \tilde{H}^\sigma(\mathbb{R}_+) \rightarrow [L^2_+(\mathbb{R})]^{2n},$$

where $l_0 : [L^2(\mathbb{R}_+)]^{2n} \rightarrow [L^2_+(\mathbb{R})]^{2n}$ is the zero extension operator.

These operators are bounded invertible (see [17, §2.10.3]). Moreover, attending to the structure of E_2 and F_2 [17], we have

$$W_{\Psi, \mathbb{R}_+} = E_2 W_{\Psi_0, \mathbb{R}_+} F_2,$$


which demonstrates operator equivalence between W_{Ψ, \mathbb{R}_+} and W_{Ψ_0, \mathbb{R}_+} . □

Corollary 3.3. *The convolution type operator $W_{t, \Omega}$ and any of the Wiener-Hopf operators W_{Ψ, \mathbb{R}_+} , W_{Ψ_0, \mathbb{R}_+} belong to the same regularity class [5]. More precisely, any of these three operators is invertible, one-sided invertible, Fredholm, semi-Fredholm, one-sided regularizable, generalized invertible or normally solvable, if and only if one of the others enjoys that property.*

Proof. From the above relations we derive that $\ker W_{t, \Omega}$, $\ker W_{\Psi, \mathbb{R}_+}$ and $\ker W_{\Psi_0, \mathbb{R}_+}$ are isomorphic and that the ranges of these operators are closed only at the same time. In addition, the presented relations allow us to conclude that

$$H^{\epsilon-3/2}(\Omega) / \text{im } W_{t, \Omega}, \quad H^\sigma(\mathbb{R}_+) / \text{im } W_{\Psi, \mathbb{R}_+}, \quad [L^2(\mathbb{R}_+)]^{2n} / \text{im } W_{\Psi_0, \mathbb{R}_+}$$

are also isomorphic. Thus, the statement follows. □



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20

4. Strong Ellipticity and Corresponding Well-posedness of the Problem

Let us consider the $2n \times 2n$ matrix function

$$\zeta^\beta = \text{diag} [\zeta^{-\epsilon+1/2}, \dots, \zeta^{-\epsilon+1/2}, \zeta^{-\epsilon+3/2}, \dots, \zeta^{-\epsilon+3/2}]$$

(where the first n elements in the main diagonal are equal and the last n elements are also all equal). We introduce the auxiliary operator

$$(4.1) \quad W_{\zeta^\beta \Psi_0, \mathbb{R}_+} = r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \zeta^\beta \Psi_0 \cdot \mathcal{F} : [L_+^2(\mathbb{R})]^{2n} \rightarrow [L^2(\mathbb{R}_+)]^{2n}.$$


This new operator will help us to arrive at the desired invertibility conditions. For this purpose, first, let us present some symmetries in the structure of $W_{\zeta^\beta \Psi_0, \mathbb{R}_+}$.

Theorem 4.1. *The Wiener-Hopf operator $W_{\zeta^\beta \Psi_0, \mathbb{R}_+}$ is equivalent to*

$$W_{\Psi_1, \mathbb{R}_+} \circ W_{\Psi_2, \mathbb{R}_+} \circ W_{\overline{\Psi_1}, \mathbb{R}_+} : [L_+^2(\mathbb{R})]^{2n} \rightarrow [L^2(\mathbb{R}_+)]^{2n}$$

where, for $j = 1, 2$,

$$W_{\Psi_j, \mathbb{R}_+} = r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \Psi_j \cdot \mathcal{F} : [L_+^2(\mathbb{R})]^{2n} \rightarrow [L^2(\mathbb{R}_+)]^{2n},$$



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit


Page 11 of 20

$$\Psi_1 = \left[\begin{array}{c|ccc} & & & \\ \hline & & & \tau_{-\gamma_{2n}} \\ & & & \tau_{-\gamma_{2n-2}} \\ & & & \dots \\ & & & \tau_{-\gamma_2} \\ \hline & \tau_{-\gamma_{2n-1}} & & \\ & & \tau_{-\gamma_{2n-3}} & \\ & & & \dots \\ & & & \tau_{-\gamma_3} \\ \hline n & 1 & 1 & \dots & 1 \end{array} \right]$$

and

$$\Psi_2 = \left[\begin{array}{c|ccc} 1 & & & \zeta^{1/2}/n \dots \zeta^{1/2}/n \\ & \dots & & \\ & & 1 & \\ \hline & & & 1 \\ & & & \dots \\ & & & 1 \end{array} \right]$$

Proof. From the identity



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 12 of 20

$$\zeta^\beta \Psi_0 = \left[\begin{array}{c|c} \begin{array}{ccc} & & 1 \\ & \ddots & \\ 1 & & \end{array} & \\ \hline & \begin{array}{c|c} & 1 \\ \hline 1 & \ddots \\ & 1 \end{array} \end{array} \right] \Psi_1 \Psi_2 \overline{\Psi_1}$$


$$\times \left[\begin{array}{c|c|c|c} \begin{array}{ccc} 1/n & -1/n & \cdots & -1/n \\ & & & 1 \end{array} & & & \\ \hline & \begin{array}{ccc} & & \ddots \\ & 1 & \\ & & \end{array} & & \\ \hline & & & \begin{array}{c} 1/n \\ \\ \\ 1 \end{array} \end{array} \right]$$

it follows that $W_{\zeta^\beta \Psi_0, \mathbb{R}_+}$ and

$$W_{\Psi_1 \Psi_2 \overline{\Psi_1}, \mathbb{R}_+} = r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \Psi_1 \Psi_2 \overline{\Psi_1} \cdot \mathcal{F} : [L_+^2(\mathbb{R})]^{2n} \rightarrow [L^2(\mathbb{R}_+)]^{2n}$$

are equivalent operators.

Therefore, in order to obtain the present result, we only have to observe that the “minus” and “plus” semi-almost periodic entries of Ψ_1 and $\overline{\Psi_1}$, respectively,



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

lead us to

$$\begin{aligned} W_{\Psi_1 \Psi_2 \overline{\Psi}_1, \mathbb{R}_+} &= (r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \Psi_1 \cdot \mathcal{F}) l_0 (r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \Psi_2 \cdot \mathcal{F}) l_0 (r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \overline{\Psi}_1 \cdot \mathcal{F}) \\ &= W_{\Psi_1, \mathbb{R}_+} l_0 W_{\Psi_2, \mathbb{R}_+} l_0 W_{\overline{\Psi}_1, \mathbb{R}_+}. \end{aligned}$$

□

We recall that an essentially bounded $m \times m$ matrix-valued function $\psi(\xi) = [\psi_{ij}(\xi)]$ is said to be *strongly elliptic* if there exist constants $\eta \in \mathbb{C}$ and $C > 0$ so that

$$\Re \eta \sum_{i,j=1}^m \psi_{ij}(\xi) \mu_j \overline{\mu_i} \geq C \sum_{i=1}^m |\mu_i|^2, \quad \forall \mu = (\mu_1, \dots, \mu_m) \in \mathbb{C}^m.$$

Theorem 4.2. *The Wiener-Hopf operator introduced in (4.1),*


$$W_{\zeta^\beta \Psi_0, \mathbb{R}_+} : [L_+^2(\mathbb{R})]^{2n} \rightarrow [L^2(\mathbb{R}_+)]^{2n},$$

is an invertible operator.

Proof. The nonnegative mean motions [14] of the semi-almost periodic entries of $\overline{\Psi}_1$ ensure us (see [14, Theorem 1]) the left invertibility of $W_{\overline{\Psi}_1, \mathbb{R}_+}$.

Moreover, we observe that Ψ_2 is strongly elliptic. In fact, from

$$\left| \frac{\zeta^{1/2}}{n} \mu_j \overline{\mu_1} \right| \leq \frac{1}{2n} (|\mu_j|^2 + |\mu_1|^2), \quad j = n+1, \dots, 2n,$$



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 14 of 20

it follows that, for $\Psi_2(\xi) = [\Theta_{ij}(\xi)]$,


$$\begin{aligned} \Re \sum_{i,j=1}^{2n} \Theta_{ij}(\xi) \mu_j \bar{\mu}_i &\geq \sum_{j=1}^{2n} |\mu_j|^2 - \frac{1}{2n} \sum_{j=n+1}^{2n} (|\mu_j|^2 + |\mu_1|^2) \\ &= \frac{1}{2} |\mu_1|^2 + \sum_{j=2}^n |\mu_j|^2 + \left(1 - \frac{1}{2n}\right) \sum_{j=n+1}^{2n} |\mu_j|^2 \\ &\geq \frac{1}{2} \sum_{j=1}^{2n} |\mu_j|^2, \end{aligned}$$

for all $(\mu_1, \dots, \mu_{2n}) \in \mathbb{C}^{2n}$.

Consequently, for $\psi \in [L^2(\mathbb{R}_+)]^{2n}$, we have

$$\begin{aligned} \Re \langle (W_{\Psi_1, \mathbb{R}_+} l_0)(W_{\Psi_2, \mathbb{R}_+} l_0)(W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi, \psi \rangle &= \Re \langle (W_{\Psi_2, \mathbb{R}_+} l_0)(W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi, (W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi \rangle \\ &= \Re \langle (r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \Psi_2 \cdot \mathcal{F} l_0)(W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi, (W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi \rangle \\ &= \Re \langle \Psi_2 \cdot \mathcal{F} l_0(W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi, \mathcal{F} l_0(W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi \rangle \\ &\geq \frac{1}{2} \|\mathcal{F} l_0(W_{\overline{\Psi_1}, \mathbb{R}_+} l_0) \psi\|^2 \\ &= \frac{1}{2} \|W_{\overline{\Psi_1}, \mathbb{R}_+} l_0 \psi\|^2 \\ &\geq \frac{1}{2} C_1 \|\psi\|^2, \end{aligned}$$

where $C_1 > 0$ is provided by the left invertibility of $W_{\overline{\Psi_1}, \mathbb{R}_+}$. This inequality allows us to conclude that $W_{\Psi_1, \mathbb{R}_+} l_0 W_{\Psi_2, \mathbb{R}_+} l_0 W_{\overline{\Psi_1}, \mathbb{R}_+} : [L^2_+(\mathbb{R})]^{2n} \rightarrow [L^2(\mathbb{R}_+)]^{2n}$



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

is a left invertible operator.

Applying the same reasoning to the conjugate operator

$$\begin{aligned} \left(l_0 W_{\Psi_1, \mathbb{R}_+} l_0 W_{\Psi_2, \mathbb{R}_+} l_0 W_{\overline{\Psi_1}, \mathbb{R}_+} \right)^* &= \left(l_0 W_{\overline{\Psi_1}, \mathbb{R}_+} \right)^* \left(l_0 W_{\Psi_2, \mathbb{R}_+} \right)^* \left(l_0 W_{\Psi_1, \mathbb{R}_+} \right)^* \\ &= l_0 W_{\Psi_1, \mathbb{R}_+} l_0 W_{\overline{\Psi_2}, \mathbb{R}_+} l_0 W_{\overline{\Psi_1}, \mathbb{R}_+} \end{aligned}$$

we obtain that this is also a left invertible operator.

Thus $W_{\Psi_1, \mathbb{R}_+} l_0 W_{\Psi_2, \mathbb{R}_+} l_0 W_{\overline{\Psi_1}, \mathbb{R}_+}$ is an invertible operator and from Theorem 4.1 our goal is achieved. \square

Corollary 4.3. *The Wiener-Hopf operator W_{Ψ_0, \mathbb{R}_+} , defined in (3.3), is a Fredholm operator with zero Fredholm index.*

Proof. From Theorem 4.2 we know that $W_{\zeta^\beta \Psi_0, \mathbb{R}_+}$ is an invertible operator. Thus (see e.g. [11, Chapter 1, Theorem 3.11]), the result is a consequence of W_{Ψ_0, \mathbb{R}_+} and $W_{\zeta^\beta \Psi_0, \mathbb{R}_+}$ being homotopic operators in the class of Fredholm operators acting from $[L^2_+(\mathbb{R})]^{2n}$ to $[L^2(\mathbb{R}_+)]^{2n}$. \square

Theorem 4.4. *The operator \mathcal{P} (associated to the problem) is bounded invertible and, therefore, our problem is well-posed for all orders of smoothness $\epsilon \in]1/2, 3/2[$.*

Proof. Due to the fact that, for negative parameters $-\gamma$ and any $s \in \mathbb{R}$,

$$r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} \tau_{-\gamma} \cdot \mathcal{F} : \widetilde{H}^s(\mathbb{R}_+) \rightarrow H^s(\mathbb{R}_+)$$



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit


Page 16 of 20

are nothing more than left shift operators composed with the restriction operator $r_{\mathbb{R} \rightarrow \mathbb{R}_+}$, we have that these operators are surjective. Therefore, taking into account the structure of Ψ (see (3.1)), we obtain that W_{Ψ, \mathbb{R}_+} is a surjective operator whenever

$$W_{t, \mathbb{R}_+} = r_{\mathbb{R} \rightarrow \mathbb{R}_+} \mathcal{F}^{-1} t \cdot \mathcal{F} : \tilde{H}^{\epsilon-1/2}(\mathbb{R}_+) \rightarrow H^{\epsilon-3/2}(\mathbb{R}_+)$$

is a surjective operator, which is true for all ϵ . As a consequence, the codimension of the image of W_{Ψ, \mathbb{R}_+} is zero.

Thus, from Corollary 3.3 and Corollary 4.3, we have that $W_{t, \Omega}$ is invertible. Therefore, the result is obtained if we take into consideration Theorem 2.1. \square



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back


Close

Quit

Page 17 of 20

References

- [1] H. BART AND V.È. TSEKANOVSKII, Matricial coupling and equivalence after extension, in *Operator Theory and Complex Analysis* (Sapporo, 1991), T. Ando (ed.) et al., Oper. Theory Adv. Appl. 59, Birkhäuser, Basel 1992, 143–160.
- [2] L.P. CASTRO, Wiener-Hopf operators on unions of finite intervals: relations and generalized inversion, in *Proceedings of the Meeting on Matrix Analysis and Applications* (Sevilla, 1997), F.J. Cobos (ed.) et al., Department of Applied Mathematics I, University of Sevilla, Sevilla 1997, 148–155.
- [3] L.P. CASTRO, A relation between convolution type operators on intervals in Sobolev spaces, *Appl. Anal.*, **74**(3-4) (2000), 393–412.
- [4] L.P. CASTRO, Solution of a wave diffraction problem by a magnetic strip, in *Proceedings of the Fifth International Conference on Mathematical and Numerical Aspects of Wave Propagation* (Santiago de Compostela, 2000), A. Bermúdez (ed.) et al., SIAM, Philadelphia 2000, 231–235.
- [5] L.P. CASTRO AND F.-O. SPECK, Regularity properties and generalized inverses of delta-related operators, *Z. Anal. Anwendungen*, **17**(3) (1998), 577–598.
- [6] L.P. CASTRO AND F.-O. SPECK, Relations between convolution type operators on intervals and on the half-line, *Integral Equations Operator Theory*, **37**(2) (2000), 169–207.



Strongly Elliptic Operators for a
Plane Wave Diffraction Problem
in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents




Go Back

Close

Quit

Page 18 of 20

- [7] A.E. HEINS, The Sommerfeld half-plane problem revisited. I: The solution of a pair of coupled Wiener-Hopf integral equations, *Math. Methods Appl. Sci.*, **4**(1) (1982), 74–90.
- [8] A.B. KUIJPER, A note on first kind convolution equations on a finite interval, *Integral Equations Operator Theory*, **14**(1) (1991), 146–152.
- [9] E. LUENEBURG, The Sommerfeld problem: Methods, generalization and frustrations, in *Modern Mathematical Methods in Diffraction Theory and its Applications in Engineering, Proceedings of the Sommerfeld '96 Workshop* (Freudenstadt 1996), E. Meister (ed.), Methoden Verfahren Math. Phys. 42, Peter Lang, Verlag der Wissenschaften, Frankfurt am Main 1997, 145–162.
- [10] E. MEISTER AND F.-O. SPECK, Modern Wiener-Hopf methods in diffraction theory, in *Ordinary and Partial Differential Equations Vol. II* (Dundee, 1988), B. Sleeman (ed.) et al., Pitman Res. Notes Math. Ser. 216, Longman Sci. Tech., Harlow 1989, 130–171.
- [11] S.G. MIKHLIN AND S. PRÖSSDORF, *Singular Integral Operators*, Springer-Verlag, Berlin-New York 1986.
- [12] A.D. RAWLINS, The solution of a mixed boundary value problem in the theory of diffraction by a semi-infinite plane, *Proc. Roy. Soc. London Ser. A*, **346** (1975), 469–484.
- [13] P.A. SANTOS AND F.S. TEIXEIRA, Sommerfeld half-plane problems with higher order boundary conditions, *Math. Nachr.*, **171** (1995), 269–



Strongly Elliptic Operators for a Plane Wave Diffraction Problem in Bessel Potential Spaces

Luis P. Castro

Title Page

Contents



Go Back


Close

Quit

Page 19 of 20

282.

- [14] D. SARASON, Toeplitz operators with semi-almost periodic symbols, *Duke Math. J.*, **44**(2) (1977), 357–364.
- [15] A. SOMMERFELD, Mathematische theorie der diffraction, *Math. Ann.*, **47** (1896), 317–374.
- [16] F.-O. SPECK, Sommerfeld diffraction problems with first and second kind boundary conditions, *SIAM J. Math. Anal.*, **20**(2) (1989), 396–407.
- [17] H. TRIEBEL, *Interpolation Theory, Function Spaces, Differential Operators*, 2nd edition, Johann Ambrosius Barth, Heidelberg 1995.



**Strongly Elliptic Operators for a
Plane Wave Diffraction Problem
in Bessel Potential Spaces**

Luis P. Castro

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20