

Journal of Inequalities in Pure and Applied Mathematics



APPLICATION OF A TRAPEZOID INEQUALITY TO NEUTRAL FREDHOLM INTEGRO-DIFFERENTIAL EQUATIONS IN BANACH SPACES

ALEXANDRU MIAHI BICA, VASILE AUREL CĂUŞ AND SORIN MUREŞAN

Department of Mathematics
University of Oradea,
Str. Universitatii no.1
410087, Oradea, Romania

EMail: abica@uoradea.ro
EMail: vcaus@uoradea.ro

volume 7, issue 5, article 173,
2006.

Received 27 January, 2006;
accepted 19 November, 2006.

Communicated by: *S.S. Dragomir*

Abstract

Contents



Home Page

Go Back

Close

Quit



Abstract

A new approach for neutral Fredholm integro-differential equations in Banach spaces, using the Perov's fixed point theorem of existence, uniqueness and approximation is presented. The approximation of the solution and of its derivative is realized using the method of successive approximations and a trapezoidal quadrature rule in Banach spaces for Lipschitzian functions. The interest is focused on the error estimation.

2000 Mathematics Subject Classification: Primary 45J05, Secondary 45N05, 45B05, 65L05.

Key words: Nonlinear neutral Fredholm integro-differential equations in Banach spaces, Perov's fixed point theorem, Method of successive approximations, Trapezoidal quadrature rule.

Contents

1	Introduction	3
2	Existence, Uniqueness and Approximation	5
3	The Main Result	11
References		

Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 2 of 24

1. Introduction

Consider the neutral Fredholm integro-differential equation

$$(1.1) \quad x(t) = \int_a^b f(t, s, x(s), x'(s)) ds + g(t), \quad t \in [a, b]$$

where $f : [a, b] \times [a, b] \times X \times X \rightarrow X$ is continuous, X is a Banach space and $g \in C^1([a, b], X)$.

To obtain the existence, uniqueness and global approximation of the solution of (1.1) we will use Perov's fixed point theorem. To this purpose, we differentiate the equation (1.1) with respect to t and assume that $f(\cdot, s, u, v) \in C^1([a, b], X)$, $\forall s \in [a, b]$, $\forall u, v \in X$, where $x' = y$. Hence (1.1) reduces to the following system of Fredholm integral equations:

$$(1.2) \quad \begin{cases} x(t) = \int_a^b f(t, s, x(s), y(s)) ds + g(t) \\ y(t) = \int_a^b \frac{\partial f}{\partial t}(t, s, x(s), y(s)) ds + g'(t) \end{cases}, \quad t \in [a, b].$$

The Perov fixed point theorem will be applied to the system (1.2) obtaining also the approximation of the solution of (1.1) and its derivative.

The Perov fixed point theorem appeared for the first time in [16] and was later used for two point boundary value problems of second order differential equations in [5]. The Perov fixed point theorem was also used in [1], [10], [18] and [19]. Bica and Muresan have used the Perov fixed point theorem for delay neutral integro-differential equations in [6] and [7]. In this paper the authors have constructed a method of approximating the solution of (1.1) and



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 3 of 24

its derivative using a sequence of successive approximations and a trapezoidal quadrature rule from [8]. Some of the existing numerical methods applied to Fredholm integro-differential equations can be found in the papers [2], [3], [4], [9], [11], [12], [13], [14], [15], [17]. The tools utilised in these papers are: the tau method, direct methods, collocation methods, Runge-Kutta methods, wavelet methods and spline approximation.

In this paper, our interest will be focused on the error estimation of the method presented in the Section 3.



**Application of a Trapezoid
Inequality to Neutral Fredholm
Integro-Differential Equations in
Banach Spaces**

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 4 of 24

2. Existence, Uniqueness and Approximation

Consider the following conditions:

(i) (continuity): $f \in C([a, b] \times [a, b] \times X \times X, X)$, $g \in C^1([a, b], X)$ and $f(\cdot, s, u, v) \in C^1([a, b], X)$, for any $s \in [a, b]$, $u, v \in X$

(ii) (Lipschitz conditions): there exist $\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2, \delta_1, \delta_2, \eta \in \mathbb{R}_+^*$ such that for any $t, t', s, s_1, s_2 \in [a, b]$ and $u, v, u_1, u_2, v_1, v_2 \in X$, we have:


$$(2.1) \quad \|f(t, s, u_1, v_1) - f(t, s, u_2, v_2)\|_X \\ \leq \alpha_1 \|u_1 - u_2\|_X + \beta_1 \|v_1 - v_2\|_X,$$

$$(2.2) \quad \left\| \frac{\partial f}{\partial t}(t, s, u_1, v_1) - \frac{\partial f}{\partial t}(t, s, u_2, v_2) \right\|_X \\ \leq \alpha_2 \|u_1 - u_2\|_X + \beta_2 \|v_1 - v_2\|_X,$$

$$(2.3) \quad \|f(t, s_1, u, v) - f(t, s_2, u, v)\|_X \leq \gamma_1 |s_1 - s_2|,$$

$$(2.4) \quad \left\| \frac{\partial f}{\partial t}(t, s_1, u, v) - \frac{\partial f}{\partial t}(t, s_2, u, v) \right\|_X \leq \gamma_2 |s_1 - s_2|,$$

$$(2.5) \quad \|f(t, s, u, v) - f(t', s, u, v)\|_X \leq \delta_1 |t - t'|,$$



Application of a Trapezoid
Inequality to Neutral Fredholm
Integro-Differential Equations in
Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 5 of 24



$$(2.6) \quad \left\| \frac{\partial f}{\partial t} (t, s, u, v) - \frac{\partial f}{\partial t} (t', s, u, v) \right\|_X \leq \delta_2 |t - t'|,$$

$$(2.7) \quad \|g'(t) - g'(t')\|_X \leq \eta \cdot |t - t'|.$$

We use Perov's fixed point theorem (see [16], [5] and [10]):

Theorem 2.1. *Let (X, d) be a complete generalized metric space such that $d(x, y) \in \mathbb{R}^n$ for $x, y \in X$. Suppose that there exists a function $A : X \rightarrow X$ such that:*

$$d(A(x), A(y)) \leq Q \cdot d(x, y)$$

for any $x, y \in X$, where $Q \in \mathcal{M}_n(\mathbb{R}_+)$. If all eigenvalues of Q lie in the open unit disc from \mathbb{R}^2 then $Q^m \rightarrow 0$ for $m \rightarrow \infty$ and the operator A has a unique fixed point $x^* \in X$. Moreover, the sequence of successive approximations $x_m = A(x_{m-1})$ converges to x^* in X for any $x_0 \in X$ and the following estimation holds:

$$(2.8) \quad d(x_m, x^*) \leq Q^m (I_n - Q)^{-1} \cdot d(x_0, x_1), \quad \text{for each } m \in \mathbb{N}^*$$

where I_n is the unity matrix in $\mathcal{M}_n(\mathbb{R})$.

We recall the notion of generalized metric, which is a function $d : Y \times Y \rightarrow \mathbb{R}^n$ on a nonempty set Y with the properties:

- a) $d(x, y) \geq 0$, for any $x, y \in Y$;

**Application of a Trapezoid
Inequality to Neutral Fredholm
Integro-Differential Equations in
Banach Spaces**

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 6 of 24

- b) $d(x, y) = 0 \Leftrightarrow x = y$, for $x, y \in Y$;
- c) $d(y, x) = d(x, y)$, for any $x, y \in Y$;
- d) $d(x, z) \leq d(x, y) + d(y, z)$, for any $x, y, z \in Y$.

Here, the order relation on \mathbb{R}^n is defined by

$$x \leq y \Leftrightarrow x_i \leq y_i, \quad \text{in } \mathbb{R}, \text{ for each } i = \overline{1, n}, \quad x_i, y_i \in \mathbb{R}$$

for $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$. The pair (Y, d) denote a generalized metric space.

In the following, we will use the notation

$$\|u\|_C = \max \{\|u(t)\|_X : t \in [a, b]\},$$

for $u \in C([a, b], X)$.


We consider the product space $Y = C([a, b], X) \times C([a, b], X)$ and define the generalized metric $d : Y \times Y \rightarrow \mathbb{R}^2$ by,

$$d((u_1, v_1), (u_2, v_2)) = (\|u_1 - u_2\|_C, \|v_1 - v_2\|_C),$$

for $(u_1, v_1), (u_2, v_2) \in Y$.

It is easy to prove that (Y, d) is a complete generalized metric space. We define the operator $A : Y \rightarrow Y$, $A = (A_1 A_2)$ by,

$$A_1(x, y)(t) = \int_a^b f(t, s, x(s), y(s)) ds + g(t)$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

**Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan**

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 7 of 24

and

$$(2.9) \quad A_2(x, y)(t) = \int_a^b \frac{\partial f}{\partial t}(t, s, x(s), y(s)) ds + gt(t), \quad t \in [a, b].$$

The following result concerning the existence and uniqueness of the solution for the equation (1.1) holds.

Theorem 2.2. *In the conditions (i), (2.1), (2.2), if $(\alpha_1 + \beta_2)(b - a) < 2$ and*

$$(2.10) \quad 1 + (b - a)^4 (\alpha_1 \beta_2 - \alpha_2 \beta_1)^2 + 2\alpha_2 \beta_1 (b - a)^2 > (b - a)^2 (\alpha_1^2 + \beta_2^2)$$

then the operator A has a unique fixed point (x^, y^*) such that*

$$x^* \in C^1([a, b], X), \quad y^* = (x^*)'$$

and x^ is the unique solution of the equation (1.1). Moreover, the sequence of the successive approximations given by,*

$$(2.11) \quad (x_0(t), y_0(t)) = (g(t), g'(t)), \quad t \in [a, b]$$

$$(2.12) \quad x_{m+1}(t) = \int_a^b f(t, s, x_m(s), y_m(s)) ds + g(t), \quad t \in [a, b]$$

and

$$(2.13) \quad y_{m+1}(t) = \int_a^b \frac{\partial f}{\partial t}(t, s, x_m(s), y_m(s)) ds + g'(t), \quad t \in [a, b],$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 8 of 24

converges in Y to (x^*, y^*) and the following error estimation holds:

$$(2.14) \quad d((x_m, y_m), (x^*, y^*)) \leq Q^m (I_n - Q)^{-1} \cdot d((x_0, y_0), (x_1, y_1)),$$

for any $m \in \mathbb{N}^*$,

where

$$Q = (b - a) \begin{pmatrix} \alpha_1 & \beta_1 \\ \alpha_2 & \beta_2 \end{pmatrix}.$$

Proof. From condition (i) we infer that $A(Y) \subset Y$. For $(u_1, v_1), (u_2, v_2) \in Y$, $t \in [a, b]$ we have


$$\begin{aligned} & \|A_1(u_1, v_1)(t) - A_1(u_2, v_2)(t)\|_X \\ & \leq \int_a^b [\alpha_1 \|u_1(s) - u_2(s)\|_X + \beta_1 \|v_1(s) - v_2(s)\|_X] \\ & \leq (b - a) \cdot [\alpha_1 \|u_1 - u_2\|_C + \beta_1 \|v_1 - v_2\|_C], \quad \text{for any } t \in [a, b] \end{aligned}$$

and

$$\begin{aligned} & \|A_2(u_1, v_1)(t) - A_2(u_2, v_2)(t)\|_X \\ & \leq (b - a) \cdot [\alpha_2 \|u_1 - u_2\|_C + \beta_2 \|v_1 - v_2\|_C], \quad \text{for any } t \in [a, b]. \end{aligned}$$

We infer that

$$\begin{aligned} d(A(u_1, v_1), A(u_2, v_2)) & \leq Q \cdot d((u_1, v_1), (u_2, v_2)), \\ & \quad \text{for any } (u_1, v_1), (u_2, v_2) \in Y, \end{aligned}$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 9 of 24

where

$$Q = \begin{pmatrix} (b-a)\alpha_1 & (b-a)\beta_1 \\ (b-a)\alpha_2 & (b-a)\beta_2 \end{pmatrix}.$$

It is easy to see that the eigenvalues of Q are real. The inequalities (2.10) and $(\alpha_1 + \beta_2)(b-a) < 2$ lead to $\mu_1, \mu_2 \in (-1, 1)$, where μ_1 and μ_2 are these eigenvalues. From the Perov fixed point theorem we infer that $Q^m \rightarrow 0$ for $m \rightarrow \infty$ and the operator A has a unique fixed point $(x^*, y^*) \in Y$. Then,

$$x^*(t) = \int_a^b f(t, s, x^*(s), y^*(s)) ds + g(t), \quad \forall t \in [a, b]$$

and

$$y^*(t) = \int_a^b \frac{\partial f}{\partial t}(t, s, x^*(s), y^*(s)) ds + g'(t), \quad \forall t \in [a, b].$$

Since $f(\cdot, s, u, v) \in C^1([a, b], X)$, for any $s \in [a, b]$, $u, v \in X$ and $g \in C^1([a, b], X)$ we infer that $x^* \in C^1([a, b], X)$. If we differentiate the first equality with respect to t we obtain $y^* = (x^*)'$. Then x^* is the unique solution of (1.1). From the relations (2.12) and (2.13) and from (2.9) we infer that the sequences given in (2.12), (2.13) fulfil the recurrence relation

$$(x_{m+1}, y_{m+1}) = A((x_m, y_m)), \quad \forall m \in \mathbb{N}.$$

Now, the inequality (2.14) follows from the estimation (2.8). Since $Q^m \rightarrow 0$ for $m \rightarrow \infty$ in $\mathcal{M}_2(\mathbb{R})$ we infer that

$$\lim_{m \rightarrow \infty} d((x_m, y_m), (x^*, y^*)) = (0, 0).$$

This proves the theorem. □



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 10 of 24](#)

3. The Main Result

To compute the terms of the sequence of successive approximations we use in the calculus of integrals in (2.12), (2.13) the trapezoidal quadrature rule for Lipschitzian functions from [8]:

$$(3.1) \quad \int_a^b F(x) dx = \frac{(b-a)}{2n} \left[F(a) + 2 \sum_{i=1}^{n-1} F\left(a + \frac{i(b-a)}{n}\right) + F(b) \right] + R_n(F)$$

with

$$(3.2) \quad \|R_n(F)\|_X \leq \frac{L(b-a)^2}{4n},$$

where L is the Lipschitz constant of $F : [a, b] \rightarrow X$.


In this respect consider the uniform partition of $[a, b]$,

$$(3.3) \quad \Delta : a = t_0 < t_1 < \cdots < t_{n-1} < t_n = b$$

with $t_i = a + \frac{i(b-a)}{n}$, $i = \overline{0, n}$ and compute $x_m(t_i), y_m(t_i)$, $i = \overline{0, n}$, $m \in \mathbb{N}^*$.

From (2.12) and (2.13) we have

$$(3.4) \quad x_{m+1}(t_i) = \int_a^b f(t_i, s, x_m(s), y_m(s)) ds + g(t_i)$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 11 of 24

and

$$(3.5) \quad y_{m+1}(t_i) = \int_a^b \frac{\partial f}{\partial t}(t_i, s, x_m(s), y_m(s)) ds + g'(t_i), \\ \text{for any } i = \overline{0, n}, \quad m \in \mathbb{N}.$$

We define the functions

$$F_{m,i}, G_{m,i} : [a, b] \rightarrow X, \quad m \in \mathbb{N}, \quad i = \overline{0, n}$$

by

$$(3.6) \quad \begin{cases} F_{m,i}(s) = f(t_i, s, x_m(s), y_m(s)) \\ G_{m,i}(s) = \frac{\partial f}{\partial t}(t_i, s, x_m(s), y_m(s)) \end{cases}, \text{ for any } s \in [a, b].$$

Definition 3.1. A set $Z \subset C([a, b], X)$ is equally Lipschitz if there exists $L \geq 0$ such that for any $h \in Z$,

$$\|h(t) - h(t')\|_X \leq L \cdot |t - t'|, \quad \text{for each } t, t' \in [a, b].$$


Theorem 3.1. The subsets

$$\{\{F_{m,i}\}_{m \in \mathbb{N}}, \quad i = \overline{0, n}\} \subset C([a, b], X)$$

and

$$\{\{G_{m,i}\}_{m \in \mathbb{N}}, \quad i = \overline{0, n}\} \subset C([a, b], X)$$

defined in (3.6), are equally Lipschitz, if the conditions (i) and (2.1) – (2.7) are true.



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 12 of 24

Proof. Let

$$\mu = \|g'\|_C = \max \{ \|g'(t)\|_X : t \in [a, b] \}.$$

For $m \in \mathbb{N}^*$ and $i = \overline{0, n}$ we have

$$\begin{aligned} & \|F_{m,i}(s_1) - F_{m,i}(s_2)\|_X \\ & \leq \gamma_1 |s_1 - s_2| + \alpha_1 \|x_m(s_1) - x_m(s_2)\|_X + \beta_1 \|y_m(s_1) - y_m(s_2)\|_X \\ & \leq \gamma_1 |s_1 - s_2| + \alpha_1 [\delta_1(b-a) + \mu] \cdot |s_1 - s_2| \\ & \quad + \beta_1 [\delta_2(b-a) \cdot |s_1 - s_2| + \|g'(s_1) - g'(s_2)\|_X] \\ & \leq [\gamma_1 + \alpha_1\mu + \beta_1\eta + (b-a)(\alpha_1\delta_1 + \beta_1\delta_2)] \cdot |s_1 - s_2|, \end{aligned}$$

for any $s_1, s_2 \in [a, b]$ and

$$\begin{aligned} & \|G_{m,i}(s_1) - G_{m,i}(s_2)\|_X \\ & \leq \gamma_2 |s_1 - s_2| + \alpha_2 \|x_m(s_1) - x_m(s_2)\|_X + \beta_2 \|y_m(s_1) - y_m(s_2)\|_X \\ & \leq [\gamma_2 + \alpha_2\mu + \beta_2\eta + (b-a)(\alpha_2\delta_1 + \beta_2\delta_2)] \cdot |s_1 - s_2|, \end{aligned}$$

for any $s_1, s_2 \in [a, b]$. Moreover, for any $i = \overline{0, n}$ we have,


$$\|F_{0,i}(s_1) - F_{0,i}(s_2)\|_X \leq (\gamma_1 + \alpha_1\mu + \beta_1\eta) \cdot |s_1 - s_2|, \quad \text{for each } s_1, s_2 \in [a, b]$$

and

$$\|G_{0,i}(s_1) - G_{0,i}(s_2)\|_X \leq (\gamma_2 + \alpha_2\mu + \beta_2\eta) \cdot |s_1 - s_2|, \quad \text{for each } s_1, s_2 \in [a, b].$$

Let

$$L_1 = \gamma_1 + \alpha_1\mu + \beta_1\eta + (b-a)(\alpha_1\delta_1 + \beta_1\delta_2),$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Go Back

Close

Quit

Page 13 of 24

$$L_2 = \gamma_2 + \alpha_2\mu + \beta_2\eta + (b-a)(\alpha_2\delta_1 + \beta_2\delta_2).$$

From the above, we infer that for any $i = \overline{0, n}$, we have,

$$\|F_{m,i}(s_1) - F_{m,i}(s_2)\|_X \leq L_1 \cdot |s_1 - s_2|, \quad \text{for each } s_1, s_2 \in [a, b]$$

and

$$\|G_{m,i}(s_1) - G_{m,i}(s_2)\|_X \leq L_2 \cdot |s_1 - s_2|, \quad \text{for each } s_1, s_2 \in [a, b] \text{ and } m \in \mathbb{N}.$$

This concludes the proof of the theorem. \square

Applying in (3.4), (3.5) the quadrature rule (3.1) – (3.2) we obtain the numerical method:

$$(3.7) \quad x_0(t_i) = g(t_i), \quad y_0(t_i) = g'(t_i), \quad \text{for } i = \overline{0, n}$$

$$(3.8) \quad x_m(t_i) = g(t_i) + \frac{(b-a)}{2n} \cdot \left[f(t_i, a, x_{m-1}(a), y_{m-1}(a)) \right. \\ \left. + 2 \sum_{j=1}^{n-1} f(t_i; t_j, x_{m-1}(t_j), y_{m-1}(t_j)) + f(t_i, b, x_{m-1}(b), y_{m-1}(b)) \right] \\ + R_{m,i}, \quad \text{for } i = \overline{0, n} \text{ and } m \in \mathbb{N}^*$$

and

$$(3.9) \quad y_m(t_i) = g'(t_i) + \frac{(b-a)}{2n} \cdot \left[\frac{\partial f}{\partial t}(t_i, a, x_{m-1}(a), y_{m-1}(a)) \right.$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Go Back](#)

[Close](#)

[Quit](#)

[Page 14 of 24](#)

$$+ 2 \sum_{j=1}^{n-1} \frac{\partial f}{\partial t} (t_i; t_j, x_{m-1}(t_j), y_{m-1}(t_j)) + \frac{\partial f}{\partial t} (t_i, b, x_{m-1}(b), y_{m-1}(b)) \Big] \\ + R_{m,i}, \quad \text{for } i = \overline{0, n} \text{ and } m \in \mathbb{N}^*.$$


with the remainder estimations

$$(3.10) \quad \|R_{m,i}\|_X \leq \frac{L_1(b-a)^2}{4n}, \quad \text{for any } m \in \mathbb{N}^* \text{ and } i = \overline{0, n},$$

$$(3.11) \quad \|\omega_{m,i}\|_X \leq \frac{L_2(b-a)^2}{4n}, \quad \text{or any } m \in \mathbb{N}^* \text{ and } i = \overline{0, n}.$$

These lead to the following algorithm:

$$(3.12) \quad x_0(t_i) = g(t_i), y_0(t_i) = g'(t_i), \\ = g(t_i) + \frac{(b-a)}{2n} \cdot \left[f(t_i, a, g(a), g'(a)) \right. \\ \left. + 2 \sum_{j=1}^{n-1} f(t_i; t_j, g(t_j), g'(t_j)) + f(t_i, b, g(b), g'(b)) \right] + R_{1,i} \\ = \bar{x}_1(t_i) + R_{1,i},$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)




[Go Back](#)

[Close](#)

[Quit](#)

[Page 15 of 24](#)

$$(3.13) \quad y_1(t_i) = g'(t_i) + \frac{(b-a)}{2n} \cdot \left[\frac{\partial f}{\partial t}(t_i, a, g(a), g'(a)) \right. \\ + 2 \sum_{j=1}^{n-1} \frac{\partial f}{\partial t}(t_i; t_j, g(t_j), g'(t_j)) \\ \left. + \frac{\partial f}{\partial t}(t_i, b, g(b), g'(b)) \right] + \omega_{1,i} \\ = \overline{y_1}(t_i) + \omega_{1,i},$$



$$(3.14) \quad x_2(t_i) = g(t_i) + \frac{(b-a)}{2n} \cdot \left[f(t_i, t_0, \overline{x_1}(t_0) + R_{1,0}, \overline{y_1}(t_0) + \omega_{1,0}) \right. \\ + 2 \sum_{j=1}^{n-1} f(t_i; t_j, \overline{x_1}(t_j) + R_{1,j}, \overline{y_1}(t_j) + \omega_{1,j}) \\ \left. + f(t_i, t_n, \overline{x_1}(t_n) + R_{1,n}, \overline{y_1}(t_n) + \omega_{1,n}) \right] + R_{2,i} \\ = g(t_i) + \frac{(b-a)}{2n} \cdot \left[f(t_i, t_0, \overline{x_1}(t_0), \overline{y_1}(t_0)) \right. \\ + 2 \sum_{j=1}^{n-1} f(t_i; t_j, \overline{x_1}(t_j), \overline{y_1}(t_j)) \\ \left. + f(t_i, t_n, \overline{x_1}(t_n), \overline{y_1}(t_n)) \right] + \overline{R}_{2,i} \\ = \overline{x_2}(t_i) + \overline{R}_{2,i},$$

**Application of a Trapezoid
Inequality to Neutral Fredholm
Integro-Differential Equations in
Banach Spaces**

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)




[Go Back](#)

[Close](#)

[Quit](#)

Page 16 of 24



$$\begin{aligned}
 (3.15) \quad y_2(t_i) &= g'(t_i) + \frac{(b-a)}{2n} \cdot \left[\frac{\partial f}{\partial t}(t_i, t_0, \bar{x}_1(t_0) + R_{1,0}, \bar{y}_1(t_0) + \omega_{1,0}) \right. \\
 &\quad + 2 \sum_{j=1}^{n-1} \frac{\partial f}{\partial t}(t_i; t_j, \bar{x}_1(t_j) + R_{1,j}, \bar{y}_1(t_j) + \omega_{1,j}) \\
 &\quad \left. + \frac{\partial f}{\partial t}(t_i, t_n, \bar{x}_1(t_n) + R_{1,n}, \bar{y}_1(t_n) + \omega_{1,n}) \right] + \omega_{2,i} \\
 &= g'(t_i) + \frac{(b-a)}{2n} \cdot \left[\frac{\partial f}{\partial t}(t_i, t_0, \bar{x}_1(t_0), \bar{y}_1(t_0)) \right. \\
 &\quad + 2 \sum_{j=1}^{n-1} \frac{\partial f}{\partial t}(t_i; t_j, \bar{x}_1(t_j), \bar{y}_1(t_j)) \\
 &\quad \left. + \frac{\partial f}{\partial t}(t_i, t_n, \bar{x}_1(t_n), \bar{y}_1(t_n)) \right] + \bar{\omega}_{2,i} \\
 &= \bar{y}_2(t_i) + \bar{\omega}_{2,i}, \quad \text{when } i = \overline{0, n}.
 \end{aligned}$$

By induction, for $m \geq 3$, we obtain for $i = \overline{0, n}$ that

$$\begin{aligned}
 (3.16) \quad x_m(t_i) &= g(t_i) + \frac{(b-a)}{2n} \cdot \left[f(t_i, t_0, \bar{x}_{m-1}(t_0) + \bar{R}_{m-1,0}, \bar{y}_{m-1}(t_0) + \bar{\omega}_{m-1,0}) \right. \\
 &\quad \left. + 2 \sum_{j=1}^{n-1} f(t_i; t_j, \bar{x}_{m-1}(t_j) + \bar{R}_{m-1,j}, \bar{y}_{m-1}(t_j) + \bar{\omega}_{m-1,j}) \right. \\
 &\quad \left. + f(t_i, t_n, \bar{x}_{m-1}(t_n) + \bar{R}_{m-1,n}, \bar{y}_{m-1}(t_n) + \bar{\omega}_{m-1,n}) \right]
 \end{aligned}$$

Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)


[Quit](#)

[Page 17 of 24](#)

$$\begin{aligned}
& + 2 \sum_{j=1}^{n-1} f(t_i; t_j, \overline{x_{m-1}}(t_j) + \overline{R}_{m-1,j}, \overline{y_{m-1}}(t_j) + \overline{\omega}_{m-1,j}) \\
& + f(t_i, t_n, \overline{x_{m-1}}(t_n) + \overline{R}_{m-1,n}, \overline{y_{m-1}}(t_n) + \overline{\omega}_{m-1,n}) \Big] + R_{m,i} \\
= & g(t_i) + \frac{(b-a)}{2n} \cdot \left[f(t_i, t_0, \overline{x_{m-1}}(t_0), \overline{y_{m-1}}(t_0)) \right. \\
& + 2 \sum_{j=1}^{n-1} f(t_i; t_j, \overline{x_{m-1}}(t_j), \overline{y_{m-1}}(t_j)) \\
& \left. + f(t_i, t_n, \overline{x_{m-1}}(t_n), \overline{y_{m-1}}(t_n)) \right] + \overline{R}_{m,i} \\
= & \overline{x_m}(t_i) + \overline{R}_{m,i}
\end{aligned}$$

and

$$\begin{aligned}
(3.17) \quad & y_m(t_i) \\
= & g'(t_i) + \frac{(b-a)}{2n} \left[\frac{\partial f}{\partial t}(t_i, t_0, \overline{x_{m-1}}(t_0) + \overline{R}_{m-1,0}, \overline{y_{m-1}}(t_0) \right. \\
& + \overline{\omega}_{m-1,0}) + 2 \sum_{j=1}^{n-1} \frac{\partial f}{\partial t}(t_i; t_j, \overline{x_{m-1}}(t_j) \\
& + \overline{R}_{m-1,j}, \overline{y_{m-1}}(t_j) + \overline{\omega}_{m-1,j})
\end{aligned}$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 18 of 24

$$\begin{aligned}
& + \frac{\partial f}{\partial t} (t_i, t_n, \overline{x_{m-1}}(t_n) + \overline{R}_{m-1,n}, \overline{y_{m-1}}(t_n) + \overline{\omega}_{m-1,n}) \Big] + \omega_{m,i} \\
& = g'(t_i) + \frac{(b-a)}{2n} \cdot \left[\frac{\partial f}{\partial t} (t_i, t_0, \overline{x_{m-1}}(t_0), \overline{y_{m-1}}(t_0)) \right. \\
& \quad + 2 \sum_{j=1}^{n-1} \frac{\partial f}{\partial t} (t_i; t_j, \overline{x_{m-1}}(t_j), \overline{y_{m-1}}(t_j)) \\
& \quad \left. + \frac{\partial f}{\partial t} (t_i, t_n, \overline{x_{m-1}}(t_n), \overline{y_{m-1}}(t_n)) \right] + \overline{\omega}_{m,i} \\
& = \overline{y_m}(t_i) + \overline{\omega}_{m,i}.
\end{aligned}$$

At the remainder estimation we have for any $i = \overline{0, n}$

$$\|R_{1,i}\|_X \leq \frac{L_1(b-a)^2}{4n}, \quad \|\omega_{1,i}\|_X \leq \frac{L_2(b-a)^2}{4n}.$$

Using in (3.14) the Lipschitz property (2.1) we obtain:

$$\|\overline{R}_{2,i}\| \leq [1 + (b-a)(\alpha_1 + \beta_1)] \cdot \frac{L_1(b-a)^2}{4n}, \quad \text{for any } i = \overline{0, n}.$$

Using in (3.15) the Lipschitz property (2.2) we obtain:

$$\|\overline{\omega}_{2,i}\| \leq [1 + (b-a)(\alpha_2 + \beta_2)] \cdot \frac{L_2(b-a)^2}{4n}, \quad \text{for each } i = \overline{0, n}.$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 19 of 24](#)

By induction, we obtain for $m \geq 2$ and $i = \overline{0, n}$,

$$(3.18) \quad \begin{aligned} \|\bar{R}_{m,i}\| &\leq [1 + (b-a)(\alpha_1 + \beta_1) + \cdots + (b-a)^{m-1}(\alpha_1 + \beta_1)^{m-1}] \\ &\quad \cdot \frac{L_1(b-a)^2}{4n} \\ &= \frac{1 - (b-a)^m (\alpha_1 + \beta_1)^m}{1 - (b-a)(\alpha_1 + \beta_1)} \cdot \frac{L_1(b-a)^2}{4n} \end{aligned}$$

and

$$(3.19) \quad \begin{aligned} \|\bar{\omega}_{m,i}\| &\leq [1 + (b-a)(\alpha_2 + \beta_2) + \cdots + (b-a)^{m-1}(\alpha_2 + \beta_2)^{m-1}] \\ &\quad \cdot \frac{L_2(b-a)^2}{4n} \\ &= \frac{1 - (b-a)^m (\alpha_2 + \beta_2)^m}{1 - (b-a)(\alpha_2 + \beta_2)} \cdot \frac{L_2(b-a)^2}{4n}. \end{aligned}$$


Finally, we can state the following result:

Theorem 3.2. *With the conditions (i), (2.1)–(2.7), (2.10), and if $(b-a)(\alpha_1 + \beta_1) < 1$ and $(b-a)(\alpha_2 + \beta_2) < 1$, then the solution of the system (1.2) is approximated on the knots of the uniform partition Δ given in (3.3), by the sequence*

$$\{(\bar{x}_m(t_i), \bar{y}_m(t_i))\}_{m \in \mathbb{N}}, \quad i = \overline{0, n}$$

obtained in (3.12) – (3.17) and the following error estimation holds:

$$(3.20) \quad \left(\begin{array}{c} \|x^*(t_i) - \bar{x}_m(t_i)\|_X \\ \|y^*(t_i) - \bar{y}_m(t_i)\|_X \end{array} \right) \leq Q^m (I_2 - Q)^{-1} d(x_0, x_1)$$



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 20 of 24

$$+ \begin{pmatrix} \frac{L_1(b-a)^2}{4n[1-(b-a)(\alpha_1+\beta_1)]} \\ \frac{L_2(b-a)^2}{4n[1-(b-a)(\alpha_2+\beta_2)]} \end{pmatrix}, \quad \text{for each } m \in \mathbb{N}^* \text{ and } i = \overline{0, n}.$$

Proof. Follows from (2.14), (3.18) and (3.19) since

$$\|x^*(t_i) - \bar{x_m}(t_i)\|_X \leq \|x^*(t_i) - x_m(t_i)\|_X + \|x_m(t_i) - \bar{x_m}(t_i)\|_X$$

for each $m \in \mathbb{N}^*$ and $i = \overline{0, n}$ and,

$$\|y^*(t_i) - \bar{y_m}(t_i)\|_X \leq \|y^*(t_i) - y_m(t_i)\|_X + \|y_m(t_i) - \bar{y_m}(t_i)\|_X$$

for each $m \in \mathbb{N}^*$ and $\forall i = \overline{0, n}$. \square

Remark 1. When $f(t, s, u, v) = H(t, s) \cdot f(s, u, v)$ with $H^1 \in C([a, b]^2, X)$ we obtain the existence, uniqueness and approximation of the solution for Hammerstein-Fredholm integro-differential equations in Banach spaces. Moreover, in the particular case $H(t, s) = G(t, s)$, the Green function, we obtain a new approach for two point boundary value problems associated to second order differential equations in Banach spaces.

Remark 2. For $X = \mathbb{R}^n$ we obtain a new method in analysing systems of Fredholm integro-differential equations and for $X = \mathbb{R}$ we obtain similar results for the approximation of the solution of a scalar Fredholm integro-differential equation.



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back


Close

Quit

Page 21 of 24

References

- [1] S. ANDRAS, A note on Perov's fixed point theorem, *Fixed Point Theory*, **4**(1) (2003), 105–108.
- [2] A. AYAD, Spline approximation for first order Fredholm delay integro-differential equations, *Int. J. Comput. Math.*, **70**(3) (1999), 467–476.
- [3] A. AYAD, Spline approximation for first order Fredholm integro-differential equations, *Studia Univ. Babes-Bolyai Math.*, **41**(3) (1996), 1–8.
- [4] S.H. BEHIRY AND H. HASHISH, Wavelet methods for the numerical solution of Fredholm integro-differential equations, *Int. J. Appl. Math.*, **11**(1) (2002), 27–35.
- [5] S.R. BERNFELD AND V. LAKSHMIKANTHAM, *An Introduction to Nonlinear Boundary Value Problems*, Acad. Press, New York, 1974.
- [6] A. BICA AND S. MUREŞAN, Periodic solutions for a delay integro-differential equations in biomathematics, *RGMIA Res. Report Coll.*, **6**(4) (2003), 755–761.
- [7] A. BICA AND S. MUREŞAN, Applications of the Perov's fixed point theorem to delay integro-differential equations, Chap. 3 in *Fixed Point Theory and Applications* (Y.J. Cho, et al., Eds), Vol. **7**, Nova Science Publishers Inc., New York, 2006.



Application of a Trapezoid Inequality to Neutral Fredholm Integro-Differential Equations in Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 22 of 24

- [8] C. BUŞE, S.S. DRAGOMIR, J. ROUMELIOTIS AND A. SOFO, Generalized trapezoid type inequalities for vector-valued functions and applications, *Math. Inequal. Appl.*, **5**(3) (2002), 435–450.
- [9] V.A. CĂUŞ, Numerical solution of the n-th order Fredholm delay integro-differential equations by spline functions, *Proc. of the Int. Sympos. on Num. Analysis and Approx. Theory*, Cluj-Napoca, Romania, May 9-11, 2002, Cluj Univ. Press, 114–127.
- [10] G. DEZSO, Fixed point theorems in generalized metric spaces, *PUMA Pure Math. Appl.*, **11**(2) (2000), 183–186.
- [11] L.A. GAREY AND C.J. GLADWIN, Direct numerical methods for first order Fredholm integro-differential equations, *Int. J. Comput. Math.*, **34**(3/4) (1990), 237–246.
- [12] S.M. HOSSEINI AND S. SHAHMORAD, Tau numerical solution of Fredholm integro-differential equations with arbitrary polynomial base, *Appl. Math. Modelling*, **27**(2) (2003), 145–154.
- [13] HU QIYA, Interpolation correction for collocation solutions of Fredholm integro-differential equations, *Math. Comput.*, **67**(223) (1998), 987–999.
- [14] L.M. LIHTARNIKOV, Use of Runge-Kutta method for solving Fredholm type integro-differential equations, *Zh. Vychisl. Mat. Fiz.*, **7** (1967), 899–905.
- [15] G. MICULA AND G. FAIRWEATHER, Direct numerical spline methods for first order Fredholm integro-differential equations, *Rev. Anal. Numer. Theor. Approx.*, **22**(1) (1993), 59–66.



Application of a Trapezoid
Inequality to Neutral Fredholm
Integro-Differential Equations in
Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

Title Page

Contents



Go Back

Close

Quit

Page 23 of 24

- [16] A.I. PEROV AND A.V. KIBENKO, On a general method to study the boundary value problems, *Iz. Acad. Nauk.*, **30** (1966), 249–264.
- [17] J. POUR MAHMOUD, M.Y. RAHIMI-ARDABILI AND S. SHAHMORAD, Numerical solution of the system of Fredholm integro-differential equations by the tau method, *Appl. Math. Comput.*, **168**(1) (2005), 465–478.
- [18] I.A. RUS, Fiber Picard operators on generalized metric spaces and an application, *Scripta Sci. Math.*, **1**(2) (1999), 355–363.
- [19] I.A. RUS, A fiber generalized contraction theorem and applications, *Mathematica*, **41**(1) (1999), 85–90.



Application of a Trapezoid
Inequality to Neutral Fredholm
Integro-Differential Equations in
Banach Spaces

Alexandru Mihai Bica,
Vasile Aurel Căuş and
Sorin Mureşan

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 24 of 24