

Journal of Inequalities in Pure and Applied Mathematics

A NOTE ON THE TRACE INEQUALITY FOR PRODUCTS OF HERMITIAN MATRIX POWER

ZHONG PENG YANG AND XIAO XIA FENG


Department of Mathematics,
Putian University,
Putian, Fujian 351100
People's Republic of China.
E-Mail: yangzhongpeng@sina.com

Department of Mathematics,
Beihua University,
Jilin, Jilin 132013,
People's Republic of China.
E-Mail: fengxiaoxia0110@163.com

©2000 School of Communications and Informatics, Victoria University of Technology

ISSN (electronic): 1443-5756

082-02



volume 3, issue 5, article 78,
2002.

*Received 6 June, 2002;
accepted 24 July, 2002.*

Communicated by: D.D. Bainov

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

Da-wei Zhang [J.M.A.A., 237 (1999): 721-725] obtained the inequality $\text{tr}(AB)^{2k} \leq \text{tr} A^{2k} B^{2k}$ for Hermitian matrices A and B , where k is natural number. Here it is proved that these results hold when the power index of the product of Hermitian matrices A and B is a nonnegative even number. In the meantime, it is pointed out that the relation between $\text{tr}(AB)^m$ and $\text{tr} A^m B^m$ is complicated when the power index m is a nonnegative odd number, therefore the above inequality cannot be generalized to all nonnegative integers. As an application, we not only improve the results of Xiaojing Yang [J.M.A.A., 250 (2000), 372-374], Xinmin Yang [J.M.A.A., 263 (2001): 327-333] and Fozil M. Dannan [J. Ineq. Pure and Appl. Math., 2(3) (2001), Art. 34], but also give the complete resolution for the question of the trace inequality about the powers of Hermitian and skew Hermitian matrices that is proposed by Zhengming Jiao.

2000 Mathematics Subject Classification: 15A42; 15A57.


Key words: Hermitian matrix, Trace, Inequality, Skew Hermitian matrix.

The research for this paper was supported by the Science Foundation of the Education Department of Fujian, (No. JB01206).

Contents

1	Introduction	3
2	Some Lemmas	6
3	Main Results	10
4	Trace of the Power on Hermitian Matrix and Skew Hermitian Matrix	19

References



A Note on the Trace Inequality for Products of Hermitian Matrix Power

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 2 of 26

1. Introduction

Let $C^{m \times n}$ be the set of all $n \times n$ matrices over the complex number field \mathbb{C} . The modulus of all diagonal entries of the matrix $A = (a_{ij}) \in C^{n \times n}$ are arranged in decreasing order as $|\delta_1(A)| \geq |\delta_2(A)| \geq \dots \geq |\delta_n(A)|$, i.e., $\delta_1(A), \delta_2(A), \dots, \delta_n(A)$ is an entire arrangement of $a_{11}, a_{22}, \dots, a_{nn}$; all its singular values satisfy $\sigma_1(A) \geq \sigma_2(A) \geq \dots \geq \sigma_n(A)$. In particular, when the eigenvalues of A are real numbers, let its eigenvalues satisfy $\lambda_1(A) \geq \lambda_2(A) \dots \geq \lambda_n(A)$; A^H , $\text{tr } A$ denote its conjugate transpose matrix and trace respectively. Further, let $H(n)$, $H_0^+(n)$, $H^+(n)$, $S(n)$ be the subsets of all *Hermitian*, *Hermitian* semi-positive definite, *Hermitian* positive definite and skew *Hermitian* matrices. Finally, let $A^{1/2}$ represent the quadratic root of $A \in H_0^+(n)$, and \mathbb{R} , \mathbb{N} denote the sets of all real numbers and nonnegative integers. The complex number $\sqrt{-1} \in C$ satisfies $(\sqrt{-1})^2 = -1$.

Recently the trace inequality of two powered *Hermitian* matrices was given in [1] as follows:


$$(1.1) \quad \text{tr}(AB)^{2k} \leq \text{tr } A^{2k} B^{2k}, \quad A, B \in H(n), \quad k \in \mathbb{N}.$$

Furthermore, the following two results were proved in [2],

$$(1.2) \quad 0 \leq \text{tr}(AB)^{2m} \leq (\text{tr } A)^2 (\text{tr } A^2)^{m-1} (\text{tr } B^2)^m, \quad m(\geq 1) \in \mathbb{N}, \quad A, B \in H_0^+(n);$$

and

$$(1.3) \quad 0 \leq \text{tr}(AB)^{2m+1} \leq \text{tr } A \text{tr } B (\text{tr } A^2)^m (\text{tr } B^2)^m, \quad m(\geq 1) \in \mathbb{N}, \quad A, B \in H_0^+(n).$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 3 of 26

Another two results appeared in [3, Theorem 1] and [4, Theorem 1]. When $A, B \in H^+(n)$, the following inequalities hold:

$$(1.4) \quad \operatorname{tr}(AB)^m \leq (\operatorname{tr} A^{2m})^{1/2} (\operatorname{tr} B^{2m})^{1/2}, \quad m \in \mathbb{N};$$

and

$$(1.5) \quad \operatorname{tr}(AB)^m \leq (\operatorname{tr} AB)^m, \quad m \in \mathbb{N}.$$

The above two results (1.1), (1.2), (1.3), (1.4) and (1.5) are related to the work of Bellman. In 1980, Bellman [5] proved:

$$(1.6) \quad \operatorname{tr}(AB)^2 \leq \operatorname{tr} A^2 B^2, \quad A, B \in H_0^+(n),$$


and proposed the conjecture whether

$$(1.7) \quad \operatorname{tr}(AB)^m \leq \operatorname{tr} A^m B^m, \quad m \in \mathbb{N}, \quad A, B \in H_0^+(n)$$

holds.

Since then, many authors have proved that the conjecture (1.7) is correct. In [6], it was pointed out that the inequality (1.7) was also proposed by Lieb and Thiring in 1976, and a similar inequality was proposed also in [7]. R.A. Brualdi [8] commented further work of the inequality (1.7) that was constructed by Lieb and Thiring in [6] and [7].

Whether or not the inequality (1.7) was a conjecture at that time, the condition in [1] was different from that in [2] – [7], which dropped the demand of “semi-positive definite property” for matrices in [1], and examined the trace



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 4 of 26


inequality on the general *Hermitian* matrix powers. Of course, it increases inevitably the discussed difficulty. In 1992 Zhengming Jiao [9] generalised inequality (1.6) for $A \in H(n)$, $B \in S(n)$ and $A, B \in S(n)$, and also presented two questions as follows:

$$(1.8) \quad \operatorname{tr}(AB)^m \geq \operatorname{tr} A^m B^m, \quad A \in H(n), \quad B \in S(n), \quad m \in \mathbb{N}?$$

and

$$(1.9) \quad \operatorname{tr}(AB)^m \leq \operatorname{tr} A^m B^m, \quad A, B \in S(n), \quad m \in \mathbb{N}?$$

We will prove that the inequality (1.1) holds when the power index is a non-negative even number. Thereby the results in [2, 3] and [4] can be obtained and improved. Moreover, a simpler proof for the inequality (1.7) may be presented. As an application of the obtained result, we answer completely two questions mentioned in [9] in the form of (1.8) and (1.9).



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 5 of 26

2. Some Lemmas

Lemma 2.1. Let $A, B \in C^{n \times n}$, then

$$(2.1) \quad \sum_{i=1}^t |\delta_i((AB)^m)| \leq \sum_{i=1}^t \lambda_i((A^H A B B^H)^m) \\ \leq \sum_{i=1}^t \lambda_i((A^H A)^m (B B^H)^m), \quad 1 \leq t \leq n, \quad m \in \mathbb{N}.$$

Proof. From [10, Theorem 8.9], it follows that


$$(2.2) \quad \sum_{i=1}^t |\delta_i(F)| \leq \sum_{i=1}^t \sigma_i(F), \quad 1 \leq t \leq n, \quad F \in C^{n \times n}$$

and by [11, Theorem 1],

$$(2.3) \quad \sum_{i=1}^t \sigma_i \left(\prod_{j=1}^p G_j \right) \\ \leq \sum_{i=1}^t \prod_{j=1}^p \sigma_i(G_j), \quad 1 \leq t \leq n, \quad G_1, G_2, \dots, G_p \in C^{m \times n}.$$

Moreover via [7, Theorem 4], it is derived that

$$(2.4) \quad \sum_{i=1}^t \lambda_i^m(FG) \leq \sum_{i=1}^t \lambda_i(F^m G^m), \quad 1 \leq t \leq n, \quad m \in \mathbb{N}, \quad F, G \in H_0^+(n).$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 6 of 26

therefore through $A^H A, BB^H \in H_0^+(n)$, it is known that all the eigenvalues of $A^H ABB^H \in C^{m \times n}$ are real. Meanwhile

$$\lambda_i \left((A^H ABB^H)^m \right) = \lambda_i^m (A^H ABB^H) = (\lambda_i (A^H ABB^H))^m, \quad i = 1, 2, \dots, n;$$


and from (2.2), (2.3), and (2.4), it holds that

$$\begin{aligned} \sum_{i=1}^t |\delta_i \left((AB)^{2m} \right)| &\leq \sum_{i=1}^t \sigma_i \left((AB)^{2m} \right) \\ &\leq \sum_{i=1}^t (\sigma_i(AB))^{2m} \\ &= \sum_{i=1}^t (\sigma_i^2(AB))^m \\ &= \sum_{i=1}^t \lambda_i \left((A^H ABB^H)^m \right) \\ &\leq \sum_{i=1}^t \lambda_i \left((A^H A)^m (BB^H)^m \right) \end{aligned}$$

that is, (2.1) holds. □

It is well known that eigenvalues of the product AB for $A, B \in H(n)$ are not real numbers, but we can obtain the following lemma.

Lemma 2.2. *Let $A = (a_{ij}), B = (b_{ij}) \in H(n)(S(n))$, then $\text{tr } AB \in \mathbb{R}$.*



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 7 of 26

Proof. When $A, B \in H(n)$, according to [10, pp. 219] it is known that $\text{tr } AB \in \mathbb{R}$.

By the simple fact,

$$(2.5) \quad F \in S(n) \text{ if and only if } \sqrt{-1}F \in H(n),$$

it follows that $\sqrt{-1}A, \sqrt{-1}B \in H(n)$ holds when $A, B \in S(n)$. Thus from the proved result, at this time, it is easy to know

$$\text{tr } AB = \text{tr} \left(-(\sqrt{-1}A)(\sqrt{-1}B) \right) = -\text{tr} \left(\sqrt{-1}A \right) \left(\sqrt{-1}B \right) \in \mathbb{R}.$$

□

By [10, Theorem 6.5.3], the following Lemma holds.

Lemma 2.3. *Let $A, B \in H_0^+(n)$, then*

$$(2.6) \quad 0 \leq \text{tr } AB \leq \text{tr } A \text{tr } B;$$

and

$$(2.7) \quad 0 \leq \text{tr } A^m \leq (\text{tr } A)^m, \quad m \in \mathbb{N}.$$

Lemma 2.4. *Let $A, B \in H(n)$, then $\text{tr}(AB)^m, \text{tr } A^m B^m \in \mathbb{R}$ for all $m \in \mathbb{N}$.*

Proof. When $m = 0, 1$, obviously the result holds by Lemma 2.2. When $m \geq 2$, via

$$\left((AB)^{m-1} A \right)^H = \left(A(BAB \cdots BAB)A \right)^H = (AB)^{m-1} A \in H(n),$$

and Lemma 2.2, it follows that $\text{tr}(AB)^m = \text{tr} \left((AB)^{m-1} A \right) B \in \mathbb{R}$.

For $A^m, B^m \in H(n)$ and from Lemma 2.2, it may be surmised that $\text{tr } A^m B^m \in \mathbb{R}$. □



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents




Go Back

Close

Quit

Page 8 of 26



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 9 of 26

Lemma 2.5. Let $A \in H(n)$, $B \in S(n)$, $m \in \mathbb{N}$, then

$$(2.8) \quad \begin{aligned} \operatorname{tr}(AB)^m &= (-\sqrt{-1})^m \operatorname{tr} (A(\sqrt{-1}B))^m, \\ \operatorname{tr} A^m B^m &= (-\sqrt{-1})^m \operatorname{tr} A^m (\sqrt{-1}B)^m; \end{aligned}$$

and for $m = 2t$ ($t \in \mathbb{N}$), $\operatorname{tr}(AB)^m$, $\operatorname{tr} A^m B^m$ are all real. Further, when $m = 2t + 1$ ($t \in \mathbb{N}$), $\operatorname{tr}(AB)^m$, $\operatorname{tr} A^m B^m$ are all zeros or pure imaginary numbers.

Proof. Without loss of generality, assume that $m \geq 2$, similarly

$$\begin{aligned} \operatorname{tr}(AB)^m &= \operatorname{tr} (A (-\sqrt{-1}(\sqrt{-1}B)))^m = (-\sqrt{-1})^m \operatorname{tr} (A (\sqrt{-1}B))^m, \\ \operatorname{tr} A^m B^m &= \operatorname{tr} A^m (-\sqrt{-1}(\sqrt{-1}B))^m = (-\sqrt{-1})^m \operatorname{tr} A^m (\sqrt{-1}B)^m, \end{aligned}$$

so that (2.8) holds.

When $m = 2t$ ($t \in \mathbb{N}$), $(-\sqrt{-1})^m = (-1)^{3t} \in \mathbb{R}$, thus by (2.8) and Lemma 2.4, one obtains that $\operatorname{tr}(AB)^m$, $\operatorname{tr} A^m B^m$ are all real. When $m = 2t + 1$ ($t \in \mathbb{N}$), $(-\sqrt{-1})^m = (-1)^{3t+1} \sqrt{-1} \notin \mathbb{R}$, then we have that $\operatorname{tr}(AB)^m$, $\operatorname{tr} A^m B^m$ are all zeros or pure imaginary numbers by (2.8) and Lemma 2.4. \square

Similar to the proof of Lemma 2.5, it also follows that:

Lemma 2.6. Let $A, B \in S(n)$, $m \in \mathbb{N}$, then

$$(2.9) \quad \begin{aligned} \operatorname{tr}(AB)^m &= (-1)^m \operatorname{tr} ((\sqrt{-1}A) (\sqrt{-1}B))^m, \\ \operatorname{tr} A^m B^m &= (-1)^m \operatorname{tr} (\sqrt{-1}A)^m (\sqrt{-1}B)^m \in \mathbb{R}. \end{aligned}$$

3. Main Results


Theorem 3.1. Let $A, B \in C^{m \times n}$, then

$$(3.1) \quad |\operatorname{tr}(AB)^{2m}| \leq \operatorname{tr}(A^H A B B^H)^m \leq \operatorname{tr}(A^H A)^m (B B^H)^m, \quad m \in \mathbb{N};$$

$$(3.2) \quad \begin{aligned} |\operatorname{tr}(AB)^{2m}| &\leq \operatorname{tr}(A^H A B B^H)^m \\ &\leq \operatorname{tr}(A^H A)^m (B B^H)^m \\ &\leq \operatorname{tr}\left((A^H A)^{1/2}\right)^2 \operatorname{tr}(A^H A)^{m-1} \operatorname{tr}(B B^H)^m \\ &\leq \left(\operatorname{tr}(A^H A)^{1/2}\right)^2 (\operatorname{tr} A^H A)^{m-1} (\operatorname{tr} B B^H)^m, \\ &\quad m(\geq 1) \in \mathbb{N}. \end{aligned}$$

Proof. Take $t = n$ in (2.1), then we have that

$$\begin{aligned} |\operatorname{tr}(AB)^{2m}| &= \left| \sum_{i=1}^n \delta_i((AB)^{2m}) \right| \\ &\leq \sum_{i=1}^n |\delta_i((AB)^{2m})| \\ &\leq \sum_{i=1}^n \lambda_i((A^H A B B^H)^m) \\ &= \operatorname{tr}(A^H A B B^H)^m \\ &\leq \sum_{i=1}^n \lambda_i((A^H A)^m (B B^H)^m) = \operatorname{tr}((A^H A)^m (B B^H)^m), \end{aligned}$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 10 of 26

giving (3.1).

By $A^H A, BB^H \in H_0^+(n)$ and (2.6), (2.7), when $m \geq 1$ one can get

$$\begin{aligned} \operatorname{tr} (A^H A)^m (BB^H)^m &\leq \operatorname{tr} (A^H A) (A^H A)^{m-1} \operatorname{tr} (BB^H)^m \\ &\leq \operatorname{tr} \left((A^H A)^{1/2} \right)^2 \operatorname{tr} (A^H A)^{m-1} \operatorname{tr} (BB^H)^m \\ &\leq \left(\operatorname{tr} (A^H A)^{1/2} \right)^2 \operatorname{tr} (A^H A)^{m-1} (\operatorname{tr} BB^H)^m, \end{aligned}$$

therefore (3.2) is correct, by (3.1). \square

Theorem 3.2. Let $A, B \in H(n)$, then


$$(3.3) \quad \operatorname{tr}(AB)^{2m} \leq |\operatorname{tr}(AB)^{2m}| \leq \operatorname{tr} (A^2 B^2)^m \leq \operatorname{tr} A^{2m} B^{2m}, \quad m \in \mathbb{N}.$$

Proof. From Lemma 2.4, it is obtained that $\operatorname{tr}(AB)^m, \operatorname{tr} A^m B^m$ are all real, then by $\operatorname{tr}(AB)^{2m} \leq |\operatorname{tr}(AB)^{2m}|$ and $A^H = A, B^H = B$, moreover applying (3.1), (3.3) holds. \square

Because 2^k ($k \in \mathbb{N}$) is a nonnegative even number, conclusion (1.1) in [1] can be achieved by (3.3). When $A, B \in H_0^+(n)$, the inequality (1.7) is obtained by replacing A, B of (3.3) with $A^{1/2}, B^{1/2}$. The above procedure indicates we can give a simple proof for the inequality (1.7).

Example 3.1. Let

$$A = \begin{bmatrix} 1 & -2 \\ -2 & -2 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \in H(2),$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 11 of 26

and so

$$(AB)^3 = \begin{bmatrix} 16 & -6 \\ 4 & 12 \end{bmatrix}, \quad A^3B^3 = \begin{bmatrix} -29 & 44 \\ -26 & 32 \end{bmatrix},$$
$$(AB)^5 = \begin{bmatrix} -56 & -60 \\ 40 & -96 \end{bmatrix}, \quad A^5B^5 = \begin{bmatrix} -481 & 765 \\ -610 & 954 \end{bmatrix},$$

giving

$$\operatorname{tr}(AB)^3 = 28 > 3 = \operatorname{tr} A^3B^3, \quad \operatorname{tr}(AB)^5 = -152 < 573 = \operatorname{tr} A^5B^5.$$

Example 3.2. Let

$$A = \begin{bmatrix} -1 & 2 \\ 2 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \in H(2),$$


giving

$$(AB)^3 = \begin{bmatrix} -16 & 6 \\ -4 & -12 \end{bmatrix}, \quad A^3B^3 = \begin{bmatrix} 29 & -44 \\ 26 & -32 \end{bmatrix},$$

and so we obtain

$$\operatorname{tr}(AB)^3 = -28 < -3 = \operatorname{tr} A^3B^3.$$

We have generalized the index of the trace inequality (1.1) on Hermitian matrix power ([1, Theorem 1]) from $2^k (k \in \mathbb{N})$ to nonnegative even numbers. Examples 3.1 and 3.2 indicate that it is complex when the power index is a positive odd number. Of course, they also show that the result cannot hold when



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 12 of 26

one gives up the “positive semi-definite” requirement of (1.7). In [12, Theorem 6.3.2], the statement “Marcus (1956) generalized this theorem as following form

$$\operatorname{tr}(AB)^m \leq \operatorname{tr} A^m B^m, \quad m \in \mathbb{N}, \quad A, B \in H(n),”$$

follows the proof of (1.6).

Although we do not have access to the article of Marcus, Examples 3.1 and 3.2 make us unsure that the generalization is not correct.

Theorem 3.3. *Let $A, B \in H(n)$, $m \in \mathbb{N}$, then*

$$(3.4) \quad \operatorname{tr}(AB)^{2m} \leq \operatorname{tr}(A^2 B^2)^m \leq \operatorname{tr} A^{2m} B^{2m} \leq (\operatorname{tr} A^{4m})^{1/2} (\operatorname{tr} B^{4m})^{1/2};$$


$$(3.5) \quad \operatorname{tr}(AB)^{2m} \leq \operatorname{tr}(A^2 B^2)^m \leq (\operatorname{tr} A^2 B^2)^m;$$

$$(3.6) \quad \begin{aligned} \operatorname{tr}(AB)^{2m} &\leq \operatorname{tr}(A^2 B^2)^m \\ &\leq \operatorname{tr} A^{2m} B^{2m} \\ &\leq \operatorname{tr} A^2 \operatorname{tr} A^{2(m-1)} \operatorname{tr} B^{2m} \\ &\leq \left(\operatorname{tr}(A^2)^{1/2}\right)^2 (\operatorname{tr} A^2)^{m-1} (\operatorname{tr} B^2)^m, \quad \text{when } m \geq 1. \end{aligned}$$

Proof. From [10, Problem 7.2.10], it is known that

$$(3.7) \quad |\operatorname{tr} FG| \leq (\operatorname{tr} F^2)^{1/2} (\operatorname{tr} G^2)^{1/2}, \quad F, G \in H(n).$$

Thus by (3.3), (3.7) and Lemma 2.4, the inequality (3.4) results.



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents




Go Back

Close

Quit

Page 13 of 26



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 14 of 26

Notice that $AB^2A \in H_0^+(n)$ and $(A^2B^2)^m = A(AB^2A)^{m-1}(AB^2)$, furthermore through (2.7), it follows that

$$\operatorname{tr} (A^2B^2)^m = \operatorname{tr} (AB^2A)^m \leq (\operatorname{tr} AB^2A)^m = (\operatorname{tr} A^2B^2)^m,$$

the inequality (3.5) then follows by (3.3).

Notice that $A^2, (A^2)^{1/2} \in H_0^+(n)$, then via (2.6) and (2.7), it follows that

$$\operatorname{tr} A^{2m} B^{2m} \leq \operatorname{tr} A^2 \operatorname{tr} A^{2(m-1)} \operatorname{tr} B^{2m} \leq \left(\operatorname{tr} (A^2)^{1/2} \right)^2 (\operatorname{tr} A^2)^{m-1} (\operatorname{tr} B^2)^m,$$

the inequality (3.6) then results by using (3.5). □

Corollary 3.4. *Let $A, B \in H_0^+(n)$, $m \in \mathbb{N}$, then the inequalities (1.4) and (1.5) hold. Moreover when $m \geq 1$, it follows that*

$$\begin{aligned} (3.8) \quad 0 &\leq \operatorname{tr}(AB)^{2m} \\ &\leq \operatorname{tr} (A^2B^2)^m \\ &\leq \operatorname{tr} A^{2m} B^{2m} \\ &\leq \operatorname{tr} A^2 \operatorname{tr} A^{2(m-1)} \operatorname{tr} B^{2m} \\ &\leq (\operatorname{tr} A)^2 (\operatorname{tr} A^2)^{m-1} (\operatorname{tr} B^2)^m; \end{aligned}$$


$$\begin{aligned} (3.9) \quad 0 &\leq \operatorname{tr}(AB)^{2m+1} \\ &\leq \operatorname{tr} A^{2m+1} B^{2m+1} \\ &\leq \operatorname{tr} A \operatorname{tr} B \operatorname{tr} A^{2m} \operatorname{tr} B^{2m} \\ &\leq \operatorname{tr} A \operatorname{tr} B (\operatorname{tr} A^2)^m (\operatorname{tr} B^2)^m. \end{aligned}$$

Proof. Using $A^{1/2}$, $B^{1/2}$ instead of A , B in the inequalities (3.4) and (3.5), (1.4) and (1.5) can be obtained. From $A \in H_0^+(n)$ it follows that $(A^2)^{1/2} = A$ and (3.8) is derived by (3.6). Furthermore through (3.6), (3.8), (2.6) and (2.7), it holds that

$$\begin{aligned}
 0 &\leq \operatorname{tr}(AB)^{2m+1} \\
 &= \operatorname{tr}\left(\left(A^{1/2}\right)^2\left(B^{1/2}\right)^2\right)^{2m+1} \\
 &\leq \operatorname{tr}\left(A^{1/2}\right)^{2(2m+1)}\left(B^{1/2}\right)^{2(2m+1)} \\
 &= \operatorname{tr}A^{2m+1}B^{2m+1} \\
 &\leq \operatorname{tr}A^{2m+1}\operatorname{tr}B^{2m+1} \\
 &= \operatorname{tr}AA^{2m}\operatorname{tr}BB^{2m} \\
 &\leq \operatorname{tr}A\operatorname{tr}A^{2m}\operatorname{tr}B\operatorname{tr}B^{2m} \\
 &= \operatorname{tr}A\operatorname{tr}B\operatorname{tr}A^{2m}\operatorname{tr}B^{2m} \\
 &\leq \operatorname{tr}A\operatorname{tr}B\left(\operatorname{tr}A^2\right)^m\left(\operatorname{tr}B^2\right)^m,
 \end{aligned}$$

giving (3.9). □

As an application of the main results, in Corollary 3.4, the basic conclusions (1.2), (1.3), (1.4) and (1.5) of [2, 3] and [4] are summarized. At the same time, in Theorem 3.3, it is shown that the trace inequality (1.2) ([2]) on semi-positive definite *Hermitian* matrix power is extended to general *Hermitian* matrix, according to the form (3.6). By following Example 3.3, it is indicated that the trace inequality (1.3) ([2]) on semi-positive definite *Hermitian* matrix power cannot be generalised in a similar fashion to Theorem 3.3.



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 15 of 26

Example 3.3. *Let*

$$A = \begin{bmatrix} -1 & 2 \\ 2 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix} \in H(2),$$

then from Examples 3.1 and 3.2, it is shown that

$$(AB)^5 = \begin{bmatrix} 56 & 60 \\ -40 & 96 \end{bmatrix}, \quad A^2 = \begin{bmatrix} 5 & 2 \\ 2 & 8 \end{bmatrix}, \quad B^2 = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}.$$

Thus it is easily seen that

$$\begin{aligned} \operatorname{tr}(AB)^5 &= \operatorname{tr}(AB)^{2 \times 2 + 1} \\ &= 152 > 1 \times (-1) \times 13^2 \times 3^2 \\ &= \operatorname{tr} A \operatorname{tr} B (\operatorname{tr} A^2)^2 (\operatorname{tr} B^2)^2. \end{aligned}$$

Example 3.4. *Let*


$$A = \begin{bmatrix} 2 & 3 \\ 3 & -4 \end{bmatrix}, \quad B = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$$

and so from

$$AB = \begin{bmatrix} 6 & 6 \\ 9 & -8 \end{bmatrix}, \quad (AB)^2 = \begin{bmatrix} 90 & -12 \\ -18 & 118 \end{bmatrix},$$

it follows that

$$\operatorname{tr}(AB)^2 = 208 > 4 = (\operatorname{tr} AB)^2.$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 16 of 26

Example 3.4 indicates that the result (1.5) ([4]) for positive definite matrix cannot be generalized to general *Hermitian* matrix, but the generalized form, similar to (3.5) in Theorem 3.3, may be obtained.

Example 3.5. *Let*

$$A = \begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix} \in H(2),$$

then from $A^2 = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = 2I$, $B^2 = 5I$, it follows that

$$(\operatorname{tr}(A^2B^2))^2 = 400 > 200 = \operatorname{tr} A^4B^4.$$

Example 3.6. *Let*

$$A = \begin{bmatrix} 3 & -1 \\ -1 & -1 \end{bmatrix}, \quad B = \begin{bmatrix} 5 & 2 \\ 2 & -1 \end{bmatrix} \in H(2),$$


from

$$A^2B^2 = \begin{bmatrix} 274 & 70 \\ -42 & -6 \end{bmatrix}, \quad A^4B^4 = \begin{bmatrix} 87592 & 26152 \\ -19544 & -5816 \end{bmatrix},$$

it is achieved that

$$(\operatorname{tr}(A^2B^2))^2 = 71824 < 81776 = \operatorname{tr} A^4B^4.$$

Examples 3.5 and 3.6 show that the two upper bounds $\operatorname{tr} A^{2m}B^{2m}$ and $(\operatorname{tr} A^2B^2)^m$ for $\operatorname{tr}(AB)^{2m}$ as given by (3.3) and (3.5), are independent of each



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close


Quit

Page 17 of 26

other, in which $\text{tr}(AB)^{2m}$ is the trace of the product power on the two *Hermitian* matrices A and B . The result of [4] can be derived by replacing the matrices A and B in Examples 3.5 and 3.6 with $A^{1/2}$ and $B^{1/2}$.

For the trace $\text{tr}(AB)^m$ of the product power on positive definite *Hermitian* matrices A and B , the upper bounds $\text{tr} A^m B^m$ and $(\text{tr} AB)^m$ given by (1.7) and (1.5) cannot be compared with each other; but from the upper bound $(\text{tr} A^{2m})^{1/2} (\text{tr} B^{2m})^{1/2}$ ([3]) determined by (1.4), via (3.4) and Corollary 3.4, it follows that

$$\text{tr} A^m B^m \leq (\text{tr} A^{2m})^{1/2} (\text{tr} B^{2m})^{1/2} .$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 18 of 26

4. Trace of the Power on Hermitian Matrix and Skew Hermitian Matrix

Theorem 4.1. Let $A \in H(n)$, $B \in S(n)$, then when $m = 4t$ or $m = 4t + 2$, $t \in \mathbb{N}$, $\operatorname{tr}(AB)^m$ and $\operatorname{tr} A^m B^m$ are all real numbers, and


$$(4.1) \quad \operatorname{tr}(AB)^m \leq \operatorname{tr}(A^2 B^2)^{m/2} \leq \operatorname{tr} A^m B^m, \quad m = 4t, \quad t \in \mathbb{N};$$

$$(4.2) \quad \operatorname{tr}(AB)^m \geq \operatorname{tr}(A^2 B^2)^{m/2} \geq \operatorname{tr} A^m B^m, \quad m = 4t + 2, \quad t \in \mathbb{N};$$

similarly when $m = 4t + 1$ or $m = 4t + 3$, $t \in \mathbb{N}$, if $\operatorname{tr}(AB)^m \neq 0$ or $\operatorname{tr} A^m B^m \neq 0$, then $\operatorname{tr}(AB)^m \notin \mathbb{R}$ or $\operatorname{tr} A^m B^m \notin \mathbb{R}$, so $\operatorname{tr}(AB)^m$ and $\operatorname{tr} A^m B^m$ cannot be compared with each other.

Proof. By Lemma 2.5, we have that both $\operatorname{tr}(AB)^m$ and $\operatorname{tr} A^m B^m$ are real numbers when $m = 4t$. Furthermore through (3.3), (2.8), it follows that

$$\begin{aligned} \operatorname{tr}(AB)^m &= (-\sqrt{-1})^{4t} \operatorname{tr}(A(\sqrt{-1}B))^{4t} \\ &= \operatorname{tr}(A(\sqrt{-1}B))^{4t} \\ &\leq \operatorname{tr}(A^2(\sqrt{-1}B)^2)^{2t} \\ &= \sqrt{-1}^{4t} \operatorname{tr}(A^2 B^2)^{2t} \\ &= \operatorname{tr}(A^2 B^2)^{m/2} \\ &\leq \operatorname{tr} A^{4t} (\sqrt{-1}B)^{4t} \\ &= \sqrt{-1}^{4t} \operatorname{tr} A^{4t} B^{4t} \\ &= \operatorname{tr} A^m B^m, \end{aligned}$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 19 of 26

giving (4.1).

In the same way, when $m = 4t + 2$, $\text{tr}(AB)^m$ and $\text{tr} A^m B^m$ are all real numbers and it holds that


$$\begin{aligned}
 \text{tr}(AB)^m &= (-\sqrt{-1})^{4t+2} \text{tr} (A (\sqrt{-1}B))^{4t+2} \\
 &= -\text{tr} (A (\sqrt{-1}B))^{2(2t+1)} \\
 &\geq -\text{tr} (A^2 (\sqrt{-1}B)^2)^{2t+1} \\
 &= -\sqrt{-1}^{4t+2} \text{tr} (A^2 (\sqrt{-1}B)^2)^{2t+1} \\
 &= \text{tr} (A^2 B^2)^{m/2} \\
 &\geq -\text{tr} A^{2(2t+1)} (\sqrt{-1}B)^{2(2t+1)} \\
 &= -\sqrt{-1}^{4t+2} \text{tr} A^{4t+2} B^{4t+2} = \text{tr} A^m B^m,
 \end{aligned}$$

producing (4.2).

When $m = 4t + 1$ or $m = 4t + 3$, $t \in N$, the result is obtained by Lemma 2.5. \square

Example 4.1. Let

$$\begin{aligned}
 A &= \begin{bmatrix} -2 & 1 \\ 1 & -1 \end{bmatrix} \in H(2), \\
 B &= \begin{bmatrix} 2\sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{bmatrix} = \sqrt{-1} \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} \in S(2).
 \end{aligned}$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 20 of 26

Hence

$$(AB)^3 = \sqrt{-1} \begin{bmatrix} 50 & 11 \\ -22 & -5 \end{bmatrix} \quad \text{and} \quad A^3B^3 = \sqrt{-1} \begin{bmatrix} 104 & 8 \\ -64 & -5 \end{bmatrix}.$$

It is known that both $\text{tr}(AB)^3$ and $\text{tr} A^3B^3$ are pure imaginary numbers by Lemma 2.5, they cannot be compared with each other, but their imaginary parts have the following relation

$$\text{Im tr}(AB)^3 = 45 < 99 = \text{Im tr} A^3B^3.$$

Let

$$C = \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix} \in H(2),$$

$$D = \sqrt{-1} \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \in S(2).$$

$$(CD)^3 = \sqrt{-1} \begin{bmatrix} 15 & -42 \\ 70 & 29 \end{bmatrix},$$


$$C^3D^3 = \sqrt{-1} \begin{bmatrix} 15 & 30 \\ -64 & -35 \end{bmatrix},$$

and hence

$$\text{Im tr}(CD)^3 = 44 > -20 = \text{Im tr} C^3D^3.$$

Example 4.2. Let

$$A = \begin{bmatrix} 1 & -2 \\ -2 & -1 \end{bmatrix} \in H(2), \quad B = \sqrt{-1} \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \in S(2).$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 21 of 26

Hence

$$(AB)^5 = \sqrt{-1} \begin{bmatrix} -435 & 543 \\ -905 & -616 \end{bmatrix} \quad \text{and} \quad A^5 B^5 = \sqrt{-1} \begin{bmatrix} 525 & 375 \\ -1775 & -670 \end{bmatrix}.$$

Furthermore their imaginary parts satisfy:

$$\text{Im tr}(AB)^5 = -1051 < -145 = \text{Im tr } A^5 B^5.$$

Let $C = -A \in H(2)$, $D = B \in S(2)$, then

$$\text{Im tr}(CD)^5 = -\text{Im tr}(AB)^5 = 1051 > 145 = -\text{Im tr } A^5 B^5 = \text{Im tr } C^5 D^5.$$

From Example 4.1 and 4.2, it is known that when $m = 4t + 1$ or $m = 4t + 3$, $t \in N$, in general, the imaginary parts of $\text{tr}(AB)^m$ and $\text{tr } A^m B^m$ as pure imaginary numbers make the positive and reverse direction of the question (1.8) not to hold. It is seen from this that in Theorem 4.1, the question (1.8) that is proposed by [9] is completely resolved.


Theorem 4.2. Let $A, B \in S(n)$, if $m = 2t$, $t \in N$, then both $\text{tr}(AB)^m$ and $\text{tr } A^m B^m$ are real numbers and

$$(4.3) \quad \text{tr}(AB)^m \leq |\text{tr}(AB)^m| \leq \text{tr}(A^2 B^2)^{m/2} \leq \text{tr } A^m B^m,$$

holds.

Proof. By (2.5) and Lemma 2.6, it is known that both $\text{tr}(AB)^m$ and $\text{tr } A^m B^m$ are real numbers and

$$(4.4) \quad \begin{aligned} \text{tr}(AB)^m &= \text{tr}((\sqrt{-1}A)(\sqrt{-1}B))^m, \\ \text{tr } A^m B^m &= \text{tr}(\sqrt{-1}A)^m (\sqrt{-1}B)^m. \end{aligned}$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 22 of 26

Moreover by (3.3), it follows that

$$\begin{aligned}
 \operatorname{tr}(AB)^m &= \operatorname{tr} \left((\sqrt{-1}A) (\sqrt{-1}B) \right)^{2t} \\
 &\leq \left| \operatorname{tr} \left((\sqrt{-1}A) (\sqrt{-1}B) \right)^{2t} \right| \\
 &= \left| \operatorname{tr}(AB)^m \right| \\
 &\leq \operatorname{tr} \left((\sqrt{-1}A)^2 (\sqrt{-1}B)^2 \right)^t \\
 &= \operatorname{tr} \left(\sqrt{-1}^4 A^2 B^2 \right)^{m/2} \\
 &= \operatorname{tr} \left(A^2 B^2 \right)^{m/2} \\
 &\leq \operatorname{tr} \left(\sqrt{-1}A \right)^{2t} \left(\sqrt{-1}B \right)^{2t} \\
 &= \operatorname{tr} \left(\sqrt{-1}^{4t} A^{2t} B^{2t} \right) = \operatorname{tr} A^m B^m,
 \end{aligned}$$

giving (4.3). □

Example 4.3. *Let*


$$A = \sqrt{-1} \begin{bmatrix} -2 & 1 \\ 1 & -1 \end{bmatrix}, \quad B = \sqrt{-1} \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} \in S(2),$$

and

$$(AB)^3 = - \begin{bmatrix} -51 & -11 \\ 22 & 5 \end{bmatrix}, \quad A^3 B^3 = - \begin{bmatrix} -104 & -8 \\ 64 & 5 \end{bmatrix},$$

so that

$$\operatorname{tr}(AB)^3 = 45 < 99 = \operatorname{tr} A^3 B^3.$$



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 23 of 26

Let

$$C = \sqrt{-1} \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix}, \quad D = \sqrt{-1} \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \in S(2),$$


according to

$$(CD)^3 = \begin{bmatrix} 15 & -42 \\ 70 & 29 \end{bmatrix}, \quad C^3D^3 = \begin{bmatrix} 15 & 30 \\ -130 & -35 \end{bmatrix},$$

it is known that

$$\operatorname{tr}(CD)^3 = 44 > -20 = \operatorname{tr} C^3D^3.$$

Via (2.9), we know that both $\operatorname{tr}(AB)^m$ and $\operatorname{tr} A^m B^m$ are real numbers when $m = 2t + 1$, $t \in \mathbb{N}$ and $A, B \in S(n)$. Thereby in Example 4.3, it is indicated that the positive and reverse direction of the question (1.9) certainly do not hold, in which the question (1.9) is proposed by [9]; when m is a nonnegative even number, the question as posed by (1.9) is confirmed by Theorem 4.2, thus we completely resolve the question (1.9) of [9].



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back


Close

Quit

Page 24 of 26

References

- [1] DA-WEI ZHANG, Note a matrix trace inequality for products of Hermitian matrices, *J. Math. Anal. Appl.*, **237** (1999), 721–725.
- [2] XIAOJING YANG, Note a matrix trace inequality, *J. Math. Anal. Appl.*, **250** (2000), 372–374.
- [3] XINMIN YANG, XIAOQI YANG AND KOK LAI TEO, A matrix trace inequality, *J. Math. Anal. Appl.*, **263** (2001), 327–333.
- [4] F.M. DANNAN, Matrix and operator inequalities, *Ineq. Pure. and Appl. Math.*, **2**(3) (2001), Art. 34. [ONLINE: http://jipam.vu.edu.au/v2n3/029_01.html]
- [5] B. BELLMAN, Some inequalities for positive definite matrices, in *General Inequalities 2*, Proceeding 2nd international conference on general inequalities (E. F. Beckenbach, Ed), Birkhauser, Basel, 1980, 89–90.
- [6] P. J. BUSHHELL AND G. B. TRUSTRUM, Trace inequalities for positive definite matrix power products, *Lin. Alg. Appl.*, **132** (1990), 172–178.
- [7] BO-YING WANG AND MING-PENG GONG, Some eigenvalue inequalities for positive semidefinite matrix power products, *Lin. Alg. Appl.*, **184** (1993), 242–249.
- [8] R.A. BRUALDI, From the Editor-in-Chief, *Lin. Alg. Appl.*, **220** (1995), 1–6.



A Note on the Trace Inequality
for Products of Hermitian Matrix
Power

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents




Go Back

Close

Quit

Page 25 of 26

- [9] ZHENGMING JIAO, On the inequalities for traces of matrix product, *Journal of Shangdong Normal University* (Natural Science Edition), **7**(4) (1992), 19–20.
- [10] FUZHEN ZHANG, *Matrix Theory: Basic and Techniques*, Springer-Verlag, New York, 1999.
- [11] LING CHEN, Inequalities for singular values and trace, *Lin. Alg. Appl.*, **171** (1992), 109–120.
- [12] SONG GUI WANG AND ZHONG ZHEN JIA, *The Inequality in Matrix Theory*, The Educational Publishing House of An’hui, Hefei, 1994.



**A Note on the Trace Inequality
for Products of Hermitian Matrix
Power**

Zhong Peng Yang and
Xiao Xia Feng

Title Page

Contents



Go Back

Close

Quit

Page 26 of 26