


SUBORDINATION RESULTS FOR THE FAMILY OF UNIFORMLY CONVEX p -VALENT FUNCTIONS

H.A. AL-KHARSANI AND S.S. AL-HAJIRY

Department of Mathematics
Faculty of Science
Girls College, Dammam
Saudi Arabia.
EMail: ssmh1@hotmail.com



volume 7, issue 1, article 20,
2006.

*Received 16 May, 2005;
accepted 02 October, 2005.*

Communicated by: G. Kohr

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

The object of the present paper is to introduce a class of p -valent uniformly convex functions UCV_p . We deduce a criteria for functions to lie in the class UCV_p and derive several interesting properties such as distortion inequalities and coefficients estimates. We confirm our results using the Mathematica program by drawing diagrams of extremal functions of this class.

2000 Mathematics Subject Classification: 30C45.


Key words: p -valent, Uniformly convex functions, Subordination.

We express our thanks to Dr.M.K.Aouf for his helpful comments.

Contents

1	Introduction	3
2	The Class PAR_p	4
3	Characterization of UCV_p	6
4	Subordination Theorem and Consequences	9
5	General Properties of Functions in UCV_p	15
	5.1 Remarks	21

References



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 2 of 22

1. Introduction

Denote by $A(p, n)$ the class of normalized functions

$$(1.1) \quad f(z) = z^p + \sum_{k=2}^{\infty} a_{k+p-1} z^{k+p-1}$$

regular in the unit disk $D = \{z : |z| < 1\}$ and $p \in \mathbb{N}$, consider also its subclasses $C(p), S^*(p)$ consisting of p -valent convex and starlike functions respectively, where $C(1) \equiv C, S^*(1) \equiv S^*$, the classes of univalent convex and starlike functions .


It is well known that for any $f \in \mathbb{C}$, not only $f(D)$ but the images of all circles centered at 0 and lying in D are convex arcs. B. Pinchuk posed a question whether this property is still valid for circles centered at other points of D . A.W. Goodman [1] gave a negative answer to this question and introduced the class UCV of univalent uniformly convex functions, $f \in C$ such that any circular arc γ lying in D , having the center $\zeta \in D$ is carried by f into a convex arc. A.W. Goodman [1] stated the criterion

$$(1.2) \quad \operatorname{Re} \left[1 + (z - \zeta) \frac{f''(z)}{f'(z)} \right] > 0, \quad \forall z, \zeta \in D \iff f \in UCV.$$

Later F. Ronning (and independently W. Ma and D. Minda) [7] obtained a more suitable form of the criterion , namely

$$(1.3) \quad \operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] > \left| \frac{zf''(z)}{f'(z)} \right|, \quad \forall z \in D \iff f \in UCV.$$

This criterion was used to find some sharp coefficients estimates and distortion theorems for functions in the class UCV .



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents




Go Back

Close

Quit

Page 3 of 22



2. The Class PAR_p

We now introduce a subfamily PAR_p of P . Let


$$(2.1) \quad \Omega = \left\{ w = \mu + iv : \frac{v^2}{p} < 2\mu - p \right\}$$

$$(2.2) \quad = \{ w : \operatorname{Re} w > |w - p| \}.$$

Note that Ω is the interior of a parabola in the right half-plane which is symmetric about the real axis and has vertex at $(p/2, 0)$. The following diagram shows Ω when $p = 3$:

Let

$$(2.3) \quad PAR_p = \{ h \in p : h(D) \subseteq \Omega \}.$$



Subordination Results for the
Family of Uniformly Convex
 p -valent Functions

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 4 of 22

Example 2.1. It is known that $z = -\tan^2\left(\frac{\pi}{2\sqrt{2p}}\sqrt{w}\right)$ maps

$$\left\{w = \mu + iv : \frac{v^2}{p} < p - 2\mu\right\}$$

conformally onto D . Hence, $z = -\tan^2\left(\frac{\pi}{2\sqrt{2p}}\sqrt{p-w}\right)$ maps Ω conformally onto D . Let $w = Q(z)$ be the inverse function. Then $Q(z)$ is a Riemann mapping function from D to Ω which satisfies $Q(0) = p$; more explicitly,

$$(2.4) \quad Q(z) = p + \frac{2p}{\pi^2} \left(\log \left(\frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right) \right)^2 = \sum_{n=0}^{\infty} B_n z^n$$

$$(2.5) \quad = p + \frac{8p}{\pi^2} z + \frac{16p}{3\pi^2} z^2 + \frac{184p}{45\pi^2} z^3 + \dots$$

Obviously, $Q(z)$ belongs to the class PAR_p . Geometrically, PAR_p consists of those holomorphic functions $h(z)$ ($h(0) = p$) defined on D which are subordinate to $Q(z)$, written $h(z) \prec Q(z)$.


The analytic characterization of the class PAR_p is shown in the following relation:

$$(2.6) \quad h(z) \in PAR_p \Leftrightarrow \operatorname{Re}\{h(z)\} \geq |h(z) - p|$$

such that $h(z)$ is a p -valent analytic function on D .

Now, we can derive the following definition.

Definition 2.1. Let $f(z) \in A(p, n)$. Then $f(z) \in UCV_p$ if $f(z) \in C(p)$ and $1 + z \frac{f''(z)}{f'(z)} \in PAR_p$.



Subordination Results for the
Family of Uniformly Convex
 p -valent Functions

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 5 of 22

3. Characterization of UCV_p

We present the necessary and sufficient condition to belong to the class UCV_p in the following theorem:

Theorem 3.1. *Let $f(z) \in A(p, n)$. Then*

$$(3.1) \quad f(z) \in UCV_p \Leftrightarrow 1 + \operatorname{Re} \left\{ z \frac{f''(z)}{f'(z)} \right\} \geq \left| z \frac{f''(z)}{f'(z)} - (p-1) \right|, \quad z \in D.$$

Proof. Let $f(z) \in UCV_p$ and $h(z) = 1 + z \frac{f''(z)}{f'(z)}$. Then $h(z) \in PAR_p$, that is, $\operatorname{Re}\{h(z)\} \geq |h(z) - p|$. Then

$$\operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} \geq \left| z \frac{f''(z)}{f'(z)} - (p-1) \right|.$$

□


Example 3.1. *We now specify a holomorphic function $K(z)$ in D by*

$$(3.2) \quad 1 + z \frac{K''(z)}{K'(z)} = Q(z),$$

where $Q(z)$ is the conformal mapping onto Ω given in Example 2.1. Then it is clear from Theorem 3.1 that $K(z)$ is in UCV_p .

Let

$$(3.3) \quad K(z) = z^p + \sum_{k=2}^{\infty} A_k z^{k+p-1}.$$



Subordination Results for the Family of Uniformly Convex p -valent Functions

H.A. Al-Kharsani and S.S. Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 6 of 22

From the relationship between the functions $Q(z)$ and $K(z)$, we obtain

$$(3.4) \quad (p + n - 1)(n - 1)A_n = \sum_{k=1}^{n-1} (k + p - 1)A_k B_{n-k}.$$

Since all the coefficients B_n are positive, it follows that all of the coefficients A_n are also positive. In particular,

$$(3.5) \quad A_2 = \frac{8p^2}{\pi^2(p + 1)},$$


and

$$(3.6) \quad A_3 = \frac{p^2}{2(p + 2)} \left(\frac{16}{3\pi^2} + \frac{64p}{\pi^4} \right).$$

Note that

$$(3.7) \quad \log \frac{k'(z)}{z^{p-1}} = \int_0^z \frac{Q(\zeta) - p}{\zeta} d\zeta.$$

By computing some coefficients of $K(z)$ when $p = 3$, we can obtain the following diagram



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents





Go Back

Close

Quit

Page 7 of 22



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 8 of 22

4. Subordination Theorem and Consequences

In this section, we first derive some subordination results from Theorem 4.1; as corollaries we obtain sharp distortion, growth, covering and rotation theorems from the family UCV_p .

Theorem 4.1. Assume that $f(z) \in UCV_p$. Then $1 + z \frac{f''(z)}{f'(z)} \prec 1 + z \frac{K''(z)}{K'(z)}$ and $\frac{f'(z)}{z^{p-1}} \prec \frac{K'(z)}{z^{p-1}}$.

Proof. Let $f(z) \in UCV_p$. Then $h(z) = 1 + z \frac{f''(z)}{f'(z)} \prec 1 + z \frac{K''(z)}{K'(z)}$ is the same as $h(z) \prec Q(z)$. Note that $Q(z) - p$ is a convex univalent function in D . By using a result of Goluzin, we may conclude that


$$(4.1) \quad \log \frac{f'(z)}{z^{p-1}} = \int_0^z \frac{h(\varsigma) - 1}{\varsigma} d\varsigma \prec \int_0^z \frac{Q(\varsigma) - p}{\varsigma} d\varsigma = \log \frac{K'(z)}{z^{p-1}}.$$

Equivalently, $\frac{f'(z)}{z^{p-1}} \prec \frac{K'(z)}{z^{p-1}}$. □

Corollary 4.2 (Distortion Theorem). Assume $f(z) \in UCV_p$ and $|z| = r < 1$. Then $K'(-r) \leq |f'(z)| \leq K'(r)$.

Equality holds for some $z \neq 0$ if and only if $f(z)$ is a rotation of $K(z)$.

Proof. Since $Q(z) - p$ is convex univalent in D , it follows that $\log K'(z)$ is also convex univalent in D . In fact, the power series for $\log K'(z)$ has positive coefficients, so the image of D under this convex function is symmetric about



Subordination Results for the
Family of Uniformly Convex
 p -valent Functions

H.A. Al-Kharsani and S.S.
Al-Hajjry

Title Page

Contents



Go Back

Close

Quit

Page 9 of 22

the real axis. As $\log \frac{f'(z)}{z^{p-1}} \prec \log \frac{K'(z)}{z^{p-1}}$, the subordination principle shows that

$$(4.2) \quad \begin{aligned} K'(-r) &= e^{\{\log K'(-r)\}} = e^{\left\{ \min_{|z|=r} \operatorname{Re}\{\log K'(z)\} \right\}} \\ &\leq e^{\{\operatorname{Re} \log K'(z)\}} = |f'(z)| \leq e^{\left\{ \max_{|z|=r} \operatorname{Re}\{\log K'(z)\} \right\}} \\ &= e^{\{\log K'(r)\}} = K'(r). \end{aligned}$$

Note that for $|z_0| = r$, either

$$\operatorname{Re}\{\log f'(z_0)\} = \min_{|z|=r} \operatorname{Re}\{\log K'(z)\}$$

or

$$\operatorname{Re}\{\log f'(z_0)\} = \max_{|z|=r} \operatorname{Re}\{\log K'(z)\}$$

for some $z_0 \neq 0$ if and only if $\log f'(z) = \log K'(e^{i\theta}z)$ for some $\theta \in R$. \square


Theorem 4.3. *Let $f(z) \in UCV_p$. Then*

$$(4.3) \quad |f'(z)| \leq |z^{p-1}| e^{\frac{14p}{\pi^2}\zeta(3)} = |z^{p-1}| L^p$$

for $|z| < 1$. ($L \approx 5.502$, $\zeta(t)$ is the Riemann Zeta function.)

Proof. Let $\phi(z) = \frac{zg'(z)}{g(z)}$, where $g(z) = zf'(z)$. Then $\phi(z) \prec Q(z)$ which means that $\phi(z) \prec p + \frac{2p}{\pi^2} \left(\log \left(\frac{1+\sqrt{z}}{1-\sqrt{z}} \right) \right)$. Moreover,

$$\log \frac{g(z)}{z^p} = \int_0^z \left(\frac{\phi(s) - p}{s} \right) ds$$



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 10 of 22

and therefore, if $z = re^{i\theta}$ and $|z| = 1$,

$$\begin{aligned} \log \left| \frac{g(z)}{z^p} \right| &= \int_0^r \Re e(\phi(te^{i\theta}) - p) \frac{dt}{t} \leq \frac{2p}{\pi^2} \int_0^r \frac{1}{t} \log \left(\frac{1 + \sqrt{t}}{1 - \sqrt{t}} \right) dt \\ &\leq \frac{2p}{\pi^2} \int_0^1 \frac{1}{t} \log \left(\frac{1 + \sqrt{t}}{1 - \sqrt{t}} \right) dt = \frac{2p}{\pi^2} (7\zeta(3)), \end{aligned}$$

where


$$\int_0^1 \frac{1}{t} \log \left(\frac{1 + \sqrt{t}}{1 - \sqrt{t}} \right) dt = 7\zeta(3) \quad [8].$$

Then we find that

$$\left| \frac{zf'(z)}{z^p} \right| \leq e^{\frac{2p}{\pi^2} (7\zeta(3))}.$$

□

The following diagram shows the boundary of $K(z)$'s derivative when $p = 2$ in a circle has the radius $(5.5)^2$:



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 11 of 22

Corollary 4.4 (Growth Theorem). Let $f(z) \in UCV_p$ and $|z| = r < 1$. Then $-K(-r) \leq |f(z)| \leq K(r)$.

Equality holds for some $z \neq 0$ if and only if $f(z)$ is a rotation of $K(z)$.

Corollary 4.5 (Covering Theorem). Suppose $f(z) \in UCV_p$. Then either $f(z)$ is a rotation of $K(z)$ or $\{w : |w| \leq -K(-1)\} \subseteq f(D)$.

Corollary 4.6 (Rotation Theorem). Let $f(z) \in UCV_p$ and $|z_0| = r < 1$. Then

$$(4.4) \quad |Arg\{f'(z_0)\}| \leq \max_{|z|=r} Arg\{K'(z)\}.$$

Equality holds for some $z \neq 0$ if and only if $f(z)$ is a rotation of $K(z)$.

Theorem 4.7. Let $f(z) = z^p + \sum_{k=2}^{\infty} a_{k+p-1} z^{k+p-1}$ and $f(z) \in UCV_p$, and let $A_{n+p-1} = \max_{f(z) \in UCV_p} |a_{n+p-1}|$. Then

$$(4.5) \quad A_{p+1} = \frac{8p^2}{\pi^2(p+1)}.$$

The result is sharp. Further, we get

$$(4.6) \quad A_{n+p-1} \leq \frac{8p^2}{(n+p-1)(n-1)\pi^2} \prod_{k=3}^n \left(1 + \frac{8p}{(k-2)\pi^2}\right).$$

Proof. Let $f(z) = z^p + \sum_{k=2}^{\infty} a_{k+p-1} z^{k+p-1}$ and $f(z) \in UCV_p$, and define

$$\phi(z) = 1 + \frac{zf''(z)}{f'(z)} = p + \sum_{k=2}^{\infty} c_k z^{k+p-1}.$$



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajjry

Title Page

Contents



Go Back

Close

Quit

Page 12 of 22

Then $\phi(z) \prec Q(z)$. $Q(z)$ is univalent in D and $Q(D)$ is a convex region, so Rogosinski's theorem applies.

$$Q(z) = p + \frac{8p}{\pi^2}z + \frac{16p}{3\pi^2}z^2 + \frac{184p}{45\pi^2}z^3 + \dots,$$

so we have $|c_n| \leq |B_1| = \frac{8p}{\pi^2} := B$. Now, from the relationship between functions $f(z)$ and $Q(z)$, we obtain

$$(n+p-1)(n-1)a_{n+p-1} = \sum_{k=1}^{n-1} (k+p-1)a_{k+p-1}c_{n-k}.$$

From this we get $|a_{p+1}| = \frac{pB}{(p+1)} = \frac{8p^2}{\pi^2(p+1)}$. If we choose $f(z)$ to be that function for which $Q(z) = 1 + \frac{zf''(z)}{f'(z)}$, then $f(z) \in UCV_p$ with $a_{p+1} = \frac{8p^2}{\pi^2(p+1)}$, which shows that this result is sharp. Now, when we put $|c_1| = B$, then

$$a_{p+2} = \frac{pa_p c_2 + (p+1)a_{p+1}c_1}{2(p+2)}$$


$$|a_{p+2}| \leq \frac{pB(1+Bp)}{2(p+2)}.$$

When $n = 3$

$$a_{p+3} = \frac{pa_p c_3 + (p+1)a_{p+1}c_2 + (p+2)a_{p+2}c_1}{3(p+3)}$$

$$|a_{p+3}| \leq \frac{1}{2} \frac{pB(1+Bp)(2+Bp)}{3(p+3)}$$

$$= \frac{1}{3(p+3)} pB(1+Bp) \left(1 + \frac{Bp}{2}\right).$$



Subordination Results for the Family of Uniformly Convex p -valent Functions

H.A. Al-Kharsani and S.S. Al-Hajiry

Title Page

Contents



Go Back

Close

Quit


Page 13 of 22

We now proceed by induction. Assume we have

$$|a_{p+n-1}| \leq \frac{1}{(n-1)(p+n-1)} pB(1+Bp) \left(1 + \frac{Bp}{2}\right) \cdots \left(1 + \frac{Bp}{n-2}\right) \\ = \frac{pB}{(n-1)(p+n-1)} \prod_{k=3}^n \left(1 + \frac{Bp}{k-2}\right).$$

Corollary 4.8. Let $f(z) = z^p + \sum_{k=2}^{\infty} a_{k+p-1} z^{k+p-1}$ and $f(z) \in UCV_p$. Then $|a_{p+n-1}| = O\left(\frac{1}{n^2}\right)$.

□



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 14 of 22

5. General Properties of Functions in UCV_p

Theorem 5.1. Let $f(z) = z^p + \sum_{k=2}^{\infty} a_{k+p-1} z^{k+p-1}$ and $f(z) \in UCV_p$. Then $f(z)$ is a p -valently convex function of order β in $|z| < r_1 = r_1(p, \beta)$, where $r_1(p, \beta)$ is the largest value of r for which

$$(5.1) \quad r^{k-1} \leq \frac{(p - \beta)(k - 1)}{(k + p - \beta - 1)B \prod_{j=3}^k \left(1 + \frac{pB}{j-2}\right)},$$

$$(k \in \mathbb{N} - \{1\}, 0 \leq \beta < p).$$

Proof. It is sufficient to show that for $f(z) \in UCV_p$,

$$\left|1 + \frac{zf''(z)}{f'(z)} - p\right| \leq p - \beta, \quad |z| < r_1(p, \beta), \quad 0 \leq \beta < p,$$

where $r_1(p, \beta)$ is the largest value of r for which the inequality (5.1) holds true. Observe that

$$\left|1 + \frac{zf''(z)}{f'(z)} - p\right| = \left|\frac{\sum_{k=2}^{\infty} (k + p - 1)(k - 1)a_{k+p-1}z^{k-1}}{p + \sum_{k=2}^{\infty} (k + p - 1)a_{k+p-1}z^{k-1}}\right|.$$

Then we have $\left|1 + \frac{zf''(z)}{f'(z)} - p\right| \leq p - \beta$ if and only if

$$\frac{\sum_{k=2}^{\infty} (k + p - 1)(k - 1) |a_{k+p-1}| r^{k-1}}{p - \sum_{k=2}^{\infty} (k + p - 1) |a_{k+p-1}| r^{k-1}} \leq p - \beta$$

$$\Rightarrow \sum_{k=2}^{\infty} (k + p - 1)(k + p - 1 - \beta) |a_{k+p-1}| r^{k-1} \leq p^2 - p\beta.$$



Subordination Results for the
Family of Uniformly Convex
 p -valent Functions

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 15 of 22

Then from Theorem 4.7 since $f(z) \in UCV_p$, we have

$$|a_{k+p-1}| \leq \frac{p\beta}{(k+p-1)(k-1)} \prod_{j=3}^k \left(1 + \frac{Bp}{j-2}\right)$$

and we may set

$$|a_{k+p-1}| = \frac{p\beta}{(k+p-1)(k-1)} \prod_{j=3}^k \left(1 + \frac{Bp}{j-2}\right) c_{k+p-1}, c_{k+p-1} \geq 0,$$

$$\left\{ k \in \mathbb{N} - \{1\}, \sum_{k=1}^{\infty} c_{k+p-1} \leq 1 \right\}.$$


Now, for each fixed r , we choose a positive integer $k_0 = k_0(r)$ for which

$$\frac{(k+p-1-\beta)}{(k-1)} r^{k-1}$$

is maximal. Then

$$\sum_{k=2}^{\infty} (k+p-1)(k+p-\beta-1) |a_{k+p-1}| r^{k-1}$$

$$\leq \frac{(k_0+p-\beta-1)}{(k_0-1)} r^{k_0-1} \prod_{j=3}^{k_0} \left(1 + \frac{Bp}{j-2}\right).$$



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 16 of 22

Consequently, the function $f(z)$ is a p -valently convex function of order β in $|z| < r_1 = r_1(p, \beta)$ provided that

$$\frac{(k_0 + p - \beta - 1)}{(k_0 - 1)} r^{k_0-1} \prod_{j=3}^k \left(1 + \frac{Bp}{j-2}\right) \leq p(p - \beta).$$

We find the value $r_0 = r_0(p, \beta)$ and the corresponding integer $k_0(r_0)$ so that

$$\frac{(k_0 + p - \beta - 1)}{(k_0 - 1)} r^{k_0-1} \prod_{j=3}^k \left(1 + \frac{Bp}{j-2}\right) = p(p - \beta), \quad (0 \leq \beta < p).$$

Then this value r_0 is the radius of p -valent convexity of order β for functions $f(z) \in UCV_p$. □

Theorem 5.2. $h(z) = z^p + b_{n+p-1}z^{n+p-1}$ is in UCV_p if and only if

$$r \leq \frac{p^2}{(p+n-1)(p+2n-2)},$$

where $|b_{n+p-1}| = r$ and $b_{n+p-1}z^{n+p-1} = re^{i\theta}$.

Proof. Let $w(z) = 1 + \frac{zh''(z)}{h'(z)}$. Then $h(z) \in UCV_p$ if and only if $w(z) \in PAR_p$ which means that $\operatorname{Re}\{w(z)\} \geq |w(z) - p|$. On the other side we have

$$\operatorname{Re} \left\{ 1 + \frac{zh''(z)}{h'(z)} \right\} \geq \left| 1 + \frac{zh''(z)}{h'(z)} - p \right|,$$



Subordination Results for the Family of Uniformly Convex p -valent Functions

H.A. Al-Kharsani and S.S. Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 17 of 22

then

$$\begin{aligned} \operatorname{Re} \left\{ 1 + \frac{zh''(z)}{h'(z)} \right\} &= \operatorname{Re} \left\{ (p-1) + \frac{p + (n+p-1)nr e^{i\theta}}{p + (n+p-1)r e^{i\theta}} \right\} \\ &= \frac{p^3 + p(n+p-1)(n+2p-1)r \cos \theta + (n+p-1)^3 r^2}{|p + (n+p-1)r e^{i\theta}|^2}. \end{aligned}$$

The right-hand side is seen to have a minimum for $\theta = \pi$ and this minimal value is

$$\frac{p^3 + p(n+p-1)(n+2p-1)r + (n+p-1)^3 r^2}{|p + (n+p-1)r e^{i\theta}|^2}.$$

Now, by computation we see that

$$\left| 1 + \frac{zh''(z)}{h'(z)} - p \right| = \frac{(n+p-1)(n-1)r}{|p + (n+p-1)r e^{i\theta}|}.$$

Then

$$(n+p-1)(n-1)r \leq \frac{p^3 + p(n+p-1)(n+2p-1)r + (n+p-1)^3 r^2}{p - (n+p-1)r},$$


which leads to

$$(n+p-1)(n-1)r \leq p^2 - (n+p-1)^2 r.$$

Hence,

$$r \leq \frac{p^2}{(n+p-1)(2n+p-2)}.$$

□



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 18 of 22

Theorem 5.3. Let $f(z) \in UCV$, then $(f(z))^p \in UCV_p$.

Proof. Let $w(z) = (f(z))^p$, then

$$1 + z \frac{w''(z)}{w'(z)} = 1 + z \frac{f''(z)}{f'(z)} + (p-1)z \frac{f'(z)}{f(z)}.$$

Then we find


$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{w''(z)}{w'(z)} \right\} - \left| z \frac{w''(z)}{w'(z)} - (p-1) \right| \\ = \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} + (p-1)z \frac{f'(z)}{f(z)} \right\} \\ - \left| z \frac{f''(z)}{f'(z)} + (p-1)z \frac{f'(z)}{f(z)} - (p-1) \right|. \end{aligned}$$

Since $f(z) \in UCV$, therefore we have

$$\begin{aligned} \operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} + (p-1)z \frac{f'(z)}{f(z)} \right\} \\ - \left| z \frac{f''(z)}{f'(z)} + (p-1)z \frac{f'(z)}{f(z)} - (p-1) \right| \\ \geq (p-1) \operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} - \left| z \frac{f'(z)}{f(z)} - 1 \right| \end{aligned}$$

$f(z) \in UCV$, then $f(z) \in SP$ [7] which means that

$$\operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} - \left| z \frac{f'(z)}{f(z)} - 1 \right| \geq 0.$$



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close


Quit

Page 19 of 22



Then

$$\operatorname{Re} \left\{ 1 + z \frac{w''(z)}{w'(z)} \right\} - \left| z \frac{w''(z)}{w'(z)} - (p-1) \right| \geq 0.$$

The following diagram shows the external function $k(z)$ of the class UCV when $(k(z))^p, p = 2$:



The following diagram shows the external function $K(z)$ of the class UCV_p when $p = 2$:



Subordination Results for the Family of Uniformly Convex p -valent Functions

H.A. Al-Kharsani and S.S. Al-Hajiry

Title Page

Contents




Go Back

Close

Quit

Page 20 of 22

And the following diagram shows that $(k(z))^p \prec K(z)$:



□


5.1. Remarks

Taking $p = 1$ in Theorem 3.1, we obtain the corresponding Theorem 1 of [7].

Taking $p = 1$ in Theorem 4.1, we obtain the corresponding Theorem 3 of [3].

Taking $p = 1$ in inequality (4.3), we obtain Theorem 6 of [7], and in inequalities (4.5), (4.6), we obtain Theorem 5 of [7].

Taking $p = 1$ in Theorem 5.2, we obtain Theorem 2 of [4].



**Subordination Results for the
Family of Uniformly Convex
 p -valent Functions**

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back


Close

Quit

Page 21 of 22

References

- [1] A.W. GOODMAN, On uniformly convex functions, *Ann. Polon. Math.*, **56**(1) (1991), 87–92.
- [2] S. KANAS AND A. WISNIOWSKA, Conic regions and k -uniform convexity, *J. Comput. Appl. Math.*, **105**(1-2) (1999), 327–336.
- [3] W.C. MA AND D. MINDA, Uniformly convex functions, *Ann. Polon. Math.*, **57**(2) (1992), 165–175.
- [4] S. OWA, On uniformly convex functions, *Math. Japonica*, **48**(3) (1998), 377–384.
- [5] M.S. ROBERTSON, On the theory of univalent functions, *Ann. Math.*, **2**(37) (1936), 347–408.
- [6] F. RONNING, A survey on uniformly convex and uniformly starlike functions, *Ann. Univ. Mariae Curie-Sklodowska Sect. A*, **47** (1993), 123–134.
- [7] F. RONNING, Uniformly convex functions and a corresponding class of starlike functions, *Proc. Amer. Math. Soc.*, **118**(1) (1993), 189–196.
- [8] F. RONNING, On uniform starlikeness and related properties of univalent functions, *Complex Variables Theory Appl.*, **24**(3-4) (1994), 233–239.



Subordination Results for the
Family of Uniformly Convex
 p -valent Functions

H.A. Al-Kharsani and S.S.
Al-Hajiry

Title Page

Contents



Go Back

Close

Quit

Page 22 of 22