


# Journal of Inequalities in Pure and Applied Mathematics



## SIMULTANEOUS APPROXIMATION FOR THE PHILLIPS-BÉZIER OPERATORS

VIJAY GUPTA, P.N. AGRAWAL AND HARUN KARSLI

School of Applied Sciences  
Netaji Subhas Institute of Technology  
Sector 3 Dwarka  
New Delhi 110075, India.  
*EMail:* vijay@nsit.ac.in

Department of Mathematics  
Indian Institute of Technology  
Roorkee 247667, India.  
*EMail:* pnappfma@iitr.ernet.in

Ankara University  
Faculty of Sciences Department of Mathematics  
06100 Tandoğan-Ankara-Turkey.  
*EMail:* karsli@science.ankara.edu.tr

volume 7, issue 4, article 141,  
2006.

*Received 28 July, 2006;  
accepted 17 November, 2006.*

*Communicated by: F. Qi*

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



## Abstract

We study the simultaneous approximation properties of the Bézier variant of the well known Phillips operators and estimate the rate of convergence of the Phillips-Bézier operators in simultaneous approximation, for functions of bounded variation.

*2000 Mathematics Subject Classification:* 41A25, 41A30.

*Key words:* Rate of convergence, Bounded variation, Total variation, Simultaneous approximation.

## Contents

1	Introduction .....	3
2	Auxiliary Results .....	5
3	Proof .....	8
References		

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 2 of 15](#)

# 1. Introduction

For  $\alpha \geq 1$ , the Phillips-Bézier operator is defined by

$$(1.1) \quad P_{n,\alpha}(f, x) = n \sum_{k=1}^{\infty} Q_{n,k}^{(\alpha)}(x) \int_0^{\infty} p_{n,k-1}(t) f(t) dt + Q_{n,0}^{(\alpha)}(x) f(0),$$

where  $n \in \mathbb{N}$ ,  $x \in [0, \infty)$ ,


$$Q_{n,k}^{(\alpha)}(x) = [J_{n,k}(x)]^{\alpha} - [J_{n,k+1}(x)]^{\alpha}, \quad J_{n,k}(x) = \sum_{j=k}^{\infty} p_{n,j}(x) \quad \text{and}$$
$$p_{n,k}(x) = e^{-nx} \frac{(nx)^k}{k!}.$$

For  $\alpha = 1$ , the operator (1.1) reduces to the Phillips operator [1]. Some approximation properties of the Phillips operators were recently studied by Finta and Gupta [2]. The rates of convergence in ordinary and simultaneous approximations on functions of bounded variation for the Phillips operators were estimated in [3], [4] and [5]. In the present paper we extend the earlier study and here we investigate and estimate the rate of convergence for the Bézier variant of the Phillips operators in simultaneous approximations by means of the decomposition technique of functions of bounded variation. We denote the class  $B_{r,\beta}$  by

$$B_{r,\beta} = \left\{ f : f^{(r-1)} \in C[0, \infty), f_{\pm}^{(r)}(x) \text{ exist everywhere} \right.$$

and are bounded on every finite subinterval of

$$\left. [0, \infty) \text{ and } f_{\pm}^{(r)}(x) = O(e^{\beta t}) (t \rightarrow \infty), \text{ for some } \beta > 0 \right\},$$



---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)

[◀◀](#)

[▶▶](#)

[◀](#)

[▶](#)

[Go Back](#)

[Close](#)

[Quit](#)

**Page 3 of 15**

$r = 1, 2, \dots$ . By  $f_{\pm}^{(0)}(x)$  we mean  $f(x\pm)$ . Our main theorem is stated as:

**Theorem 1.1.** Let  $f \in B_{r,\beta}$ ,  $r = 1, 2, \dots$  and  $\beta > 0$ . Then for every  $x \in (0, \infty)$  and  $n \geq \max \{r^2 + r, 4\beta\}$ , we have

$$\begin{aligned} \left| P_{n,\alpha}^{(r)}(f, x) - \frac{1}{\alpha+1} \left\{ f_+^{(r)}(x) + \alpha f_-^{(r)}(x) \right\} \right| &\leq \frac{r+\alpha-1}{\sqrt{2enx}} \left| f_+^{(r)}(x) - f_-^{(r)}(x) \right| \\ &+ \frac{1}{n} \left( 1 + \frac{2\alpha(1+2x)}{x^2} \right) \sum_{k=1}^n \bigvee_{x-x/\sqrt{k}}^{x+x/\sqrt{k}} (g_{r,x}) + \alpha \frac{\sqrt{2x+1}}{x\sqrt{n}} 2^{r/2} e^{2\beta x}, \end{aligned}$$

where  $g_{r,x}$  is the auxiliary function defined by

$$g_{r,x}(t) = \begin{cases} f^{(r)}(t) - f_+^{(r)}(x), & x < t < \infty \\ 0, & t = x \\ f^{(r)}(t) - f_-^{(r)}(x), & 0 \leq t < x \end{cases},$$

$\bigvee_a^b(g_{r,x}(t))$  is the total variation of  $g_{r,x}(t)$  on  $[a, b]$ . In particular  $g_{0,x}(t) \equiv g_x(t)$ , defined in [4].




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 15

## 2. Auxiliary Results

In this section we give certain lemmas, which are necessary for proving the main theorem.

**Lemma 2.1.** *For all  $x \in (0, \infty)$ ,  $\alpha \geq 1$  and  $k \in \mathbb{N} \cup \{0\}$ , we have*

$$p_{n,k}(x) \leq \frac{1}{\sqrt{2enx}}$$

and

$$Q_{n,k}^{(\alpha)}(x) \leq \frac{\alpha}{\sqrt{2enx}},$$

where the constant  $1/\sqrt{2e}$  and the estimation order  $n^{-1/2}$  (for  $n \rightarrow \infty$ ) are the best possible.


**Lemma 2.2 ([3]).** *If  $f \in L_1[0, \infty)$ ,  $f^{(r-1)} \in A.C._{loc}$ ,  $r \in \mathbb{N}$  and  $f^{(r)} \in L_1[0, \infty)$ , then*

$$P_n^{(r)}(f, x) = n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) f^{(r)}(t) dt.$$

**Lemma 2.3 ([3]).** *For  $m \in \mathbb{N} \cup \{0\}$ ,  $r \in \mathbb{N}$ , if we define the  $m$ -th order moment by*

$$\mu_{r,n,m}(x) = n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) (t-x)^m dt$$

then  $\mu_{r,n,0}(x) = 1$ ,  $\mu_{r,n,1}(x) = \frac{r}{n}$  and  $\mu_{r,n,2}(x) = \frac{2nx+r(r+1)}{n^2}$ .



---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 5 of 15**

Also there holds the following recurrence relation

$$n\mu_{r,n,m+1}(x) = x[\mu_{r,n,m}^{(1)}(x) + 2m\mu_{r,n,m-1}(x)] + (m+r)\mu_{r,n,m}(x).$$

Consequently by the recurrence relation, for all  $x \in [0, \infty)$ , we have

$$\mu_{r,n,m}(x) = O(n^{-(m+1)/2}).$$

**Remark 1.** In particular, by Lemma 2.3, for given any number  $n \geq r^2 + r$  and  $0 < x < \infty$ , we have

$$(2.1) \quad \mu_{r,n,2}(x) \leq \frac{2x+1}{n}.$$

**Remark 2.** We can observe from Lemma 2.2 and Lemma 2.3 that for  $r = 0$ , the summation over  $k$  starts from 1. For  $r = 0$ , Lemma 2.3 may be defined as [5, Lemma 2], with  $c = 0$ .

**Lemma 2.4.** Suppose  $x \in (0, \infty)$ ,  $r \in \mathbb{N} \cup \{0\}$ ,  $\alpha \geq 1$  and

$$K_{r,n,\alpha}(x, t) = n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) p_{n,k+r-1}(t).$$

Then for  $n \geq r^2 + r$ , there hold

$$(2.2) \quad \int_0^y K_{r,n,\alpha}(x, t) dt \leq \frac{\alpha(2x+1)}{n(x-y)^2}, \quad 0 \leq y < x,$$

$$(2.3) \quad \int_z^{\infty} K_{r,n,\alpha}(x, t) dt \leq \frac{\alpha(2x+1)}{n(z-x)^2}, \quad x < z < \infty.$$




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

[Go Back](#)

[Close](#)

[Quit](#)

**Page 6 of 15**

*Proof.* We first prove (2.2) as follows:

$$\begin{aligned}\int_0^y K_{r,n,\alpha}(x,t)dt &\leq \int_0^y \frac{(x-t)^2}{(x-y)^2} K_{r,n,\alpha}(x,t)dt \\ &\leq \frac{\alpha}{(x-y)^2} P_n((t-x)^2, x) \\ &\leq \frac{\alpha \mu_{r,n,2}(x)}{(x-y)^2} \leq \frac{\alpha(2x+1)}{n(x-y)^2},\end{aligned}$$

by using (2.1). The proof of (2.3) follows along similar lines. □



---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

**Page 7 of 15**


### 3. Proof

*Proof of Theorem 1.1.* Clearly

$$(3.1) \quad \left| P_{n,\alpha}^{(r)}(f, x) - \frac{1}{\alpha+1} \left\{ f_+^{(r)}(x) + \alpha f_-^{(r)}(x) \right\} \right| \\ \leq n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) \int_0^{\infty} p_{n+r-1,k}(x) |g_{r,x}(t)| dt \\ + \frac{1}{\alpha+1} \left| f_+^{(r)}(x) - f_-^{(r)}(x) \right| \left| P_{n,\alpha}^{(r)}(\operatorname{sgn}_{\alpha}(t-x), x) \right| .$$

We first estimate  $P_{n,\alpha}^{(r)}(\operatorname{sgn}_{\alpha}(t-x), x)$  as follows:

$$\begin{aligned} & P_{n,\alpha}^{(r)}(\operatorname{sgn}_{\alpha}(t-x), x) \\ &= n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) \left( \int_0^{\infty} \alpha p_{n,k+r-1}(t) dt - (1+\alpha) \int_0^x p_{n,k+r-1}(t) dt \right) \\ &= \alpha - (1+\alpha)n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) \int_0^{\infty} p_{n,k+r-1}(t) dt \\ &= \alpha - (1+\alpha)n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) \left( 1 - \sum_{j=0}^{k+r-1} p_{n,j}(x) \right) \\ &= (1+\alpha)n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) \sum_{j=0}^{k+r-1} p_{n,j}(x) - 1 \end{aligned}$$



---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 15

$$\begin{aligned}
&= (1 + \alpha)n \sum_{k=0}^{\infty} Q_{n,k}^{(\alpha)}(x) \left( \sum_{j=0}^k p_{n,j}(x) + \frac{r-1}{\sqrt{2enx}} \right) - 1 \\
&= (1 + \alpha) \left[ \sum_{j=0}^{\infty} p_{n,j}(x) \sum_{k=j}^{\infty} Q_{n,k}^{(\alpha)}(x) + \frac{r-1}{\sqrt{2enx}} \right] - 1 \\
&= (1 + \alpha) \left[ \sum_{j=0}^{\infty} p_{n,j}(x) [J_{n,j}(x)]^{\alpha} + \frac{r-1}{\sqrt{2enx}} \right] - \sum_{j=0}^{\infty} Q_{n,j}^{(\alpha)}(x).
\end{aligned}$$

By the mean value theorem, we find that

$$Q_{n,j}^{(\alpha+1)}(x) = [J_{n,j}(x)]^{\alpha+1} - [J_{n,j+1}(x)]^{\alpha+1} = (\alpha+1)p_{n,j}(x)[\gamma_{n,j}(x)]^{\alpha},$$

where

$$J_{n,j+1}(x) < \gamma_{n,j}(x) < J_{n,j}(x).$$

Hence by Lemma 2.1, we have

$$\begin{aligned}
&\left| P_{n,\alpha}^{(r)}(\operatorname{sgn}_{\alpha}(t-x), x) \right| \\
&\leq \left| (1 + \alpha) \left[ \sum_{j=0}^{\infty} p_{n,j}(x) ([J_{n,j}(x)]^{\alpha} - [\gamma_{n,j}(x)]^{\alpha}) \right] \right| + \frac{(1 + \alpha)(r-1)}{\sqrt{2enx}} \\
&\leq (1 + \alpha) \left[ \sum_{j=0}^{\infty} p_{n,j}(x) Q_{n,k}^{(\alpha)}(x) + \frac{r-1}{\sqrt{2enx}} \right] \\
(3.2) \quad &\leq (1 + \alpha) \left[ \frac{\alpha}{\sqrt{2enx}} + \frac{r-1}{\sqrt{2enx}} \right] = \frac{(1 + \alpha)(r + \alpha - 1)}{\sqrt{2enx}}.
\end{aligned}$$




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 9 of 15](#)

Next we estimate  $P_n^{(r)}(g_{r,x}, x)$ . By the Lebesgue-Stieltjes integral representation, we have

$$\begin{aligned}
 P_{n,\alpha}^{(r)}(g_{r,x}, x) &= \int_0^\infty g_{r,x}(t) K_{r,n,\alpha}(x, t) dt \\
 &= \left( \int_{I_1} + \int_{I_2} + \int_{I_3} + \int_{I_4} \right) g_{r,x}(t) K_{r,n,\alpha}(x, t) dt \\
 (3.3) \quad &= R_1 + R_2 + R_3 + R_4,
 \end{aligned}$$

say, where  $I_1 = [0, x - x/\sqrt{n}]$ ,  $I_2 = [x - x/\sqrt{n}, x + x/\sqrt{n}]$ ,  $I_3 = [x + x/\sqrt{n}, 2x]$  and  $I_4 = [2x, \infty)$ . Let us define

$$\eta_{r,n,\alpha}(x, t) = \int_0^t K_{r,n,\alpha}(x, u) du.$$

We first estimate  $R_1$ . Writing  $y = x - x/\sqrt{n}$  and using integration by parts, we have

$$\begin{aligned}
 R_1 &= \int_0^y g_{r,x}(t) d_t(\eta_{r,n,\alpha}(x, t)) \\
 &= g_{r,x}(y) \eta_{r,n,\alpha}(x, y) - \int_0^y \eta_{r,n,\alpha}(x, t) d_t(g_{r,x}(t)).
 \end{aligned}$$

By Remark 1, it follows that

$$|R_1| \leq \sqrt[y]{(g_{r,x})} \eta_{r,n,\alpha}(x, y) + \int_0^y \eta_{r,n,\alpha}(x, t) d_t \left( -\sqrt[t]{(g_{r,x})} \right)$$




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 10 of 15](#)

$$\leq \sqrt[n]{\int_y^x (g_{r,x})} \frac{\alpha(2x+1)}{n(x-y)^2} + \frac{\lambda x}{n} \int_0^y \frac{1}{(x-t)^2} dt \left( - \sqrt[n]{\int_t^x (g_{r,x})} \right).$$

Integrating by parts the last term, we have after simple computation,

$$|R_1| \leq \frac{\alpha(2x+1)}{n} \left[ \frac{\sqrt[n]{0}(g_{r,x})}{x^2} + 2 \int_0^y \frac{\sqrt[n]{t}(g_{r,x})}{(x-t)^3} dt \right].$$

Now replacing the variable  $y$  in the last integral by  $x - x/\sqrt{t}$ , we get


$$(3.4) \quad |R_1| \leq \frac{2\alpha(2x+1)}{nx^2} \sum_{k=1}^n \sqrt[n]{x-x/\sqrt{k}} (g_{r,x}).$$

Next we estimate  $R_2$ . For  $t \in [x - x/\sqrt{n}, x + x/\sqrt{n}]$ , we have

$$|g_{r,x}(t)| = |g_{r,x}(t) - g_{r,x}(x)| \leq \sqrt[n]{x+x/\sqrt{n}} (g_{r,x}).$$

Also by the fact that  $\int_a^b d_t(\eta_{r,n}(x, t)) \leq 1$  for  $(a, b) \subset [0, \infty)$ , therefore

$$(3.5) \quad |R_2| \leq \sqrt[n]{x-x/\sqrt{n}} (g_{r,x}) \leq \frac{1}{n} \sum_{k=1}^n \sqrt[n]{x+x/\sqrt{k}} (g_{r,x}).$$




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 11 of 15](#)

To estimate  $R_3$ , we take  $z = x + x/\sqrt{n}$ , thus


$$\begin{aligned}
 R_3 &= \int_z^{2x} K_{r,n,\alpha}(x, t) g_{r,x}(t) dt \\
 &= - \int_z^{2x} g_{r,x}(t) d_t(1 - \eta_{r,n,\alpha}(x, t)) \\
 &= -g_{r,x}(2x)(1 - \eta_{r,n,\alpha}(x, 2x)) + g_{r,x}(z)(1 - \eta_{r,n,\alpha}(x, z)) \\
 &\quad + \int_z^{2x} (1 - \eta_{r,n,\alpha}(x, t)) d_t(g_{r,x}(t)).
 \end{aligned}$$

Thus arguing similarly as in the estimate of  $R_1$ , we obtain

$$(3.6) \quad |R_3| \leq \frac{2\alpha(2x+1)}{nx^2} \sum_{k=1}^n \bigvee_{x}^{x+x/\sqrt{k}} (g_{r,x}).$$

Finally we estimate  $R_4$  as follows

$$\begin{aligned}
 |R_4| &= \left| \int_{2x}^{\infty} K_{r,n,\alpha}(x, t) g_{r,x}(t) dt \right| \\
 &\leq n\alpha \sum_{k=0}^{\infty} p_{n,k}(x) \int_{2x}^{\infty} p_{n,k+r-1}(t) e^{\beta t} dt \\
 &\leq \frac{n\alpha}{x} \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) e^{\beta t} |t-x| dt
 \end{aligned}$$




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 12 of 15](#)


$$\leq \frac{\alpha}{x} \left( n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) (t-x)^2 dt \right)^{\frac{1}{2}} \\ \times \left( n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) e^{2\beta t} dt \right)^{\frac{1}{2}}.$$

For the first expression above we use Remark 1. To evaluate the second expression, we proceed as follows:

$$n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) e^{2\beta t} dt \\ = n \sum_{k=0}^{\infty} p_{n,k}(x) \frac{n^{k+r-1}}{(k+r-1)!} \int_0^{\infty} t^{k+r-1} e^{-(n-2\beta)t} dt,$$

$$n \sum_{k=0}^{\infty} p_{n,k}(x) \frac{n^{k+r-1}}{(k+r-1)!} \frac{\Gamma(k+r)}{(n-2\beta)^{k+r}} \\ = \frac{n^r}{(n-2\beta)^r} \sum_{k=0}^{\infty} \left( \frac{n}{n-2\beta} \right)^k p_{n,k}(x),$$

$$\frac{n^r}{(n-2\beta)^r} e^{-nx} \sum_{k=0}^{\infty} \left( \frac{n^2 x}{n-2\beta} \right)^k \frac{1}{k!} = \frac{n^r}{(n-2\beta)^r} e^{2nx\beta/(n-2\beta)} \leq 2^r e^{4\beta x}$$




---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 13 of 15](#)

for  $n > 4\beta$ . Thus

$$\begin{aligned} |R_4| &\leq \frac{\alpha}{x} \left( n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) (t-x)^2 dt \right)^{\frac{1}{2}} \\ &\quad \times \left( n \sum_{k=0}^{\infty} p_{n,k}(x) \int_0^{\infty} p_{n,k+r-1}(t) e^{2\beta t} dt \right)^{\frac{1}{2}} \\ (3.7) \quad &\leq \frac{\alpha \sqrt{2x+1}}{x \sqrt{n}} 2^{r/2} e^{2\beta x}. \end{aligned}$$

Combining the estimates of (3.1)-(3.7), we get the required result. □



---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 14 of 15](#)

## References

- [1] R.S. PHILLIPS, An inversion formula for semi-groups of linear operators, *Ann. Math. (Ser. 2)*, **59** (1954), 352–356.
- [2] Z. FINTA AND V. GUPTA, Direct and inverse estimates for Phillips type operators, *J. Math Anal Appl.*, **303** (2005), 627–642.
- [3] N.K. GOVIL, V. GUPTA AND M.A. NOOR, Simultaneous approximation for the Phillips operators, *International Journal of Math and Math Sci.*, Vol. **2006** Art Id. 49094 92006, 1-9.
- [4] V. GUPTA AND G.S. SRIVASTAVA, On the rate of convergence of Phillips operators for functions of bounded variation, *Ann. Soc. Math. Polon. Ser. I Commentat. Math.*, **36** (1996), 123–130.
- [5] H.M. SRIVASTAVA AND V. GUPTA, A certain family of summation-integral type operators, *Math. Comput. Modelling*, **37** (2003), 1307–1315.
- [6] X.M. ZENG AND J.N. ZHAO, Exact bounds for some basis functions of approximation operators, *J. Inequal. Appl.*, **6** (2001), 563–575.



---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

Vijay Gupta, P.N. Agrawal and  
Harun Karsli

---

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 15 of 15](#)