

MONOTONICITY AND CONCAVITY PROPERTIES OF SOME FUNCTIONS INVOLVING THE GAMMA FUNCTION WITH APPLICATIONS

SENLIN GUO

Department of Mathematics
University of Manitoba
Winnipeg, MB, R3T 2N2
Canada.

EMail: umguos@cc.umanitoba.ca



volume 7, issue 2, article 45,
2006.

*Received 28 October, 2005;
accepted 27 November, 2005.*

Communicated by: F. Qi

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


In this article, we give the monotonicity and concavity properties of some functions involving the gamma function and some equivalence sequences to the sequence $n!$ with exact equivalence constants.

2000 Mathematics Subject Classification: Primary 33B15; Secondary 26D07, 26D20.

Key words: Gamma function, Monotonicity, Concavity, Equivalence

Contents

1	Introduction and Main Results	3
2	Lemmas	8
3	Proof of Main Results	9
	References	



Monotonicity and Concavity Properties of Some Functions Involving the Gamma Function with Applications

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 2 of 15

1. Introduction and Main Results

Throughout the paper, let \mathbb{N} denote the set of all positive integers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

We say $a_n \simeq b_n$ ($n \geq n_0$) if there exist two constants $c_1 > 0$ and $c_2 > 0$ such that

$$(1.1) \quad c_1 b_n \leq a_n \leq c_2 b_n$$

hold for all $n \geq n_0$. The fixed numbers c_1 and c_2 in (1.1) are called equivalence constants.

The incomplete gamma function is defined for $\operatorname{Re} z > 0$ by


$$(1.2) \quad \Gamma(z, x) = \int_x^\infty t^{z-1} e^{-t} dt, \quad \gamma(z, x) = \int_0^x t^{z-1} e^{-t} dt,$$

and $\Gamma(z, 0) = \Gamma(z)$ is called the gamma function. The logarithmic derivative of $\Gamma(z)$, denoted by $\psi(z) = \Gamma'(z)/\Gamma(z)$, is called the psi or digamma function, and $\psi^{(k)}$ for $k \in \mathbb{N}$ are called the polygamma functions. One of the elementary properties of the gamma function is $\Gamma(x+1) = x\Gamma(x)$. In particular, $\Gamma(n+1) = n!$.

In [13], it was proved by F. Qi that the functions

$$(1.3) \quad f(s, r) = \left[\frac{\Gamma(s)}{\Gamma(r)} \right]^{\frac{1}{s-r}},$$

$$(1.4) \quad f(s, r, x) = \left[\frac{\Gamma(s, x)}{\Gamma(r, x)} \right]^{\frac{1}{s-r}}$$



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 3 of 15

and

$$(1.5) \quad g(s, r, x) = \left[\frac{\gamma(s, x)}{\gamma(r, x)} \right]^{\frac{1}{s-r}}$$

are increasing with respect to $r > 0$, $s > 0$, or $x > 0$.

E. A. Karatsuba [9] proved that the function

$$(1.6) \quad f_1(x) = [g(x)]^6 - (8x^3 + 4x^2 + x),$$

where

$$(1.7) \quad g(x) = \left(\frac{e}{x}\right)^x \frac{\Gamma(1+x)}{\sqrt{\pi}},$$

is strictly increasing from $[1, \infty)$ onto $[f_1(1), f_1(\infty))$ with

$$f_1(1) = \frac{e^6}{\pi^3} - 13 \quad \text{and} \quad f_1(\infty) = \frac{1}{30}.$$

In 2003, in [1], H. Alzer proved that


$$\alpha \leq f_1(x) < \frac{1}{30}, \quad x \in (0, \infty),$$

where

$$\alpha = \min_{x>0} f_1(x) = 0.0100450 \dots = f_1(x_0)$$

for some $x_0 \in [0.6, 0.7]$. Since $f_1(x_0) < f_1(1)$ and

$$f_1(x_0) < \lim_{x \rightarrow 0^+} f_1(x) = \frac{1}{\sqrt{\pi}},$$



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 4 of 15

his result shows that $f_1(x)$ is not still monotonic on $(0, 1]$.

In [3], it was shown in 1997, by G. Anderson and S. Qiu, that the function

$$(1.8) \quad f_2(x) = \frac{\ln \Gamma(x+1)}{x \ln x}$$

is strictly increasing from $(1, \infty)$ onto $(1-\gamma, 1)$, where γ is the Euler-Mascheroni constant. H. Alzer, in 1998 in [2], proved that $f_2(x)$, with

$$(1.9) \quad f_2(1) = \lim_{x \rightarrow 1} f_2(x) = 1 - \gamma,$$

is strictly increasing on $(0, \infty)$. Also note that the function $f_2(x)$ was proved to be concave on $(1, \infty)$ in [6] in 2000 by A. Elbert and A. Laforgia.

In [5, 8, 10, 12, 14, 17], monotonicity properties of other functions related to the (di)gamma function were obtained.

In this article, we shall give some monotonicity and concavity properties of several functions involving the gamma function and, as applications, deduce some equivalence sequences to the sequence $n!$ with best equivalence constants.


Our main results are as follows.

Theorem 1.1. *The functions*

$$(1.10) \quad f(x) = \frac{x^{x+\frac{1}{2}}}{e^x \Gamma(x+1)}$$

and

$$(1.11) \quad F(x) = \frac{e^x \Gamma(x+1)}{x^x}$$



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 5 of 15

are strictly logarithmically concave and strictly increasing from $(0, \infty)$, respectively, onto $(0, 1/\sqrt{2\pi})$ and onto $(1, \infty)$.

Theorem 1.2. *The function*

$$(1.12) \quad g(x) = \frac{e^x \Gamma(x+1)}{\left(x + \frac{1}{2}\right)^{x+\frac{1}{2}}}$$

is strictly logarithmically concave and strictly increasing from $(-\frac{1}{2}, \infty)$ onto $(\sqrt{\pi/e}, \sqrt{2\pi/e})$.

Theorem 1.3. *The function*

$$(1.13) \quad h(x) = \frac{e^x \Gamma(x+1) \sqrt{x-1}}{x^{x+1}}$$

is strictly logarithmically concave and strictly increasing from $(1, \infty)$ onto $(0, \sqrt{2\pi})$.

As applications of these theorems, we have the following corollaries.


Corollary 1.4. *For $n \in \mathbb{N}$,*

$$(1.14) \quad n! \simeq e^{-n} n^{n+1/2}.$$

Moreover, for all $n \in \mathbb{N}$,

$$(1.15) \quad \sqrt{2\pi} \cdot e^{-n} n^{n+1/2} < n! \leq e \cdot e^{-n} n^{n+1/2}.$$

The equivalence constants $\sqrt{2\pi}$ and e in (1.15) are best possible.



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 6 of 15

Corollary 1.5. For $n \in \mathbb{N}_0$,

$$(1.16) \quad n! \simeq e^{-n} \left(n + \frac{1}{2} \right)^{n+\frac{1}{2}}.$$

Moreover, for all $n \in \mathbb{N}_0$,

$$(1.17) \quad \sqrt{2} e^{-n} \left(n + \frac{1}{2} \right)^{n+\frac{1}{2}} \leq n! < \sqrt{\frac{2\pi}{e}} e^{-n} \left(n + \frac{1}{2} \right)^{n+\frac{1}{2}}.$$

The equivalence constants $\sqrt{2}$ and $\sqrt{2\pi/e}$ in (1.17) are best possible.

Corollary 1.6. For $n \geq 2$,

$$(1.18) \quad n! \simeq \sqrt{\frac{n}{n-1}} e^{-n} n^{n+1/2}.$$

Furthermore, for all $n \geq 2$,


$$(1.19) \quad \left(\frac{e}{2} \right)^2 \sqrt{\frac{n}{n-1}} e^{-n} n^{n+1/2} \leq n! < \sqrt{2\pi} \sqrt{\frac{n}{n-1}} e^{-n} n^{n+1/2}.$$

The equivalence constants $(e/2)^2$ and $\sqrt{2\pi}$ in (1.19) are best possible.

Remark 1. In [16, Theorem 5], it was proved that for $n \geq 2$,

$$(1.20) \quad \sqrt{2\pi} e^{-n} n^{n+1/2} < n! < \left(\frac{n}{n-1} \right)^{\frac{1}{2}} \sqrt{2\pi} e^{-n} n^{n+1/2},$$

which can be directly deduced from (1.15) and (1.19).



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 7 of 15

2. Lemmas

We need the following lemmas to prove our results.

Lemma 2.1 ([4, p. 20]). As $x \rightarrow \infty$,

$$(2.1) \quad \ln \Gamma(x) = \left(x - \frac{1}{2}\right) \ln x - x + \ln \sqrt{2\pi} + O\left(\frac{1}{x}\right).$$

Lemma 2.2 ([7, p. 892] and [11, p. 17]). For $x > 0$,

$$(2.2) \quad \psi(x) = \ln x - \frac{1}{2x} - 2 \int_0^\infty \frac{t \, dt}{(t^2 + x^2)(e^{2\pi t} - 1)},$$


$$(2.3) \quad \psi\left(x + \frac{1}{2}\right) = \ln x + 2 \int_0^\infty \frac{t \, dt}{(t^2 + 4x^2)(e^{\pi t} + 1)}.$$

Lemma 2.3. The function

$$(2.4) \quad \varphi(x) = \ln \frac{x+1}{x+\frac{1}{2}} - \frac{1}{2x}$$

is strictly increasing from $(0, \infty)$ onto $(-\infty, 0)$.

Proof. We omit the proof of this lemma due to its simplicity. □



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 8 of 15

3. Proof of Main Results

Proof of Theorem 1.1. Taking the logarithm of $f(x)$ defined by (1.10) and differentiating directly yields

$$(3.1) \quad \ln f(x) = \left(x - \frac{1}{2}\right) \ln x - x - \ln \Gamma(x),$$

$$(3.2) \quad [\ln f(x)]' = \ln x - \frac{1}{2x} - \psi(x).$$

Then by formula (2.2) of Lemma 2.2,

$$(3.3) \quad [\ln f(x)]' = 2 \int_0^\infty \frac{t dt}{(t^2 + x^2)(e^{2\pi t} - 1)}, \quad x > 0.$$

Hence, $[\ln f(x)]' > 0$ for $x \in (0, \infty)$, which means that $\ln f(x)$, and then $f(x)$, is strictly increasing on $(0, \infty)$.

It is easy to see that $\lim_{x \rightarrow 0^+} f(x) = 0$. By (3.1) and Lemma 2.1, we have


$$(3.4) \quad \ln f(x) = -\ln \sqrt{2\pi} + O\left(\frac{1}{x}\right) \rightarrow \ln \frac{1}{\sqrt{2\pi}}, \quad x \rightarrow \infty,$$

which implies $\lim_{x \rightarrow \infty} f(x) = 1/\sqrt{2\pi}$.

Taking the logarithm of $F(x)$ defined by (1.11) and differentiating easily gives

$$(3.5) \quad \ln F(x) = x + \ln \Gamma(x+1) - x \ln x,$$

$$(3.6) \quad [\ln F(x)]' = \psi(x+1) - \ln x.$$



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents




Go Back

Close

Quit

Page 9 of 15



Then by (2.3) of Lemma 2.2, for all $x > 0$,

$$(3.7) \quad [\ln F(x)]' = \ln \left(1 + \frac{1}{2x} \right) + 2 \int_0^\infty \frac{t \, dt}{\left[t^2 + 4 \left(x + \frac{1}{2} \right)^2 \right] (e^{\pi t} + 1)} > 0.$$

Hence, $\ln F(x)$, and then $F(x)$, is strictly increasing on $(0, \infty)$.

It is easy to see that $\lim_{x \rightarrow 0^+} F(x) = 1$. By using Lemma 2.1, from (3.5),

$$(3.8) \quad \ln F(x) = \frac{1}{2} \ln x + \ln \sqrt{2\pi} + O\left(\frac{1}{x}\right), \quad x \rightarrow \infty.$$

Therefore, $\ln F(x)$, and then $F(x)$ tends to ∞ as $x \rightarrow \infty$.

Formulas (3.3) and (3.7) tell us that $[\ln f(x)]'$ and $[\ln F(x)]'$ are both strictly decreasing. Therefore, $\ln f(x)$ and $\ln F(x)$ are strictly concave, that is, the function $f(x)$ and $F(x)$ are both logarithmically concave. \square

Proof of Theorem 1.2. Taking the logarithm of $g(x)$ defined by (1.12) and differentiating shows

$$(3.9) \quad \ln g(x) = x + \ln \Gamma(x+1) - \left(x + \frac{1}{2}\right) \ln \left(x + \frac{1}{2}\right),$$

$$(3.10) \quad [\ln g(x)]' = \psi(x+1) - \ln \left(x + \frac{1}{2}\right).$$

Then, by formula (2.3) of Lemma 2.2, we have

$$(3.11) \quad [\ln g(x)]' = 2 \int_0^\infty \frac{t \, dt}{[t^2 + (2x+1)^2](e^{\pi t} + 1)}, \quad x > -\frac{1}{2}.$$

**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 10 of 15

So

$$(3.12) \quad [\ln g(x)]' > 0, \quad x \in \left(-\frac{1}{2}, \infty\right),$$

which means that $\ln g(x)$, then $g(x)$, is strictly increasing on $(-\frac{1}{2}, \infty)$.

Since $\Gamma(1/2) = \sqrt{\pi}$, it is easy to verify that $\lim_{x \rightarrow -1/2} g(x) = \sqrt{\pi/e}$.

From (3.9) and Lemma 2.1, it is obtained that

$$(3.13) \quad \ln g(x) = \left(x + \frac{1}{2}\right) \ln \frac{x + 1}{x + \frac{1}{2}} + \ln \sqrt{2\pi} - 1 + O\left(\frac{1}{x}\right), \quad x \rightarrow \infty.$$

Hence $\ln g(x) \rightarrow \ln \sqrt{2\pi/e}$ as $x \rightarrow \infty$, and then $\lim_{x \rightarrow \infty} g(x) = \sqrt{2\pi/e}$.

Formula (3.11) shows that $[\ln g(x)]'$ is strictly decreasing. Therefore, $\ln g(x)$ is strictly concave, that is, the function $g(x)$ is logarithmically concave. \square

Proof of Theorem 1.3. Taking the logarithm of $h(x)$ defined by (1.13) and differentiating straightforwardly reveals

$$(3.14) \quad \ln h(x) = \ln \Gamma(x) + x + \frac{1}{2} \ln(x-1) - x \ln x,$$

$$(3.15) \quad [\ln h(x)]' = \psi(x) + \frac{1}{2(x-1)} - \ln x.$$

By setting $x = u + 1$ with $u > 0$, we have

$$(3.16) \quad [\ln h(x)]' = \psi(u+1) + \frac{1}{2u} - \ln(u+1) = [\ln g(u)]' - \varphi(u),$$



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 11 of 15

where $g(u)$ and $\varphi(u)$ are respectively defined by (1.12) and (2.4). From (3.12) and Lemma 2.3, it is deduced that $[\ln h(x)]' > 0$ for $x > 1$. Therefore, $\ln h(x)$, and then $h(x)$, is strictly increasing on $(1, \infty)$.

It is obvious that $\lim_{x \rightarrow 1^+} h(x) = 0$. From (3.14) and Lemma 2.1, we see

$$(3.17) \quad \ln h(x) = \frac{1}{2} \ln \frac{x-1}{x} + \ln \sqrt{2\pi} + O\left(\frac{1}{x}\right) \rightarrow \ln \sqrt{2\pi}, \quad x \rightarrow \infty.$$

So $\lim_{x \rightarrow \infty} h(x) = \sqrt{2\pi}$.

Considering the logarithmic concavity of $g(x)$ and the increasing monotonicity of $\varphi(x)$ in (3.16) reveals that $[\ln h(x)]'$ is strictly decreasing. Therefore, $\ln h(x)$ is strictly concave, that is, the function $h(x)$ is logarithmically concave. \square


Proof of Corollary 1.4. By Theorem 1.1, we know that the function $f(x)$ is strictly increasing from $(0, \infty)$ onto $(0, \frac{1}{\sqrt{2\pi}})$, hence

$$(3.18) \quad \frac{1}{e} = f(1) \leq f(n) = \frac{n^{n+1/2}}{e^n n!} < \frac{1}{\sqrt{2\pi}}$$

for $n \in \mathbb{N}$, and

$$(3.19) \quad \lim_{n \rightarrow \infty} \frac{n^{n+1/2}}{e^n n!} = \frac{1}{\sqrt{2\pi}}.$$

From (3.18) and (3.19), we see that Corollary 1.4 is true. \square



Monotonicity and Concavity Properties of Some Functions Involving the Gamma Function with Applications

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 12 of 15

Proof of Corollary 1.5. By Theorem 1.2, we see that the function $g(x)$ is strictly increasing from $(-\frac{1}{2}, \infty)$ onto $(\sqrt{\pi/e}, \sqrt{2\pi/e})$. So

$$(3.20) \quad \sqrt{2} = g(0) \leq g(n) = \frac{e^n n!}{(n + \frac{1}{2})^{n+1/2}} < \sqrt{\frac{2\pi}{e}}, \quad n \in \mathbb{N}_0$$

and

$$(3.21) \quad \lim_{n \rightarrow \infty} \frac{e^n n!}{(n + \frac{1}{2})^{n+1/2}} = \sqrt{\frac{2\pi}{e}}.$$

Inequality (3.20) is equivalent to (1.17). Since the constants $\sqrt{2}$ and $\sqrt{2\pi/e}$ are best possible in (3.20), they are also best possible in (1.17). \square


Proof of Corollary 1.6. The monotonicity of $h(x)$ by Theorem 1.3 implies

$$(3.22) \quad \left(\frac{e}{2}\right)^2 = h(2) \leq h(n) = \frac{e^n n! \sqrt{n-1}}{n^{n+1}} < \sqrt{2\pi}, \quad n \geq 2$$

and

$$(3.23) \quad \lim_{n \rightarrow \infty} \frac{e^n n! \sqrt{n-1}}{n^{n+1}} = \sqrt{2\pi}.$$

From (3.22) and (3.23), we see that Corollary 1.6 is valid. \square



Monotonicity and Concavity Properties of Some Functions Involving the Gamma Function with Applications

Senlin Guo

Title Page

Contents



Go Back


Close

Quit

Page 13 of 15

References

- [1] H. ALZER, On Ramanujan's double inequality for the gamma function, *Bull. London Math. Soc.*, **35** (2003), 601–607.
- [2] H. ALZER, Inequalities for gamma and polygamma functions, *Abh. Math. Sem. Hamburg*, **68** (1998), 363–372.
- [3] G.D. ANDERSON AND S.-L. QIU, A monotonicity property of the gamma function, *Proc. Amer. Math. Soc.*, **125** (1997), 3355–3362.
- [4] G.E. ANDREWS, R. ASKEY AND R. ROY, *Special Functions*, Cambridge University Press, Cambridge, 1999.
- [5] Ch.-P. CHEN AND F. QI, Monotonicity results for the gamma function, *J. Inequal. Pure Appl. Math.*, **4**(2) (2003), Art. 44. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=282>].
- [6] A. ELBERT AND A. LAFORGIA, On some properties of the gamma function, *Proc. Amer. Math. Soc.*, **128** (2000), 2667–2673.
- [7] I.S. GRADSHTEYN AND I.M. RYZHIK, *Table of Integrals, Series, and Products*, Sixth Edition, Academic Press, New York, 2000.
- [8] B.-N. GUO AND F. QI, Inequalities and monotonicity for the ratio of gamma functions, *Taiwanese J. Math.*, **7** (2003), 239–247.
- [9] E.A. KARATSUBA, On asymptotic representation of the Euler gamma function by Ramanujan, *J. Comp. Appl. Math.*, **135** (2001), 225–240.



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents




Go Back

Close

Quit

Page 14 of 15

- [10] D. KERSHAW AND A. LAFORGIA, Monotonicity results for the gamma function, *Atti. Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.*, **119** (1985), 127–133.
- [11] W. MAGNUS, F. OBERHETTINGER AND R.P. SONI, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer, Berlin-New York, 1966.
- [12] M.E. MULDOON, Some monotonicity properties and characterization of the gamma function, *Aequat. Math.*, **18** (1978), 54–63.
- [13] F. QI, Monotonicity results and inequalities for the gamma and incomplete gamma functions, *Math. Inequal. Appl.*, **5**(1) (2002), 61–67.
- [14] S.L. QIU AND M. VUORINEN, Some properties of the gamma and psi functions, with applications, *Math. Comp.*, **74** (2005), 723–742.
- [15] A.W. ROBERTS AND D.E. VARBERG, *Convex Functions*, Academic Press, New York, 1973.
- [16] J. SÁNDOR AND L. DEBNATH, On certain inequalities involving the constant e and their applications, *J Math. Anal. Appl.*, **249** (2000) 569–592.
- [17] H. VOGT AND J. VOIGT, A monotonicity property of the Γ -function, *J. Inequal. Pure Appl. Math.*, **3**(5) (2002), Art. 73. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=225>].



**Monotonicity and Concavity
Properties of Some Functions
Involving the Gamma Function
with Applications**

Senlin Guo

Title Page

Contents



Go Back

Close

Quit

Page 15 of 15