

UNIMODAL SINGULARITIES AND DIFFERENTIAL OPERATORS

by

Yayoi Nakamura & Shinichi Tajima

Abstract. — An algebraic local cohomology class attached to a hypersurface isolated singularity is considered from the view point of algebraic analysis. A holonomic system derived from first order differential equations associated to a cohomology class and its solutions are studied. For the unimodal singularities case, it is shown that the multiplicity of the holonomic system associated to the cohomology class, which generates the dual space of Milnor algebra, is equal to two.

Résumé (Singularités unimodulaires et opérateurs différentiels). — On considère une classe de cohomologie locale algébrique attachée à une hypersurface à singularités isolées, du point de vue de l'analyse algébrique. On étudie le système holonome des équations aux dérivées partielles du premier ordre associé à la classe de cohomologie ainsi que ses solutions. On décrit une méthode générale pour examiner le système holonome associé. Il est montré que, dans le cas de singularités isolées unimodales, la multiplicité du système holonome associé à la classe génératrice de l'espace dual de l'algèbre de Milnor est égale à deux. Une description explicite des solutions du système holonome est donnée.

1. Introduction

We consider algebraic local cohomology classes attached to hypersurface isolated singularities by using first order differential operators. The purpose is to clarify the difference between quasihomogeneity and non-quasihomogeneity of the singularity from a view point of \mathcal{D} -modules theory.

In [3], we gave a characterization of quasihomogeneity of hypersurface isolated singularities based on \mathcal{D} -modules theory. We considered an algebraic local cohomology class attached to a given singularity which generates the dual space of Milnor algebra, and an associated holonomic system derived from first order annihilators of the cohomology class in consideration. We showed that the simplicity of the associated

2000 Mathematics Subject Classification. — Primary 32S25; Secondary 32C36, 32C38.

Key words and phrases. — Isolated hypersurface singularities, algebraic local cohomology classes, holonomic \mathcal{D} -modules.

first order holonomic system is equivalent to the quasihomogeneity of the singularity. For non-quasihomogeneous case, the structure of associated first order holonomic system is not fully investigated. By putting the idea in [3] to practical use to non-quasihomogeneous singularities, we consider relations between non-quasihomogeneous singularities and a structure of first order holonomic systems. In this paper, we give a practical method of computations. Applying the method, we give an explicit description of algebraic local cohomology solution space of the holonomic system in question and also present a detailed result of computations on normal forms of exceptional unimodal singularities.

In §2, we briefly recall the dual space of Milnor algebra with respect to the Grothendieck local duality and introduce a holonomic system derived from differential operators of order at most one which annihilate a generator of the dual space. In §3, we give a method for describing the solution space of the first order holonomic system. We recall our results on the quasihomogeneous singularities and the unimodal singularities concerning to the solution space of the holonomic system. We show, for the unimodal singularities case, that the solution space of the holonomic system derived from first order differential equations is of dimension two. In §4, we give a method for examining semiquasihomogeneous singularities from the computational point of view. We show that the computation of the solution space of the first order holonomic system can be carried out in finite dimensional vector spaces. In §5, we give results of the computations for each normal form of exceptional unimodal singularities. For proves of results stated in §2 and §3, please refer to [3].

2. The dual space of Milnor algebra and first order differential operators

Let X be an open neighborhood of the origin O in the n dimensional affine space \mathbb{C}^n and \mathcal{O}_X the sheaf of holomorphic functions on X . Let $f(z) \in \mathcal{O}_{X,O}$ be a holomorphic function on X with an isolated singularity at the origin O . Denote by \mathcal{I} the ideal in $\mathcal{O}_{X,O}$ generated by the partial derivatives $f_j = \partial f(z)/\partial z_j$ ($j = 1, \dots, n$) of $f(z)$:

$$\mathcal{I} = \langle f_1, \dots, f_n \rangle_O.$$

From the Grothendieck local duality, we have a non-degenerate perfect pairing

$$(1) \quad \Omega_{X,O}^n / \mathcal{I} \Omega_{X,O}^n \times \mathcal{E}xt_{\mathcal{O}_{X,O}}^n(\mathcal{O}_{X,O}/\mathcal{I}, \mathcal{O}_{X,O}) \rightarrow \mathbb{C}$$

where Ω_X^n is the sheaf of holomorphic differential n -forms on X . Let Σ be the space of algebraic local cohomology classes annihilated by the ideal \mathcal{I} :

$$\Sigma = \{\eta \in \mathcal{H}_{[O]}^n(\mathcal{O}_X) \mid g(z)\eta = 0, g(z) \in \mathcal{I}\}.$$

We can identify the space Σ with $\mathcal{E}xt_{\mathcal{O}_{X,O}}^n(\mathcal{O}_{X,O}/\mathcal{I}, \mathcal{O}_{X,O})$ as a finite dimensional vector spaces over \mathbb{C} . Then, by identifying Milnor algebra $\mathcal{O}_{X,O}/\mathcal{I}$ and $\Omega_{X,O}^n / \mathcal{I} \Omega_{X,O}^n$, we find that the space Σ is the dual space of Milnor algebra.

The space Σ is generated by a single cohomology class over $\mathcal{O}_{X,O}$. For instance, one can take the cohomology class $\sigma_f = \left[\frac{1}{f_1 \dots f_n} \right] \in \mathcal{H}_{[O]}^n(\mathcal{O}_X)$ as a generator over $\mathcal{O}_{X,O}$ of Σ , where the notation $\left[\frac{a}{b_1 \dots b_n} \right]$ for functions $a, b_1 \dots b_n \in \mathcal{O}_{X,O}$ stands for the algebraic local cohomology class associated to the residue symbol $\left[\frac{a}{b_1 \dots b_n} \right] \in \mathcal{E}xt_{\mathcal{O}_{X,O}}^n(\mathcal{O}_{X,O}/\mathcal{I}, \mathcal{O}_{X,O})$.

Let σ be a generator of Σ over $\mathcal{O}_{X,O}$:

$$\Sigma = \mathcal{O}_{X,O}\sigma.$$

Since the algebraic local cohomology group $\mathcal{H}_{[O]}^n(\mathcal{O}_X)$ has a structure of $\mathcal{D}_{X,O}$ -module, we can consider annihilators of σ in $\mathcal{D}_{X,O}$ where \mathcal{D}_X is the sheaf of linear partial differential operators.

Let \mathcal{L}_f be the set of linear partial differential operators of order at most one which annihilate σ :

$$\mathcal{L}_f = \{P = \sum_{j=1}^n a_j(z) \frac{\partial}{\partial z_j} + a_0(z) \mid P\sigma = 0, a_j(z) \in \mathcal{O}_{X,O}, j = 0, 1, \dots, n\}.$$

Let $\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma)$ be the left ideal in $\mathcal{D}_{X,O}$ generated by \mathcal{L}_f ; $\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma) = \mathcal{D}_{X,O}\mathcal{L}_f$. Then $\mathcal{D}_{X,O}/\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma)$ defines a holonomic \mathcal{D}_X module supported at the origin.

Let $P \in \mathcal{L}_f$ be a first order partial differential operator annihilating the algebraic local cohomology class σ . Such an operator has the following property :

Lemma 2.1. — Let σ be a generator of Σ over $\mathcal{O}_{X,O}$. Let P be a first order linear partial differential operator annihilating the cohomology class σ . Then, the space Σ is closed under the action of P , i.e., $P(\Sigma) \subseteq \Sigma$.

It is obvious that the condition whether a given first order differential operator acts on Σ or not depends only on its first order part. We introduce Θ_f to be the set of differential operators of the form $\sum_{j=1}^n a_j(z) \frac{\partial}{\partial z_j}$ acting on Σ . Then, an operator v is in Θ_f if and only if v satisfies the condition $vg(z) \in \mathcal{I}$ for every $g(z) \in \mathcal{I}$, i.e.,

$$\Theta_f = \{v = \sum_{j=1}^n a_j(z) \frac{\partial}{\partial z_j} \mid vg(z) \in \mathcal{I}, \forall g(z) \in \mathcal{I}, a_j(z) \in \mathcal{O}_{X,O}, j = 1, \dots, n\}.$$

Lemma 2.2. — The mapping, from \mathcal{L}_f to Θ_f , which associates the first order part $v_p \in \Theta_f$ to $P \in \mathcal{L}_f$ is a surjective mapping.

3. Solution space of the holonomic system

Let $\sigma \in \Sigma$ be a generator of Σ over $\mathcal{O}_{X,O}$. Let η be another algebraic local cohomology class in Σ and $h(z) \in \mathcal{O}_{X,O}$ a holomorphic function satisfying $\eta = h(z)\sigma$.

It is obvious that, to represent $\eta \in \Sigma$ in the form $\eta = h\sigma$, it suffices to take the modulo class $h \bmod \mathcal{I}$ of the holomorphic function $h(z) \in \mathcal{O}_{X,O}$. Let P be an annihilator of σ in \mathcal{L}_f . Now let us consider the condition that an algebraic local cohomology class $\eta \in \Sigma$ becomes a solution of homogeneous differential equation $P\eta = 0$. An element $v \in \Theta_f$ induces a linear operator acting on $\mathcal{O}_{X,O}/\mathcal{I}$ which is also denoted by v :

$$v : \mathcal{O}_{X,O}/\mathcal{I} \rightarrow \mathcal{O}_{X,O}/\mathcal{I}.$$

For the first order part $v_P = \sum_{j=1}^n a_j(z) \partial/\partial z_j \in \Theta_f$ of an annihilator P , we have

$$v_P h = \sum_{j=1}^n a_j(z) \frac{\partial h}{\partial z_j} = 0 \bmod \mathcal{I}.$$

Let \mathcal{H}_f be a set of modulo classes by \mathcal{I} of functions $h(z)$ that satisfies $vh(z) \in \mathcal{I}$ for $v \in \Theta_f$:

$$\mathcal{H}_f = \{h \in \mathcal{O}_{X,O}/\mathcal{I} \mid vh = 0, \forall v \in \Theta_f\}.$$

Concerning to the algebraic local cohomology solutions of the holonomic system $\mathcal{D}_{X,O}/\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma)$, we have the following result ([3]):

Theorem 3.1. — *Let $f(z)$ be a function defining an isolated singularity at the origin. Let $\sigma \in \Sigma$ be a generator over $\mathcal{O}_{X,O}$ of Σ . Then,*

$$\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X)) = \{h\sigma \mid h \in \mathcal{H}_f\}.$$

The space \mathcal{H}_f does not depend on a choice of a generator σ of Σ . Actually, the space \mathcal{H}_f is completely determined by the ideal \mathcal{I} . Thus, in this sense, the space \mathcal{H}_f is an intrinsic object in the study of the solution space $\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X))$.

For the quasihomogeneous isolated singularity case, we have the following result ([3]) :

Proposition 3.2. — *Let $f(z)$ be a function defining a quasihomogeneous isolated singularity at the origin. Then*

$$\mathcal{H}_f = \text{Span}_{\mathbb{C}}\{1\}.$$

Let $\mathcal{A}nn_{\mathcal{D}_{X,O}}(\sigma)$ be the left ideal in $\mathcal{D}_{X,O}$ of annihilators of the cohomology class σ .

Theorem 3.3 ([3]). — *Let $f = f(z)$ be a function defining an isolated singularity at the origin O and σ a generator of Σ . The following three conditions are equivalent :*

- (i) $\mathcal{O}_{X,O}\langle f, f_1, \dots, f_n \rangle = \mathcal{O}_{X,O}\langle f_1, \dots, f_n \rangle$.
- (ii) $\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma) = \mathcal{A}nn_{\mathcal{D}_{X,O}}(\sigma)$.
- (iii) $\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\mathcal{A}nn_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X)) = \text{Span}_{\mathbb{C}}\{\sigma\}$.

This result can be regarded as a counterpart in the algebraic local cohomology theory of a result by K.Saito on a characterization of quasihomogeneity of singularities ([5]).

In contrast, for a non-quasihomogeneous function $f = f(z)$,

$$(2) \quad \mathcal{O}_{X,O}\langle f_1, \dots, f_n \rangle \neq \mathcal{O}_{X,O}\langle f, f_1, \dots, f_n \rangle$$

and thus

$$\dim \text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\text{Ann}_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X)) \geq 2.$$

It seems natural to expect that the solution space

$$\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\text{Ann}_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X))$$

is related to non-quasihomogeneity of a given hypersurface isolated singularity. Let us consider the structure of the solution spaces for exceptional unimodal singularities which are most typical non-quasihomogeneous singularities.

We have the following result :

Proposition 3.4. — *For a function $f(z)$ defining an exceptional unimodal singularity,*

$$\mathcal{H}_f = \text{Span}_{\mathbb{C}}\{1, f \bmod \mathcal{I}\}.$$

Proposition 3.4 is proved by direct computations for each normal form of exceptional unimodal singularities. We shall explain a method we used for computations in the next section.

We arrive at the following theorem ([6]):

Theorem 3.5. — *Let $f(z)$ be a function defining an exceptional unimodal singularity. Then,*

$$\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\text{Ann}_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X)) = \text{Span}_{\mathbb{C}}\{\sigma, \delta\},$$

where δ is the delta function with support at the origin O .

Proof. — Theorem 3.1 together with Proposition 3.4 yields that the solution space $\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\text{Ann}_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X))$ is spanned by σ and $f\sigma$. Since the ideal quotient $\mathcal{I} : \langle f \rangle$ is the maximal ideal m in $\mathcal{O}_{X,O}$ for any exceptional unimodal singularities, the cohomology class $f\sigma$ is annihilated by m . This implies that $f\sigma = \text{const.}\delta$ where $\delta = \left[\frac{1}{z_1 \cdots z_n} \right]$. It completes the proof. \square

We note here that it is possible to characterize the cohomology class σ attached to an exceptional unimodal (and bimodal) singularity as the solution of a *second order* holonomic system. We shall treat this subject elsewhere.

4. Strategy of computations

In this section, we describe a method for computing the space \mathcal{H}_f which works for the case $f \in \mathbb{Q}[z]$.

We assume that $f \in \mathbb{Q}[z]$ is a function defining an isolated singularity at the origin. For the ideal of the polynomial ring $\mathbb{Q}[z]$ generated by partial derivatives of a given function f , let I denote its primary component at the origin and M be a set of basis monomials of the quotient space $\mathbb{Q}[z]/I$. Put $E = \text{Span}_{\mathbb{Q}}\{e \in M\} \subset \mathbb{Q}[z]$. In this section, let Σ be a space of algebraic local cohomology classes with support at the origin corresponding to $\text{Ext}_{\mathbb{Q}[z]}^n(\mathbb{Q}[z]/I, \mathbb{Q}[z])$. Let σ denote a generator of the space Σ over $\mathbb{Q}[z]$. From a computational point of view, we introduce finite dimensional vector spaces L_f , V_f and H_f that correspond to \mathcal{L}_f , Θ_f and \mathcal{H}_f . Let A_n be the Weyl algebra $\mathbb{Q}[z]\langle\partial/\partial z_1, \dots, \partial/\partial z_n\rangle$. For two first order linear partial differential operators

$$Q = q_1(z) \frac{\partial}{\partial z_1} + \dots + q_n(z) \frac{\partial}{\partial z_n} + q_0(z)$$

and

$$R = r_1(z) \frac{\partial}{\partial z_1} + \dots + r_n(z) \frac{\partial}{\partial z_n} + r_0(z)$$

($q_k(z)$, $r_k(z) \in \mathbb{Q}[z]$), suppose that $r_k(z) \in E$ and $q_k(z) = r_k(z) \pmod{I}$, $k = 1, \dots, n$. Then, one has

$$\begin{aligned} Q &= R + \frac{\partial}{\partial z_1}(q_1(z) - r_1(z)) + \dots + \frac{\partial}{\partial z_n}(q_n(z) - r_n(z)) \\ &\quad + (q_0(z) - r_0(z)) - \left(\frac{\partial q_1(z)}{\partial z_1} + \dots + \frac{\partial q_n(z)}{\partial z_n} - \frac{\partial r_1(z)}{\partial z_1} - \dots - \frac{\partial r_n(z)}{\partial z_n} \right) \\ &= R + (q_0(z) - r_0(z)) - \left(\frac{\partial q_1(z)}{\partial z_1} + \dots + \frac{\partial q_n(z)}{\partial z_n} - \frac{\partial r_1(z)}{\partial z_1} - \dots - \frac{\partial r_n(z)}{\partial z_n} \right) \end{aligned}$$

modulo $A_n I$. If we put

$$s_0(z) = (q_0(z) - r_0(z)) - \left(\frac{\partial q_1(z)}{\partial z_1} + \dots + \frac{\partial q_n(z)}{\partial z_n} - \frac{\partial r_1(z)}{\partial z_1} - \dots - \frac{\partial r_n(z)}{\partial z_n} \right),$$

we have $Q\eta = (R + s_0(z))\eta$ for any $\eta \in \Sigma$. Accordingly, as first order annihilators $\sum_{j=1}^n a_j(z) \frac{\partial}{\partial z_j} + a_0(z)$ of the cohomology class σ , it suffices to take their coefficients $a_i(z)$ ($i = 0, 1, \dots, n$) from the vector space E . Put

$$L_f = \{P = \sum_{j=1}^n a_j(z) \frac{\partial}{\partial z_j} + a_0(z) \mid P\sigma = 0, a_j(z) \in E, j = 0, \dots, n\}.$$

Note that, a cost of computations for constructing annihilators of a given algebraic local cohomology class is expensive in general case ([4]). But, for the zero-dimensional case, the use of the above property provides us an efficient method for computing first order annihilators of a given cohomology class ([2]).

Similarly, we put

$$V_f = \{v = \sum_{j=1}^n a_j(z) \frac{\partial}{\partial z_j} \mid vg \in I, \forall g \in I, a_j(z) \in E, j = 1, \dots, n\}.$$

We have the following:

Lemma 4.1. — *There is a bijective morphism between L_f and V_f :*

$$L_f \cong V_f.$$

Proof. — For any $v \in V_f$, there exist $h(z) \in E$ s.t., $v\sigma = h(z)\sigma$. By putting $a_0(z) = -h(z) \in E$, we have $(v + a_0(z))\sigma = 0$. \square

Since $\mathcal{O}_{X,O}/\mathcal{I} \cong \mathbb{C} \otimes E$ as finite dimensional vector spaces and the defining function f is in $\mathbb{Q}[z]$, it suffices to consider L_f and V_f instead of \mathcal{L}_f and Θ_f . Let H_f be a vector space defined by

$$H_f = \{h(z) \in \mathbb{Q}[z]/I \mid vh(z) = 0, \forall v \in V_f\}.$$

As the space H_f corresponds to the space \mathcal{H}_f , we have the following result :

Proposition 4.2. — $\mathcal{H}_f \cong \mathbb{C} \otimes H_f$.

Proposition 4.2 reduces a determination of the solution space $Hom_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/Ann_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X))$ to the computation of the space H_f . The following procedure, that can be performed by the use of Gröbner basis, computes the space H_f :

- (i) Take basis monomials M of $\mathbb{Q}[z]/I$.
- (ii) Construct V_f with coefficients from E .
- (iii) Choose generators v_1, \dots, v_ℓ of V_f over $\mathbb{Q}[z]/I$.
- (iv) Compute the solution space H_f by solving simultaneous differential equations $v_1 h(z) = \dots = v_\ell h(z) = 0$.

We present detailed data of computations for the E_{12} singularity for illustration.

Example 4.3 (E_{12} singularity). — Let f be a two variables polynomial $f = x^3 + y^7 + axy^5$ with a parameter a . This is the normal form of E_{12} singularities. The primary decomposition of the ideal generated by partial derivatives $f_x = 3x^2 + ay^5$ and $f_y = 7y^6 + 5axy^4$ of f is $I \cap \langle 25a^3y + 147, 3125a^7x + 151263 \rangle$ where I is the primary component at the origin and its Gröbner basis with respect to the lexicographical ordering $x \succ y$ is

$$\{y^8, 7y^6 + 5ay^4x, 3x^2 + ay^5\}.$$

As the basis of the vector space $\mathbb{Q}[x,y]/I$, we use the following monomials:

$$M = \{y^5x, y^4x, y^3x, y^2x, yx, x, y^5, y^4, y^3, y^2, y, 1\}$$

which are constructed from the quasihomogeneous part $x^3 + y^7$ of f .

A basis of Σ is given by the following 12 cohomology classes which constitute the dual basis of the monomial basis M :

$$\begin{aligned} & \left[\frac{1}{y^6x^2} + a\left(-\frac{5}{7}\frac{1}{y^8x} - \frac{1}{3}\frac{1}{yx^4}\right) + \frac{5}{21}a^2\frac{1}{y^3x^3} \right], \left[\frac{1}{y^5x^2} - \frac{5}{7}a\frac{1}{y^7x} + \frac{5}{21}a^2\frac{1}{y^2x^3} \right], \\ & \left[\frac{1}{y^4x^2} \right], \left[\frac{1}{y^6x} - \frac{1}{3}a\frac{1}{yx^3} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{y^5x} \right], \left[\frac{1}{y^2x^2} \right], \\ & \left[\frac{1}{y^4x} \right], \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right]. \end{aligned}$$

where $[]$ is a relative Čech cohomology representation of an algebraic local cohomology class with respect to the standard covering $\{X, \{x \neq 0\}, \{y \neq 0\}\}$. The first cohomology class in the above list, denoted by τ , generates Σ over $\mathbb{Q}[x, y]$. In the above 12 cohomology classes, only τ is not annihilated by f , i.e., $f\tau \neq 0$ and $f\eta = 0$ for other cohomology classes η . If one uses τ , the above classes can be represented in the following way:

$$\begin{aligned} & \tau, \\ & y\tau, \\ & \{y^2 + \frac{5}{7}a(x - \frac{5}{21}a^2(y^3 + \frac{5}{7}a(xy - \frac{5}{21}a^2(y^4 + \frac{5}{7}a(y^2x - \frac{5}{21}a^2(y^5 + \frac{5}{7}ay^3x)))))))\}\tau, \\ & \{x - \frac{5}{21}a^2(y^3 + \frac{5}{7}a(xy - \frac{5}{21}a^2(y^4 + \frac{5}{7}a(y^2x - \frac{5}{21}a^2(y^5 + \frac{5}{7}ay^3x)))))\}\tau, \\ & \{y^3 + \frac{5}{7}a(xy - \frac{5}{21}a^2(y^4 + \frac{5}{7}a(y^2x - \frac{5}{21}a^2(y^5 + \frac{5}{7}ay^3x))))\}\tau, \\ & \{xy - \frac{5}{21}a^2(y^4 + \frac{5}{7}a(y^2x - \frac{5}{21}a^2(y^5 + \frac{5}{7}ay^3x)))\}\tau, \\ & \{y^4 + \frac{5}{7}a(y^2x - \frac{5}{21}a^2(y^5 + \frac{5}{7}ay^3x))\}\tau, \\ & \{y^2x - \frac{5}{21}a^2(y^5 + \frac{5}{7}ay^3x)\}\tau, \\ & \{y^5 + \frac{5}{7}ay^3x\}\tau, \\ & y^3x\tau, \\ & y^4x\tau, \\ & y^5x\tau. \end{aligned}$$

The vector space V_f is spanned by the following 14 operators:

$$\begin{aligned} & 252yx\frac{\partial}{\partial x} - 30ax\frac{\partial}{\partial y} - 35a^2y^4\frac{\partial}{\partial x}, 63y^2\frac{\partial}{\partial y} + 15ax\frac{\partial}{\partial y} + 25a^2y^4\frac{\partial}{\partial x}, \\ & 7y^2x\frac{\partial}{\partial x} - 2ayx\frac{\partial}{\partial y}, 7y^3\frac{\partial}{\partial y} + 5ayx\frac{\partial}{\partial y}, 6yx\frac{\partial}{\partial y} - 5ay^5\frac{\partial}{\partial x}, \\ & y^4\frac{\partial}{\partial y}, y^3x\frac{\partial}{\partial x}, y^2x\frac{\partial}{\partial y}, y^5\frac{\partial}{\partial y}, y^4x\frac{\partial}{\partial x}, y^3x\frac{\partial}{\partial y}, y^5x\frac{\partial}{\partial x}, y^4x\frac{\partial}{\partial y}, y^5x\frac{\partial}{\partial y}. \end{aligned}$$

It is easy to see that first three operators generates V_f over $\mathbb{Q}[x, y]/I$. Thus by solving the system of differential equations

$$\begin{cases} (252yx\frac{\partial}{\partial x} - 30ax\frac{\partial}{\partial y} - 35a^2y^4\frac{\partial}{\partial x})h(x, y) = 0, \\ (63y^2\frac{\partial}{\partial y} + 15ax\frac{\partial}{\partial y} + 25a^2y^4\frac{\partial}{\partial x})h(x, y) = 0, \\ (7y^2x\frac{\partial}{\partial x} - 2ayx\frac{\partial}{\partial y})h(x, y) = 0, \end{cases}$$

we have

$$H_f = \text{Span}_{\mathbb{Q}}\{1, y^5x\}.$$

Let σ be a generator of Σ over $\mathbb{Q}[z]$. Since the cohomology class σ contains τ as its non-zero direct summand, $xy^5\sigma \neq 0$. As $x(y^5x) \in I$ and $y(y^5x) \in I$, we have

$$\text{Hom}_{\mathcal{D}_{X,O}}(\mathcal{D}_{X,O}/\text{Ann}_{\mathcal{D}_{X,O}}^{(1)}(\sigma), \mathcal{H}_{[O]}^n(\mathcal{O}_X)) = \text{Span}_{\mathbb{C}}\{\sigma, \delta\},$$

where δ stands for the delta function $[1/xy]$ with support at the origin. One can also verify the above result by direct computation. Actually, since

$$\begin{cases} y^5x \left[\frac{1}{y^6x^2} + a(-\frac{5}{7}\frac{1}{y^8x} - \frac{1}{3}\frac{1}{yx^4}) + \frac{5}{21}a^2\frac{1}{y^3x^3} \right] = \left[\frac{1}{xy} \right], \\ y^5x \left[\frac{1}{y^5x^2} - \frac{5}{7}a\frac{1}{y^7x} + \frac{21}{5}a^2\frac{1}{y^2x^3} \right] = 0, \\ \dots, \\ y^5x \left[\frac{1}{yx} \right] = 0, \end{cases}$$

we have $y^5x\sigma = c[1/xy]$ with some non-zero constant c .

5. Computations for normal forms

In this section, we give results of computations which we used to derive Proposition 3.4. We apply the procedure introduced in the above section to the following normal forms of non-quasihomogeneous unimodal singularities classified by V.I.Arnold ([1]):

2variables

| | | |
|----------|-----------------------------------|---|
| E_{12} | $f(x, y) = x^3 + y^7 + axy^5$ | 3variables |
| E_{13} | $f(x, y) = x^3 + xy^5 + ay^8$ | $Q_{10} : f(x, y, z) = x^3 + y^4 + yz^2 + axy^3$ |
| E_{14} | $f(x, y) = x^3 + y^8 + axy^6$ | $Q_{11} : f(x, y, z) = x^3 + y^2z + xz^3 + az^5$ |
| Z_{11} | $f(x, y) = x^3y + y^5 + axy^4$ | $Q_{12} : f(x, y, z) = x^3 + y^5 + yz^2 + axy^4$ |
| Z_{12} | $f(x, y) = x^3y + xy^4 + ax^2y^3$ | $S_{11} : f(x, y, z) = x^4 + y^2z + xz^2 + ax^3$ |
| Z_{13} | $f(x, y) = x^3y + y^6 + axy^5$ | $S_{12} : f(x, y, z) = x^2y + y^2z + xz^3 + az^5$ |
| W_{12} | $f(x, y) = x^4 + y^5 + ax^2y^3$ | $U_{12} : f(x, y, z) = x^3 + y^3 + z^4 + axyz^2$ |
| W_{13} | $f(x, y) = x^4 + xy^4 + ay^6$ | |

Let $\Delta = H_f \cap m$ where m is the maximal ideal in $\mathbb{Q}[z]$. Then, $\text{Span}_{\mathbb{Q}}\{1\} \oplus \Delta = H_f$ holds. From the output of computations, we find that $\dim \Delta = 1$. Data in the following list is an explicit form of a generator of each Δ :

a generator of Δ

| | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|
| E_{12} | E_{13} | E_{14} | Z_{11} | Z_{12} | Z_{13} | W_{12} | W_{13} |
| y^5x | y^8 | y^6x | y^4x | y^6 | y^5x | y^3x^2 | y^6 |
| Q_{10} | Q_{11} | Q_{12} | S_{11} | S_{12} | U_{12} | | |
| y^3x | z^5 | y^4x | y^2x^2 | z^5 | z^2yx | | |

Thus, for each case, $\dim H_f = 2$. Combining this result and the fact that $\{1, f \bmod \mathcal{I}\} \subseteq \mathcal{H}_f$ for non-quasihomogeneous singularities ([3]), one can prove Proposition 3.4, i.e.,

$$\mathcal{H}_f = \{1, f \bmod \mathcal{I}\}.$$

5.1. data. — In the rest of this paper, we give the following resulting data of our computation:

- a Gröbner basis of the ideal I in $\mathbb{Q}[z]$ (with a parameter a),
- basis monomials M of $\mathbb{Q}[z]/I$ constructed from the quasihomogeneous part of f ,
- a basis of Σ which constitutes the dual basis of M w.r.t. Grothendieck local duality,

and

- a basis of V_f whose coefficients are in $E = \text{Span}_{\mathbb{Q}}\{e \in M\}$ and the number k of generators over $\mathbb{Q}[z]/I$ in the list. The first k operators are generators.

Note here that the basis monomials M of Z_{12} , Q_{10} , S_{11} and S_{12} are different from that in [1]. The first two operators (resp. three operators) in the list of V_f and I are generators over A_n for two variables (resp. three variables) case.

5.1.1. $E_{12} : x^3 + y^7 + axy^5$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^8, 7y^6 + 5ay^4x, 3x^2 + ay^5$$

Basis monomials M :

$$y^5x, y^4x, y^3x, y^5, y^2x, y^4, yx, y^3, x, y^2, y, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{y^6x^2} + a(-\frac{5}{7}\frac{1}{y^8x} - \frac{1}{3}\frac{1}{yx^4}) + \frac{5}{21}a^2\frac{1}{y^3x^3} \right], \left[\frac{1}{y^5x^2} - \frac{5}{7}a\frac{1}{y^7x} + \frac{5}{21}a^2\frac{1}{y^2x^3} \right], \\ & \left[\frac{1}{y^4x^2} \right], \left[\frac{1}{y^6x} - \frac{1}{3}a\frac{1}{yx^3} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{y^5x} \right], \left[\frac{1}{y^2x^2} \right], \left[\frac{1}{y^4x} \right], \\ & \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right] \end{aligned}$$

A basis of V_f : (# of generators : 3)

$$\begin{aligned} & 252yx\frac{\partial}{\partial x} - 30ax\frac{\partial}{\partial y} - 35a^2y^4\frac{\partial}{\partial x}, 63y^2\frac{\partial}{\partial y} + 15ax\frac{\partial}{\partial y} + 25a^2y^4\frac{\partial}{\partial x}, \\ & 7y^2x\frac{\partial}{\partial x} - 2ayx\frac{\partial}{\partial y}, 7y^3\frac{\partial}{\partial y} + 5ayx\frac{\partial}{\partial y}, 6yx\frac{\partial}{\partial y} - 5ay^5\frac{\partial}{\partial x}, \\ & y^4\frac{\partial}{\partial y}, y^3x\frac{\partial}{\partial x}, y^2x\frac{\partial}{\partial y}, y^5\frac{\partial}{\partial y}, y^4x\frac{\partial}{\partial x}, y^3x\frac{\partial}{\partial y}, y^5x\frac{\partial}{\partial x}, y^4x\frac{\partial}{\partial y}, y^5x\frac{\partial}{\partial y} \end{aligned}$$

5.1.2. $E_{13} : x^3 + xy^5 + ay^8$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^9, 5y^4x + 8ay^7, 3x^2 + y^5$$

Basis monomials M :

$$y^8, y^7, y^6, y^3x, y^5, y^2x, y^4, yx, y^3, x, y^2, y, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{y^9x} - \frac{1}{3} \frac{1}{y^4x^3} + a\left(-\frac{8}{5} \frac{1}{y^6x^2} + \frac{8}{15} \frac{1}{yx^4}\right) \right], \left[\frac{1}{y^8x} - \frac{1}{3} \frac{1}{y^3x^3} - \frac{8}{5} a \frac{1}{y^5x^2} \right], \\ & \left[\frac{1}{y^7x} - \frac{1}{3} \frac{1}{y^2x^3} \right], \left[\frac{1}{y^4x^2} \right], \left[\frac{1}{y^6x} - \frac{1}{3} \frac{1}{yx^3} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{y^5x} \right], \\ & \left[\frac{1}{y^2x^2} \right], \left[\frac{1}{y^4x} \right], \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right] \end{aligned}$$

A basis of V_f : (# of generators : 3)

$$\begin{aligned} & 125yx \frac{\partial}{\partial x} + 50y^2 \frac{\partial}{\partial y} - 40ay^4 \frac{\partial}{\partial x} + 192a^2y^2x \frac{\partial}{\partial x}, 20y^4 \frac{\partial}{\partial x} + 15x \frac{\partial}{\partial y} - 156ay^2x \frac{\partial}{\partial x}, \\ & 5y^2x \frac{\partial}{\partial x} + 2y^3 \frac{\partial}{\partial y}, yx \frac{\partial}{\partial y} - 4ay^3x \frac{\partial}{\partial x}, 5y^5 \frac{\partial}{\partial x} - 24ay^3x \frac{\partial}{\partial x}, 5y^3x \frac{\partial}{\partial x} + 2y^4 \frac{\partial}{\partial y}, \\ & y^2x \frac{\partial}{\partial y}, y^6 \frac{\partial}{\partial x}, y^5 \frac{\partial}{\partial y}, y^3x \frac{\partial}{\partial y}, y^7 \frac{\partial}{\partial x}, y^6 \frac{\partial}{\partial y}, y^8 \frac{\partial}{\partial x}, y^7 \frac{\partial}{\partial y}, y^8 \frac{\partial}{\partial y} \end{aligned}$$

5.1.3. $E_{14} : x^3 + y^8 + axy^6$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^9, 4y^7 + 3ay^5x, 3x^2 + ay^6$$

Basis monomials M :

$$y^6x, y^5x, y^4x, y^6, y^3x, y^5, y^2x, y^4, yx, y^3, x, y^2, y, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{y^7x^2} + a\left(-\frac{3}{4} \frac{1}{y^9x} - \frac{1}{3} \frac{1}{yx^4}\right) + \frac{1}{4} a^2 \frac{1}{y^3x^3} \right], \left[\frac{1}{y^6x^2} - \frac{3}{4} a \frac{1}{y^8x} + \frac{1}{4} a^2 \frac{1}{y^2x^3} \right], \\ & \left[\frac{1}{y^5x^2} \right], \left[\frac{1}{y^7x} - \frac{1}{3} a \frac{1}{yx^3} \right], \left[\frac{1}{y^4x^2} \right], \left[\frac{1}{y^6x} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{y^5x} \right], \\ & \left[\frac{1}{y^2x^2} \right], \left[\frac{1}{y^4x} \right], \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right] \end{aligned}$$

A basis of V_f : (# of generators : 3)

$$\begin{aligned} & 28yx \frac{\partial}{\partial x} - 3ax \frac{\partial}{\partial y} - 4a^2y^5 \frac{\partial}{\partial x}, 28y^2 \frac{\partial}{\partial y} + 6ax \frac{\partial}{\partial y} + 15a^2y^5 \frac{\partial}{\partial x}, \\ & 4y^2x \frac{\partial}{\partial x} - ayx \frac{\partial}{\partial y}, 4y^3 \frac{\partial}{\partial y} + 3ayx \frac{\partial}{\partial y}, yx \frac{\partial}{\partial y} - ay^6 \frac{\partial}{\partial x}, \\ & y^4 \frac{\partial}{\partial y}, y^3x \frac{\partial}{\partial x}, y^2x \frac{\partial}{\partial y}, y^5 \frac{\partial}{\partial y}, y^4x \frac{\partial}{\partial x}, y^3x \frac{\partial}{\partial y}, \end{aligned}$$

$$y^6 \frac{\partial}{\partial y}, y^5 x \frac{\partial}{\partial x}, y^4 x \frac{\partial}{\partial y}, y^6 x \frac{\partial}{\partial x}, y^5 x \frac{\partial}{\partial y}, y^6 x \frac{\partial}{\partial y}$$

5.1.4. $Z_{11} : x^3 y + y^5 + a x y^4$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^6, 15y^5 + 11ay^4x, 3yx^2 + ay^4, x^3 + 5y^4 + 4ay^3x$$

Basis monomials M :

$$y^4 x, y^3 x, y^4, y^2 x, y^3, x^2, yx, y^2, x, y, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{y^5 x^2} - 5 \frac{1}{y x^5} + a \left(-\frac{11}{15} \frac{1}{y^6 x} - \frac{1}{3} \frac{1}{y^2 x^4} \right) + \frac{11}{45} a^2 \frac{1}{y^3 x^3} \right], \\ & \left[\frac{1}{y^4 x^2} - \frac{4}{5} a \frac{1}{y^5 x} + \frac{4}{15} a^2 \frac{1}{y^2 x^3} \right], \left[\frac{1}{y^5 x} - 5 \frac{1}{y x^4} - \frac{1}{3} a \frac{1}{y^2 x^3} \right], \\ & \left[\frac{1}{y^3 x^2} \right], \left[\frac{1}{y^4 x} \right], \left[\frac{1}{y x^3} \right], \left[\frac{1}{y^2 x^2} \right], \left[\frac{1}{y^3 x} \right], \left[\frac{1}{y x^2} \right], \left[\frac{1}{y^2 x} \right], \left[\frac{1}{y x} \right] \end{aligned}$$

A basis of V_f : (# of generators : 4)

$$\begin{aligned} & 15yx \frac{\partial}{\partial x} - a(61x^2 \frac{\partial}{\partial x} + 48yx \frac{\partial}{\partial y}), 15y^2 \frac{\partial}{\partial y} + a(108x^2 \frac{\partial}{\partial x} + 83yx \frac{\partial}{\partial y}), \\ & 60x^2 \frac{\partial}{\partial x} + 45yx \frac{\partial}{\partial y} + ax^2 \frac{\partial}{\partial y}, 5y^3 \frac{\partial}{\partial x} + 2x^2 \frac{\partial}{\partial y}, y^3 \frac{\partial}{\partial y}, y^2 x \frac{\partial}{\partial x}, \\ & y^2 x \frac{\partial}{\partial y}, y^4 \frac{\partial}{\partial x}, y^4 \frac{\partial}{\partial y}, y^3 x \frac{\partial}{\partial x}, y^3 x \frac{\partial}{\partial y}, y^4 x \frac{\partial}{\partial x}, y^4 x \frac{\partial}{\partial y} \end{aligned}$$

5.1.5. $Z_{12} : x^3 y + xy^4 + ax^2 y^3$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^7, 33y^4x - 7ay^6, 3yx^2 + y^4 + 2ay^3x, 33x^3 + 132y^3x - 33ay^5 - 14a^3y^6$$

Basis monomials M :

$$y^6, y^5, y^3 x, y^4, y^2 x, y^3, x^2, yx, y^2, y, x, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{y^7 x} - \frac{1}{3} \frac{1}{y^4 x^3} + \frac{4}{3} \frac{1}{y x^5} + a \left(\frac{6}{11} \frac{1}{y^2 x^4} - \frac{18}{11} \frac{1}{y^5 x^2} \right) \right], \\ & \left[\frac{1}{y^6 x} - \frac{1}{3} \frac{1}{y^3 x^3} - \frac{3}{2} a \frac{1}{y^4 x^2} \right], \left[\frac{1}{y^4 x^2} - 4 \frac{1}{y x^4} \right], \left[\frac{1}{y^5 x} - \frac{1}{3} \frac{1}{y^2 x^3} \right], \\ & \left[\frac{1}{y^3 x^2} \right], \left[\frac{1}{y^4 x} \right], \left[\frac{1}{y x^3} \right], \left[\frac{1}{y^2 x^2} \right], \left[\frac{1}{y^3 x} \right], \left[\frac{1}{y x^2} \right], \left[\frac{1}{y^2 x} \right], \left[\frac{1}{y x} \right] \end{aligned}$$

A basis of V_f : (# of generators : 3)

$$2904x \frac{\partial}{\partial y} + 3872y^2 \frac{\partial}{\partial x} + 5324axy \frac{\partial}{\partial x} + a^2(260y^3 + 725x^2) \frac{\partial}{\partial x},$$

$$\begin{aligned}
& 792y^2 \frac{\partial}{\partial y} + 1188xy \frac{\partial}{\partial x} + a(-68y^3 + 225x^2) \frac{\partial}{\partial x}, (8y^3 - 9x^2) \frac{\partial}{\partial x} + 12axy^2 \frac{\partial}{\partial x}, \\
& 72xy \frac{\partial}{\partial y} + (8y^3 + 99x^2) \frac{\partial}{\partial x}, (-8y^3 + 9x^2) \frac{\partial}{\partial x} + 9ax^2 \frac{\partial}{\partial y}, (-8y^3 + 9x^2) \frac{\partial}{\partial x} + 8ay^3 \frac{\partial}{\partial y}, \\
& y^4 \frac{\partial}{\partial x}, y^2x \frac{\partial}{\partial y}, y^3x \frac{\partial}{\partial x}, y^4 \frac{\partial}{\partial y}, y^5 \frac{\partial}{\partial x}, y^3x \frac{\partial}{\partial y}, y^5 \frac{\partial}{\partial y}, y^6 \frac{\partial}{\partial x}, y^6 \frac{\partial}{\partial y}
\end{aligned}$$

5.1.6. $Z_{13} : x^3y + y^6 + axy^5$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^7, 9y^6 + 7ay^5x, 3yx^2 + ay^5, x^3 + 6y^5 + 5ay^4x$$

Basis monomials M :

$$y^5x, y^4x, y^5, y^3x, y^4, y^2x, x^2, y^3, yx, y^2, x, y, 1$$

A basis of Σ :

$$\begin{aligned}
& \left[\frac{1}{y^6x^2} - 6 \frac{1}{yx^5} + a \left(-\frac{7}{9} \frac{1}{y^7x} + \frac{7}{27} \frac{1}{y^3x^3} - \frac{1}{3} \frac{1}{y^2x^4} \right) \right], \left[\frac{1}{y^5x^2} - \frac{5}{6}a \frac{1}{y^6x} + \frac{5}{18}a^2 \frac{1}{y^2x^3} \right], \\
& \left[\frac{1}{y^5x^2} - 5a \frac{1}{yx^4} \right], \left[\frac{1}{y^4x^2} \right], \left[\frac{1}{y^5x} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{yx^3} \right], \left[\frac{1}{y^4x} \right], \\
& \left[\frac{1}{y^2x^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right]
\end{aligned}$$

A basis of V_f : (# of generators : 4)

$$\begin{aligned}
& 9yx \frac{\partial}{\partial x} - a(23x^2 \frac{\partial}{\partial x} + 15yx \frac{\partial}{\partial y}), 9y^2 \frac{\partial}{\partial y} + a(60x^2 \frac{\partial}{\partial x} + 37yx \frac{\partial}{\partial y}), \\
& 45x^2 \frac{\partial}{\partial x} + 27yx \frac{\partial}{\partial y} + ax^2 \frac{\partial}{\partial y}, 3y^4 \frac{\partial}{\partial x} + x^2 \frac{\partial}{\partial y}, y^3 \frac{\partial}{\partial y}, y^2x \frac{\partial}{\partial x}, \\
& y^2x \frac{\partial}{\partial y}, y^4 \frac{\partial}{\partial y}, y^3x \frac{\partial}{\partial x}, y^5 \frac{\partial}{\partial x}, y^3x \frac{\partial}{\partial y}, y^5 \frac{\partial}{\partial y}, y^4x \frac{\partial}{\partial x}, y^4x \frac{\partial}{\partial y}, y^5x \frac{\partial}{\partial x}, y^5x \frac{\partial}{\partial y}
\end{aligned}$$

5.1.7. $W_{12} : x^4 + y^5 + ax^2y^3$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^6, y^4x, 5y^4 + 3ay^2x^2, 2x^3 + ay^3x$$

Basis monomials M :

$$y^3x^2, y^2x^2, y^3x, yx^2, y^2x, y^3, x^2, yx, y^2, x, y, 1$$

A basis of Σ :

$$\begin{aligned}
& \left[\frac{1}{y^4x^3} + a \left(-\frac{3}{5} \frac{1}{y^6x} - \frac{1}{2} \frac{1}{yx^5} \right) \right], \left[\frac{1}{y^3x^3} - \frac{3}{5}a \frac{1}{y^5x} \right], \left[\frac{1}{y^4x^2} - \frac{1}{2}a \frac{1}{yx^4} \right], \\
& \left[\frac{1}{y^2x^3} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{y^4x} \right], \left[\frac{1}{yx^3} \right], \left[\frac{1}{y^2x^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right]
\end{aligned}$$

A basis of V_f : (# of generators : 4)

$$yx \frac{\partial}{\partial x} + y^2 \frac{\partial}{\partial y}, 3x^2 \frac{\partial}{\partial x} + 2yx \frac{\partial}{\partial y}, 10y^2 \frac{\partial}{\partial y} + 3ax^2 \frac{\partial}{\partial y}, 2yx \frac{\partial}{\partial y} - ay^3 \frac{\partial}{\partial x},$$

$$\begin{aligned} & y^3 \frac{\partial}{\partial y}, y^2 x \frac{\partial}{\partial x}, y^2 x \frac{\partial}{\partial y}, yx^2 \frac{\partial}{\partial x}, yx^2 \frac{\partial}{\partial y}, \\ & y^3 x \frac{\partial}{\partial x}, y^3 x \frac{\partial}{\partial y}, y^2 x^2 \frac{\partial}{\partial x}, y^2 x^2 \frac{\partial}{\partial y}, y^3 x^2 \frac{\partial}{\partial x}, y^3 x^2 \frac{\partial}{\partial y} \end{aligned}$$

5.1.8. $W_{13} : x^4 + xy^4 + ay^6$. — The Gröbner basis of I w.r.t. the lexicographic ordering $x \succ y$:

$$y^7, 2y^3x + 3ay^5, 4x^3 + y^4$$

Basis monomials M :

$$y^6, y^5, y^2x^2, y^4, yx^2, y^2x, y^3, x^2, yx, y^2, x, y, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{y^7x} - \frac{1}{4} \frac{1}{y^3x^4} + a(-\frac{3}{2} \frac{1}{y^5x^2} + \frac{3}{8} \frac{1}{yx^5}) \right], \left[\frac{1}{y^6x} - \frac{1}{4} \frac{1}{y^2x^4} - \frac{3}{2}a \frac{1}{y^4x^2} \right], \\ & \left[\frac{1}{y^3x^3} \right], \left[\frac{1}{y^5x} - \frac{1}{4} \frac{1}{yx^4} \right], \left[\frac{1}{y^2x^3} \right], \left[\frac{1}{y^3x^2} \right], \left[\frac{1}{y^4x} \right], \\ & \left[\frac{1}{yx^3} \right], \left[\frac{1}{y^2x^2} \right], \left[\frac{1}{y^3x} \right], \left[\frac{1}{yx^2} \right], \left[\frac{1}{y^2x} \right], \left[\frac{1}{yx} \right] \end{aligned}$$

A basis of V_f : (# of generators : 4)

$$\begin{aligned} & 4yx \frac{\partial}{\partial x} + 3y^2 \frac{\partial}{\partial y} - 3ay^3 \frac{\partial}{\partial x}, 8x^2 \frac{\partial}{\partial x} - 9ay^3 \frac{\partial}{\partial y}, 2yx \frac{\partial}{\partial y} + 3ay^3 \frac{\partial}{\partial y}, 3y^3 \frac{\partial}{\partial x} + 4x^2 \frac{\partial}{\partial y}, \\ & 3y^3 \frac{\partial}{\partial y} + 4y^2x \frac{\partial}{\partial x}, y^2x \frac{\partial}{\partial y}, yx^2 \frac{\partial}{\partial x}, yx^2 \frac{\partial}{\partial y}, y^4 \frac{\partial}{\partial x}, y^4 \frac{\partial}{\partial y}, \\ & y^2x^2 \frac{\partial}{\partial x}, y^2x^2 \frac{\partial}{\partial y}, y^5 \frac{\partial}{\partial x}, y^5 \frac{\partial}{\partial y}, y^6 \frac{\partial}{\partial x}, y^6 \frac{\partial}{\partial y} \end{aligned}$$

5.1.9. $Q_{10} : x^3 + y^4 + yz^2 + axy^3$. — The Gröbner basis of I w.r.t. the lexicographic ordering $z \succ x \succ y$:

$$y^5, 4y^4 + 3axy^3, 3x^2 + ay^3, yz, z^2 + 4y^3 + 3axy^2$$

Basis monomials M :

$$y^3x, y^2x, y^3, zx, yx, y^2, z, x, y, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{zy^4x^2} - 4 \frac{1}{z^3yx^2} + a(-\frac{3}{4} \frac{1}{zy^5x} - \frac{1}{3} \frac{1}{zyx^4}) + \frac{1}{4}a^2 \frac{1}{zy^2x^3} \right], \\ & \left[\frac{1}{zy^3x^2} - \frac{3}{4}a \frac{1}{zy^4x} - \frac{1}{4}a^2 \frac{1}{zyx^3} \right], \left[\frac{1}{zy^4x} - 4 \frac{1}{z^3yx} + \frac{1}{3}a \frac{1}{zyx^3} \right], \\ & \left[\frac{1}{z^2yx^2} \right], \left[\frac{1}{zy^2x^2} \right], \left[\frac{1}{zy^3x} \right], \left[\frac{1}{z^2yx} \right], \left[\frac{1}{zyx^2} \right], \left[\frac{1}{zy^2x} \right], \left[\frac{1}{zyx} \right] \end{aligned}$$

A basis of V_f : (# of generators : 5)

$$z \frac{\partial}{\partial y} + 4y^2 \frac{\partial}{\partial z} + 3axy \frac{\partial}{\partial z}, 8xy \frac{\partial}{\partial x} + 3axz \frac{\partial}{\partial z} - 2a^2y^3 \frac{\partial}{\partial x}, 2xy \frac{\partial}{\partial y} - 3xz \frac{\partial}{\partial z} + ay^3 \frac{\partial}{\partial x},$$

$$16y^2 \frac{\partial}{\partial y} + 24xy \frac{\partial}{\partial x} + 3axz \frac{\partial}{\partial z}, xz \frac{\partial}{\partial x}, 4y^3 \frac{\partial}{\partial z} + 3axy^2 \frac{\partial}{\partial z}, 16y^3 \frac{\partial}{\partial z} - 3axz \frac{\partial}{\partial y}, \\ xy^2 \frac{\partial}{\partial x}, y^3 \frac{\partial}{\partial y}, xy^2 \frac{\partial}{\partial y}, xy^3 \frac{\partial}{\partial z}, xy^3 \frac{\partial}{\partial x}, xy^3 \frac{\partial}{\partial y},$$

5.1.10. $Q_{11} : x^3 + y^2z + xz^3 + az^5$. — The Gröbner basis of I w.r.t. the lexicographic ordering $y \succ x \succ z$:

$$z^6, 3xz^3 + 5az^5, z^3 + 3x^2, yz, 3xz^2 + y^2 + 5az^4$$

Basis monomials M :

$$z^5, z^4, z^2x, yx, z^3, zx, z^2, y, x, z, 1$$

A basis of Σ :

$$\left[\frac{1}{z^6yx} + \frac{1}{zy^3x^2} - \frac{1}{3} \frac{1}{z^3yx^3} + a(-\frac{5}{3} \frac{1}{z^4yx^2} + \frac{5}{9} \frac{1}{zyx^4}) \right], \\ \left[\frac{1}{z^5yx} - \frac{1}{3} \frac{1}{z^2yx^3} - 5a \frac{1}{zy^3x} \right], \left[\frac{1}{zy^3x} - \frac{1}{3} \frac{1}{z^3yx^2} \right], \left[\frac{1}{zy^2x^2} \right], \\ \left[\frac{1}{z^4yx} - \frac{1}{3} \frac{1}{zyx^3} \right], \left[\frac{1}{z^2yx^2} \right], \left[\frac{1}{z^3yx} \right], \left[\frac{1}{zy^2x} \right], \left[\frac{1}{zyx^2} \right], \left[\frac{1}{z^2yx} \right], \left[\frac{1}{zyx} \right]$$

A basis of V_f : (# of generators : 5)

$$18xz \frac{\partial}{\partial x} + 12z^2 \frac{\partial}{\partial z} + 15axz \frac{\partial}{\partial z} + 25a^2 z^3 \frac{\partial}{\partial z}, y \frac{\partial}{\partial z} + (5az^3 + 3xz) \frac{\partial}{\partial y}, \\ 3xz \frac{\partial}{\partial z} + 3xy \frac{\partial}{\partial y} + 5az^3 \frac{\partial}{\partial z}, 6z^3 \frac{\partial}{\partial x} + 9xz \frac{\partial}{\partial z} + 35az^3 \frac{\partial}{\partial z}, xy \frac{\partial}{\partial x}, 3xz^2 \frac{\partial}{\partial y} + 5axy \frac{\partial}{\partial z}, \\ 3xz^2 \frac{\partial}{\partial x} + 2z^3 \frac{\partial}{\partial z}, z^4 \frac{\partial}{\partial y} - xy \frac{\partial}{\partial z}, xz^2 \frac{\partial}{\partial z}, z^4 \frac{\partial}{\partial x}, z^5 \frac{\partial}{\partial z}, z^5 \frac{\partial}{\partial x}, z^5 \frac{\partial}{\partial y},$$

5.1.11. $Q_{12} : x^3 + y^5 + yz^2 + axy^4$. — The Gröbner basis of I w.r.t. the lexicographic ordering $z \succ x \succ y$:

$$y^6, 5y^5 + 4axy^4, 3x^2 + ay^4, yz, z^2 + 5y^4 + 4axy^3$$

Basis monomials M :

$$y^4x, y^3x, y^4, y^2x, zx, y^3, yx, y^2, z, x, y, 1$$

A basis of Σ :

$$\left[\frac{1}{zy^5x^2} - 5 \frac{1}{z^3yx^2} + a(-\frac{4}{5} \frac{1}{zy^6x} - \frac{1}{3} \frac{1}{zyx^4}) + \frac{4}{15} a^2 \frac{1}{zy^2x^3} \right], \left[\frac{1}{zy^4x^2} - 4a \frac{1}{z^3yx} \right], \\ \left[\frac{1}{zy^5x} - 5 \frac{1}{z^3yx} - \frac{a}{3} \frac{1}{zyx^3} \right], \left[\frac{1}{zy^3x^2} \right], \left[\frac{1}{z^2yx^2} \right], \left[\frac{1}{zy^4x} \right] \left[\frac{1}{zy^2x^2} \right], \\ \left[\frac{1}{zy^3x} \right], \left[\frac{1}{z^2yx} \right], \left[\frac{1}{zyx^2} \right], \left[\frac{1}{zy^2x} \right], \left[\frac{1}{zyx} \right],$$

A basis of V_f : (# of generators : 5)

$$15xy \frac{\partial}{\partial x} - 3axy \frac{\partial}{\partial y} - 2a^2 y^4 \frac{\partial}{\partial x}, 5y^2 \frac{\partial}{\partial y} + axy \frac{\partial}{\partial y} + 2a^2 y^4 \frac{\partial}{\partial x}, z \frac{\partial}{\partial y} + 5y^3 \frac{\partial}{\partial z} + 4axy^2 \frac{\partial}{\partial z},$$

$$5y^2 \frac{\partial}{\partial y} + a(4xy \frac{\partial}{\partial y} + 6xz \frac{\partial}{\partial z}), xz \frac{\partial}{\partial x}, y^3 \frac{\partial}{\partial y}, xy^2 \frac{\partial}{\partial x}, xy^2 \frac{\partial}{\partial y}, \\ 25y^4 \frac{\partial}{\partial z} - 4axz \frac{\partial}{\partial y}, xz \frac{\partial}{\partial y} + 5xy^3 \frac{\partial}{\partial z}, y^4 \frac{\partial}{\partial y}, xy^3 \frac{\partial}{\partial x}, xy^3 \frac{\partial}{\partial y}, xy^4 \frac{\partial}{\partial z} xy^4 \frac{\partial}{\partial x}, xy^4 \frac{\partial}{\partial y},$$

5.1.12. $S_{11} : x^4 + y^2z + xz^2 + ax^3z$. — The Gröbner basis of I w.r.t. the lexicographic ordering $z \succ y \succ x$:

$$x^5, x^3y, 16x^4 - 5ax^2y^2, y^3, 2xz + y^2 + ax^3, yz, 2z^2 + 8x^3 - 3axy^2 - 3a^2x^4$$

Basis monomials M :

$$y^2x^2, y^2x, yx^2, x^3, y^2, yx, x^2, z, y, x, 1$$

A basis of Σ :

$$\left[\frac{1}{zy^3x^3} + 2\frac{1}{z^4yx} - \frac{1}{2}\frac{1}{z^2yx^4} + a(-\frac{1}{z^2y^3x} + \frac{1}{2}\frac{1}{z^3yx^2} - \frac{5}{8}\frac{1}{zyx^5}) \right], \\ \left[\frac{1}{zy^3x^2} - \frac{1}{2}\frac{1}{z^2yx^3} - \frac{1}{2}a\frac{1}{zyx^4} \right], \left[\frac{1}{zy^2x^3} - a\frac{1}{z^2y^2x} \right], \left[\frac{1}{zyx^4} - 4\frac{1}{z^3yx} \right], \\ \left[\frac{1}{zy^3x} - \frac{1}{2}\frac{1}{z^2yx^2} \right], \left[\frac{1}{zy^2x^2} \right], \left[\frac{1}{zyx^3} \right], \left[\frac{1}{z^2yx} \right], \left[\frac{1}{zy^2x} \right], \left[\frac{1}{zyx^2} \right], \left[\frac{1}{zyx} \right]$$

A basis of V_f : (# of generators : 6)

$$16xy \frac{\partial}{\partial y} - 16y^2 \frac{\partial}{\partial z} - 48ax^3 \frac{\partial}{\partial z} + 15a^2xy^2 \frac{\partial}{\partial z}, \\ 64x^2 \frac{\partial}{\partial x} + 32y^2 \frac{\partial}{\partial z} + 144ax^3 \frac{\partial}{\partial z} - 65a^2xy^2 \frac{\partial}{\partial z}, xy \frac{\partial}{\partial x}, \\ y^2 \frac{\partial}{\partial y}, 4y^2 \frac{\partial}{\partial x} + 48x^3 \frac{\partial}{\partial z} - 21axy^2 \frac{\partial}{\partial z}, 2x^3 \frac{\partial}{\partial y} + x^2y \frac{\partial}{\partial z}, x^2y \frac{\partial}{\partial y} - xy^2 \frac{\partial}{\partial z}, \\ 2x^3 \frac{\partial}{\partial x} + xy^2 \frac{\partial}{\partial z}, x^2y \frac{\partial}{\partial x}, xy^2 \frac{\partial}{\partial y}, x^2y^2 \frac{\partial}{\partial x}, x^2y^2 \frac{\partial}{\partial z}, x^2y^2 \frac{\partial}{\partial y}, x^2y^2 \frac{\partial}{\partial x}.$$

5.1.13. $S_{12} : x^2y + y^2z + xz^3 + az^5$. — The Gröbner basis of I w.r.t. the lexicographic ordering $y \succ x \succ z$:

$$z^6, 13xz^3 + 20az^5, x^2z^2, z^4 - x^3, 2yz + x^2, z^3 + 2xy, 3xz^2 + y^2 + 5az^4$$

Basis monomials M :

$$z^5, z^4, zx^2, z^2x, z^3, x^2, zx, z^2, y, x, z, 1$$

A basis of Σ :

$$\left[\frac{1}{z^6yx} + \frac{3}{2}\frac{1}{zy^4x} - \frac{1}{2}\frac{1}{z^3y^2x^2} + \frac{1}{z^2yx^4} + a(-\frac{5}{13}\frac{1}{z^2y^3x} - \frac{20}{13}\frac{1}{z^4yx^2} + \frac{10}{13}\frac{1}{zy^2x^3}) \right], \\ \left[\frac{1}{z^5yx} - \frac{1}{2}\frac{1}{z^2y^2x^2} + \frac{1}{zyx^4} - 5a\frac{1}{zy^3x} \right], \left[\frac{1}{z^2yx^3} - \frac{1}{2}\frac{1}{z^3y^2x} \right], \left[\frac{1}{z^3yx^2} - 3\frac{1}{zy^3x} \right], \\ \left[\frac{1}{z^4yx} - \frac{1}{2}\frac{1}{zy^2x^2} \right], \left[\frac{1}{zyx^3} - \frac{1}{2}\frac{1}{z^2y^2x} \right], \left[\frac{1}{z^2yx^2} \right], \left[\frac{1}{z^3yx} \right],$$

$$\left[\frac{1}{zy^2x} \right], \left[\frac{1}{zyx^2} \right], \left[\frac{1}{z^2yx} \right], \left[\frac{1}{zyx} \right]$$

A basis of V_f : (# of generators : 5)

$$\begin{aligned} & 39z^2 \frac{\partial}{\partial x} + 39xz \frac{\partial}{\partial y} + 26y \frac{\partial}{\partial z} - 260axz \frac{\partial}{\partial z} - 400a^2z^3 \frac{\partial}{\partial z}, \\ & 156xz \frac{\partial}{\partial x} + 195yz \frac{\partial}{\partial y} + 117z^2 \frac{\partial}{\partial z} + 130ayz \frac{\partial}{\partial z} - 100a^2xz^2 \frac{\partial}{\partial z}, \\ & 39yz \frac{\partial}{\partial x} - 39xz \frac{\partial}{\partial z} - 50az^3 \frac{\partial}{\partial z}, 39z^3 \frac{\partial}{\partial y} + 78xz \frac{\partial}{\partial z} + 160az^3 \frac{\partial}{\partial z}, \\ & 13z^3 \frac{\partial}{\partial x} + 13yz \frac{\partial}{\partial z} - 50axz^2 \frac{\partial}{\partial z}, 39xz^2 \frac{\partial}{\partial y} - 13yz \frac{\partial}{\partial z} - 110axz^2 \frac{\partial}{\partial z}, \\ & 3xz^2 \frac{\partial}{\partial x} + z^3 \frac{\partial}{\partial z}, 3yz^2 \frac{\partial}{\partial y} + z^3 \frac{\partial}{\partial z}, z^4 \frac{\partial}{\partial y} + 2xz^2 \frac{\partial}{\partial z}, yz^2 \frac{\partial}{\partial x} - xz^2 \frac{\partial}{\partial z}, \\ & z^4 \frac{\partial}{\partial x}, yz^2 \frac{\partial}{\partial z}, z^4 \frac{\partial}{\partial z}, z^5 \frac{\partial}{\partial y} z^5 \frac{\partial}{\partial x}, z^5 \frac{\partial}{\partial z} \end{aligned}$$

5.1.14. $U_{12} : x^3 + y^3 + z^4 + axyz^2$. — The Gröbner basis of I w.r.t. the lexicographic ordering $y \succ x \succ z$:

$$z^5, xz^3, x^2z, 2z^4 - 3x^3, 3x^2 + ayz^2, 2z^3 + axyz, x^2y, 3y^2 + axz^2$$

Basis monomials M :

$$z^2yx, zyx, z^2x, z^2y, yx, zx, zy, z^2, x, y, z, 1$$

A basis of Σ :

$$\begin{aligned} & \left[\frac{1}{z^3y^2x^2} - \frac{1}{3}a \frac{1}{zyx^4} + a \left(-\frac{1}{2} \frac{1}{z^5yx} - \frac{1}{3} \frac{1}{zy^4x} \right) \right], \left[\frac{1}{z^2y^2x^2} - \frac{1}{2}a \frac{1}{z^4yx} \right], \\ & \left[\frac{1}{z^3y^2x} - \frac{1}{3}a \frac{1}{zyx^3} \right], \left[\frac{1}{z^3yx^2} - \frac{1}{3}a \frac{1}{zy^3x} \right], \left[\frac{1}{zy^2x^2} \right], \left[\frac{1}{z^2yx^2} \right], \\ & \left[\frac{1}{z^2y^2x} \right], \left[\frac{1}{z^3yx} \right], \left[\frac{1}{zyx^2} \right], \left[\frac{1}{zy^2x} \right], \left[\frac{1}{z^2yx} \right], \left[\frac{1}{zyx} \right] \end{aligned}$$

A basis of V_f : (# of generators : 8)

$$\begin{aligned} & 2xy \frac{\partial}{\partial x} - yz \frac{\partial}{\partial z}, 2yz \frac{\partial}{\partial y} + z^2 \frac{\partial}{\partial z}, 2xy \frac{\partial}{\partial y} - xz \frac{\partial}{\partial z}, \\ & 3z^2 \frac{\partial}{\partial z} + axy \frac{\partial}{\partial z}, 2xz \frac{\partial}{\partial x} + z^2 \frac{\partial}{\partial z}, 3xz \frac{\partial}{\partial z} - ayz^2 \frac{\partial}{\partial x}, \\ & 3yz \frac{\partial}{\partial z} - axz^2 \frac{\partial}{\partial y}, xz^2 \frac{\partial}{\partial x}, yz^2 \frac{\partial}{\partial y}, yz^2 \frac{\partial}{\partial z}, xz^2 \frac{\partial}{\partial z}, \\ & xyz \frac{\partial}{\partial x}, xyz \frac{\partial}{\partial y}, xyz \frac{\partial}{\partial z}, xyz^2 \frac{\partial}{\partial x}, xyz^2 \frac{\partial}{\partial y}, xyz^2 \frac{\partial}{\partial z} \end{aligned}$$

References

- [1] V.I. ARNOLD, S.M. GUSEIN-ZADE & A.N. VARCHENKO – *Singularities of Differentiable Maps, I*, Monographs in Mathematics, 82, Birkhäuser, Boston, 1985.
- [2] Y. NAKAMURA & S. TAJIMA – A method for constructiong holomic systems for algebraic local cohomology classes with support on a zero dimensional variety, in *Proceedings of the First International Congress of Mathematical Software*, World Scientific, 2002, p. 158–168.
- [3] _____, Algebraic local cohomology classes attached to quasi-homogeneous isolated singularities, *Publ. Res. Inst. Math. Sci.* **41** (to appear).
- [4] T. OAKU – Algorithms for b-functions, restrictions, and algebraic local cohomology groups of D -modules, *Adv. in Appl. Math.* **19** (1997), p. 61–105.
- [5] K. SAITO – Quasihomogene isolierte Singularitäten von Hyperflächen, *Invent. Math.* **14** (1971), p. 123–142.
- [6] S. TAJIMA & Y. NAKAMURA – Algebraic local cohomology classes attached to unimodal singularities, preprint.

Y NAKAMURA, Department of Mathematics, Faculty of Science and Engineering, Kinki University, 3-4-1, Kowakae, Higashi-Osaka, Osaka, 577-8502, Japan • E-mail : yayoi@math.kindai.ac.jp
 S. TAJIMA, Department of Information Engineering, Faculty of Engineering, Niigata University, 2-8050, Ikarashi, Niigata, 950-2181, Japan • E-mail : tajima@ie.niigata-u.ac.jp