THREE CLASSICAL RESULTS ON REPRESENTATIONS OF A NUMBER

Michael D. Hirschhorn

Introduction

Three classical results concern the number of representations of the positive integer n in the form x^2+3y^2 with $x,y\in\mathbb{Z}$, the form $(x^2+x)/2+3(y^2+y)/2$ with $x,y\in\mathbb{Z}^+$ and the form x^2+xy+y^2 with $x,y\in\mathbb{Z}$.
Indeed, if $s(n)$, $t(n)$ and $u(n)$ respectively denote the three numbers, then

$$s(n)=2\Big(d_{1,3}(n)-d_{2,3}(n)\Big)+4\Big(d_{4,12}(n)-d_{8,12}(n)\Big), \tag{1}$$

$$t(n)=d_{1,3}(2n+1)-d_{2,3}(2n+1) \tag{2}$$

and

$$u(n)=6\Big(d_{1,3}(n)-d_{2,3}(n)\Big). \tag{3}$$

where $d_{r,m}(n)$ is the number of divisors d of n with $d\equiv r\pmod m$.
(1) is equivalent to the q-series identity

$$\sum_{m,n\in\mathbb{Z}}q^{m^2+3n^2}=1+2\sum_{n\geq0}\left(\frac{q^{3n+1}}{1-q^{3n+1}}-\frac{q^{3n+2}}{1-q^{3n+2}}\right)$$
$$+4\sum_{n\geq0}\left(\frac{q^{12n+4}}{1-q^{12n+4}}-\frac{q^{12n+8}}{1-q^{12n+8}}\right) \tag{4}$$

or to

$$\sum_{m,n\in\mathbb{Z}}q^{m^2+3n^2}=1+2\sum_{n\geq0}\left(\frac{q^{3n+1}}{1-(-1)^nq^{3n+1}}-\frac{q^{3n+2}}{1+(-1)^nq^{3n+2}}\right), \tag{5}$$

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

(2) is equivalent to the q-series identity

$$\sum_{m,n\in\mathbb{Z}^+} q^{(m^2+m)/2+3(n^2+n)/2} = \sum_{n\geq 0}\left(\frac{q^{3n}}{1-q^{6n+1}} - \frac{q^{3n+2}}{1-q^{6n+5}}\right) \tag{6}$$

and (3) is equivalent to the q-series identity

$$\sum_{m,n\in\mathbb{Z}} q^{m^2+mn+n^2} = 1 + 6\sum_{n\geq 0}\left(\frac{q^{3n+1}}{1-q^{3n+1}} - \frac{q^{3n+2}}{1-q^{3n+2}}\right). \tag{7}$$

Both (5) and (6) appear in Ramanujan's second Notebook [12, p.239], and Berndt [2, pp.223-224 and p.116] shows how they follow from Ramanujan's ${}_1\psi_1$ summation [12, p.196], [2, p.32]. (7) appears in Berndt and Rankin [5, p.196] and a proof is given by Berndt [3]. The reader is referred also to [6], [7], [9] and [4] for related developments and generalisations, and to [1] and [10] for applications in statistical mechanics.

It seems that Dirichlet (1840) may have known (1), since he gives [8, p.463] the corresponding results for the forms x^2+y^2 and x^2+2y^2, and continues "And so on in similar fashion." ("Et ainsi de suite.")

However, Lorenz (1871) [11, p.420] states both (4) and (1), and in reference to (1) says (my translation) "From this equation one can deduce a theorem which must be considered new in the theory of numbers because it cannot immediately be deduced from known theorems:

If a number N contains prime factors $p_1, p_2, \cdots$ of the form $3m+1$ with exponents $a_1, a_2, \cdots$ and if the prime factors of the form $3m+2$ appear to nothing but even powers, the number of solutions of the equation $m^2+3n^2 = N$ is given by

$$\rho_N = 2(a_1+1)(a_2+1)\ldots$$

if N is odd and by

$$\rho_N = 6(a_1+1)(a_2+1)\ldots$$

if N is even. If, on the contrary, N contains a prime factor of the form $3m+2$ to an odd power, one has $\rho_N = 0$."

Lorenz also [11, p.424] states (7), and a proof is provided by his reviewer/ translator Valentiner [11, p.430].

So perhaps credit rests with Lorenz.

We shall give proofs of (4), (6) and (7) which demonstrate that all three results are intimately related.

2. Proof of the result involving $s(n)$

Let $a(q)$ denote the left hand side of (7). Then

$$a(q) + 2a(q^4) = 3 \sum_{k,l \in \mathbb{Z}} q^{k^2+3l^2}. \tag{8}$$

For,

$$a(q) = \sum_{\substack{m \text{ odd} \\ n \text{ even}}} q^{m^2+mn+n^2} + \sum_{\substack{m \text{ odd} \\ n \text{ odd}}} q^{m^2+mn+n^2} + \sum_{\substack{m \text{ even} \\ n \text{ odd}}} q^{m^2+mn+n^2} + \sum_{\substack{m \text{ even} \\ n \text{ even}}} q^{m^2+mn+n^2}.$$

In the first sum, let $k = m + \frac{n}{2},\ l = \frac{n}{2}$ (and conversely, $m = k - l,\ n = 2l$), in the second sum, $k = \frac{m-n}{2},\ l = \frac{m+n}{2}$ (conversely $m = k+l,\ n = l-k$), in the third sum, $k = \frac{m}{2} + n,\ l = \frac{m}{2}$ (conversely $m = 2l,\ n = k - l$) and in the fourth sum, $k = \frac{m-n}{2},\ l = \frac{m+n}{2}$ (conversely $m = k + l,\ n = l - k$),
and we find

$$a(q) = 3 \sum_{k \not\equiv l \pmod 2} q^{k^2+3l^2} + \sum_{k \equiv l \pmod 2} q^{k^2+3l^2}. \tag{9}$$

Also,

$$a(q^4) = \sum_{m,n \in \mathbb{Z}} q^{4m^2+4mn+4n^2} = \sum_{m,n \text{ even}} q^{m^2+mn+n^2} = \sum_{k \equiv l \pmod 2} q^{k^2+3l^2}, \tag{10}$$

as with the fourth sum above. (8) follows from (9) and (10). (4) follows from (7) and (8).

3. Proof of the result involving $t(n)$

We can write (4)

$$\sum_{k,l \in \mathbb{Z}} q^{k^2+3l^2} = 1 + 2 \sum_{n \geq 0} \left(\frac{q^{6n+1}}{1 - q^{12n+2}} - \frac{q^{6n+5}}{1 - q^{12n+10}} \right) + 6 \sum_{n \geq 0} \left(\frac{q^{12n+4}}{1 - q^{12n+4}} - \frac{q^{12n+8}}{1 - q^{12n+8}} \right).$$

If we extract the even powers of q we obtain

$$\sum_{k\equiv l \pmod 2} q^{k^2+3l^2} = 1 + 6\sum_{n\geq 0}\left(\frac{q^{12n+4}}{1-q^{12n+4}} - \frac{q^{12n+8}}{1-q^{12n+8}}\right). \tag{11}$$

(Note, incidentally, that (7) follows from (10) and (11), and that (11) follows from (7) and (10)!)

From (11) we deduce

(12)

$$\sum_{k,l\in\mathbb{Z}} q^{k^2+3l^2} + 4q\sum_{k,l\in\mathbb{Z}^+} q^{k^2+k+3l^2+3l} = 1 + 6\sum_{n\geq 0}\left(\frac{q^{3n+1}}{1-q^{3n+1}} - \frac{q^{3n+2}}{1-q^{3n+2}}\right).$$

If we subtract (4) from (12) we find

(13)

$$\begin{aligned} 4q\sum_{k,l\in\mathbb{Z}^+} q^{k^2+k+3l^2+3l} &= 4\sum_{n\geq 0}\left(\frac{q^{3n+1}}{1-q^{3n+1}} - \frac{q^{3n+2}}{1-q^{3n+2}}\right) \\ &\quad - 4\sum_{n\geq 0}\left(\frac{q^{12n+4}}{1-q^{12n+4}} - \frac{q^{12n+8}}{1-q^{12n+8}}\right) \\ &= 4\sum_{n\geq 0}\left(\frac{q^{6n+1}}{1-q^{12n+2}} - \frac{q^{6n+5}}{1-q^{12n+10}}\right). \end{aligned}$$

∎

Finally, if we divide (13) by $4q$ and replace q^2 by q we obtain (6).

4. Proof of the result involving $u(n)$

We begin by showing that

$$\sum_{m,n\in\mathbb{Z}} \omega^{m-n} q^{m^2+mn+n^2} = \frac{(q)_\infty^3}{(q^3)_\infty} \tag{14}$$

where $\omega^3 = 1,\ \omega \neq 1$.

Let $\mathbf{CT}_a\left\{\sum_{-\infty}^{\infty} a^n f_n(q)\right\}$ denote $f_0(q)$, the "**C**onstant **T**erm" of the Laurent series in a. Then

$$\begin{aligned} \sum_{m,n\in\mathbb{Z}} \omega^{m-n} q^{m^2+mn+n^2} &= \sum_{m+n+p=0} \omega^{m-n} q^{(m^2+n^2+p^2)/2} \\ &= \mathbf{CT}_a\left\{\sum_{-\infty}^{\infty} a^m \omega^m q^{m^2/2} \sum_{-\infty}^{\infty} a^n \omega^{-n} q^{n^2/2} \sum_{-\infty}^{\infty} a^p q^{p^2/2}\right\} \end{aligned}$$

$$= \mathbf{CT}_a \left\{ \prod_{n\geq 1} (1+a\omega q^{n-\frac{1}{2}})(1+a^{-1}\omega^{-1}q^{n-\frac{1}{2}})(1-q^n). \right.$$
$$\left. . \prod_{n\geq 1} (1+a\omega^{-1}q^{n-\frac{1}{2}})(1+a^{-1}\omega q^{n-\frac{1}{2}})(1-q^n). \right.$$
$$\left. . \prod_{n\geq 1} (1+aq^{n-\frac{1}{2}})(1+a^{-1}q^{n-\frac{1}{2}})(1-q^n) \right\}$$
$$= \mathbf{CT}_a \left\{ \prod_{n\geq 1} (1+a^3q^{3n-\frac{3}{2}})(1+a^{-3}q^{3n-\frac{3}{2}})(1-q^n)^3 \right\}$$
$$= \frac{(q)_\infty^3}{(q^3)_\infty} . \mathbf{CT}_a \left\{ \prod_{n\geq 1} (1+a^3q^{3n-\frac{3}{2}})(1+a^{-3}q^{3n-\frac{3}{2}})(1-q^{3n}) \right\}$$
$$= \frac{(q)_\infty^3}{(q^3)_\infty} . \mathbf{CT}_a \left\{ \sum_{-\infty}^{\infty} a^{3n} q^{3n^2/2} \right\}$$
$$= \frac{(q)_\infty^3}{(q^3)_\infty},$$

as claimed.

Now the left hand side of (14) can be written

$$\sum_{m-n\equiv 0 \pmod 3} q^{m^2+mn+n^2} + \omega \sum_{m-n\equiv 1 \pmod 3} q^{m^2+mn+n^2} + \omega^{-1} \sum_{m-n\equiv -1 \pmod 3} q^{m^2+mn+n^2}.$$

In the first sum, let $k = \dfrac{m-n}{3},\ l = \dfrac{m+2n}{3}$ (conversely $m = 2k+l,\ n = l-k$),
in the second sum, $k = \dfrac{m-n-1}{3},\ l = \dfrac{m+2n-1}{3}$ (conversely $m = 2k+l+1,\ n = l-k$) and
in the third sum, $k = \dfrac{n-m-1}{3},\ l = \dfrac{n+2m-1}{3}$ ($m = l-k,\ n = 2k+l+1$),
and the left hand side of (14) is seen to be

$$\sum_{k,l\in\mathbb{Z}} q^{3k^2+3kl+3l^2} + \omega \sum_{k,l\in\mathbb{Z}} q^{3k^2+3kl+3l^2+3k+3l+1} + \omega^{-1} \sum_{k,l\in\mathbb{Z}} q^{3k^2+3kl+3l^2+3k+3l+1}$$

$$= a(q^3) - qc(q^3),$$

where

$$c(q) = \sum_{m,n\in\mathbb{Z}} q^{m^2+mn+n^2+m+n}.$$

Thus (14) becomes

$$a(q^3) - qc(q^3) = \frac{(q)_\infty^3}{(q^3)_\infty}. \tag{15}$$

Now, it is a celebrated identity of Jacobi that

$$(q)_\infty^3 = \sum_{n\geq 0}(-1)^n(2n+1)q^{(n^2+n)/2}. \tag{16}$$

We split this sum according to the residue modulo 3 of n. For $n \equiv 0 \pmod 3$, we write $3n$ $(n \geq 0)$, for $n \equiv 1 \pmod 3$, we write $3n+1$ $(n \geq 0)$, and for $n \equiv -1 \pmod 3$ we write $-3n-1$ $(n \leq -1)$, and the right hand side of (16) becomes

$$\begin{aligned}\sum_{n\geq 0}(-1)^n(6n+1)q^{(9n^2+3n)/2} &- \sum_{n\geq 0}(-1)^n(6n+3)q^{(9n^2+9n+2)/2}\\ &- \sum_{n\leq -1}(-1)^n(-6n-1)q^{(9n^2+3n)/2}\\ &= \sum_{-\infty}^{\infty}(-1)^n(6n+1)q^{(9n^2+3n)/2} - 3q(q^9)_\infty^3.\end{aligned}$$

So (15) becomes

$$a(q^3) - qc(q^3) = \frac{1}{(q^3)_\infty}\left\{\sum_{-\infty}^{\infty}(-1)^n(6n+1)q^{3(3n^2+n)/2} - 3q(q^9)_\infty^3\right\}. \tag{17}$$

It follows that

$$a(q) = \frac{1}{(q)_\infty}\sum_{-\infty}^{\infty}(-1)^n(6n+1)q^{(3n^2+n)/2} \tag{18}$$

and

$$c(q) = 3\frac{(q^3)_\infty^3}{(q)_\infty}.$$

Now (18) becomes

$$\begin{aligned}
a(q) &= \frac{1}{(q)_\infty}\left[\frac{d}{da}\sum_{-\infty}^{\infty}(-1)^n a^{6n+1}q^{(3n^2+n)/2}\right]_{a=1}\\
&= \frac{1}{(q)_\infty}\left[\frac{d}{da}\left\{a\prod_{n\geq 1}(1-a^6q^{3n-1})(1-a^{-6}q^{3n-2})(1-q^{3n})\right\}\right]_{a=1}\\
&= \frac{1}{(q)_\infty}\left[\prod_{n\geq 1}(1-a^6q^{3n-1})(1-a^{-6}q^{3n-2})(1-q^{3n})\times\right.\\
&\qquad\left.\times\left\{1+6\sum_{n\geq 1}\left(\frac{a^{-6}q^{3n-2}}{1-a^{-6}q^{3n-2}}-\frac{a^6q^{3n-1}}{1-a^6q^{3n-1}}\right)\right\}\right]_{a=1}\\
&= \frac{1}{(q)_\infty}(q)_\infty\left\{1+6\sum_{n\geq 1}\left(\frac{q^{3n-2}}{1-q^{3n-2}}-\frac{q^{3n-1}}{1-q^{3n-1}}\right)\right\}\\
&= 1+6\sum_{n\geq 0}\left(\frac{q^{3n+1}}{1-q^{3n+1}}-\frac{q^{3n+2}}{1-q^{3n+2}}\right)
\end{aligned}$$

which is (7).

References

[1] R. J. Baxter, Some hyperelliptic function identities that occur in the chiral Potts model, J. Phys. A. Math. Gen. 31 (1998), 6806–6818.

[2] B. C. Berndt, Ramanujan's Notebooks, Part III, Springer-Verlag, New York, 1991.

[3] B. C. Berndt, On a certain theta-function in a letter of Ramanujan from Fitzroy House, Ganita 43 (1992), 33–43.

[4] B. C. Berndt, S. Bhargava and F. G. Garvan, Ramanujan's theories of elliptic functions to alternative bases, Trans. Amer. Math. Soc., 347 (1995), 4163–4244.

[5] B. C. Berndt and R. A. Rankin, Ramanujan: Letters and Commentary, History of Mathematics, Vol. 9, Amer. Math. Soc., London Math. Soc., 1995.

[6] J. M. Borwein and P. B. Borwein, A cubic counterpart of Jacobi's identity and the AGM, Trans. Amer. Math. Soc., 323 (1991), 691–701.

[7] J. M. Borwein, P. B. Borwein and F. G. Garvan, Some cubic modular identities of Ramanujan, Trans. Amer. Math. Soc., 343 (1994), 35–47.

[8] G. Lejeune Dirichlet, Mathematische Werke, Chelsea, New York, 1969.

[9] M. D. Hirschhorn, F. G. Garvan and J. M. Borwein, Cubic analogues of the Jacobian theta function $\theta(z,q)$, Can. J. Math., 45 (1993), 673–694.

[10] M. D. Hirschhorn, Proofs of some hyperelliptic function identities, Bull. Austral. Math. Soc., 58 (1998), 465–468.

[11] L. Lorenz, Oeuvres Scientifiques, Revues et annotées par H. Valentiner, Tome second, la Fondation Carlsberg, Librarie Lehmann & Stage, Copenhague, 1904.

[12] S. Ramanujan, Notebooks (2 volumes), Tata Institute of Fundamental Research, Bombay, 1957.