ON THE COMPOSITION OF FORCES

$\mathbf{B}\mathbf{y}$

William Rowan Hamilton

(Proceedings of the Royal Irish Academy, 2 (1844), pp. 166–168.)

Edited by David R. Wilkins 2000

On the Composition of Forces. By Sir WILLIAM R. HAMILTON.

Communicated November 8, 1841.

[Proceedings of the Royal Irish Academy, vol. 2 (1844), pp. 166–168.]

The Chair having been taken, *pro tempore*, by the Rev. J.H. Todd, D.D., V.P., the President communicated the following proof of the known law of Composition of Forces.

Two rectangular forces, x and y, being supposed to be equivalent to a single resultant force p, inclined at an angle v to the force x, it is required to determine the law of the dependence of this angle on the ratio of the two component forces x and y.

Denoting by p' any other single force, intermediate between x and y, and inclined to x at an angle v', which we shall suppose to be greater than v; and denoting by x' and y' the rectangular components of this new force p', in the directions of x and y, we may, by easy decompositions and recompositions, obtain a new pair of rectangular forces, x'' and y'', which are together equivalent to p', and have for components

$$x'' = \frac{x}{p}x' + \frac{y}{p}y';$$

$$y'' = \frac{x}{p}y' - \frac{y}{p}x';$$

the direction of x'' coinciding with that of p', but the direction of y'' being perpendicular thereto. Hence,

 $\frac{y''}{x''} = \frac{xy' - yx'}{xx' + yy'};$

that is,

$$\tan^{-1}\frac{y''}{x''} = \tan^{-1}\frac{y'}{x'} - \tan^{-1}\frac{y}{x};$$

or finally,

$$f(v' - v) = f(v') - f(v),$$
 (A)

at least for values of v, v', and v' - v, which are each greater than 0, and less than $\frac{\pi}{2}$; if f be a function so chosen that the equation

$$\frac{y}{x} = \tan f(v)$$

expresses the sought law of connexion between the ratio $\frac{y}{x}$ and the angle v. The functional equation (A) gives

$$f(mv) = mf(v) = \frac{m}{n}f(nv),$$

1	1	
1		

m and n being any whole numbers; and the case of equal components gives evidently

$$f\left(\frac{\pi}{4}\right) = \frac{\pi}{4};$$

hence

$$f\left(\frac{m}{n}\frac{\pi}{4}\right) = \frac{m}{n}\frac{\pi}{4},$$
$$f(v) = v,$$
(B)

and ultimately,

because it is evident, by the nature of the question, that while v increases from 0 to $\frac{\pi}{2}$, the function f(v) increases therewith, and therefore could not be equal thereto for all values of v commensurate with $\frac{\pi}{4}$, unless it had the same property also for all intermediate incommensurable values. We find, therefore, that for all values of the component forces x and y, the equation

$$\frac{y}{x} = \tan v \tag{C}$$

holds good; that is, the resultant force coincides in direction with the diagonal of the rectangle constructed with lines representing x and y as sides.

The other part of the known law of the composition of forces, namely, that this resultant is represented also *in magnitude* by the same diagonal, may easily be proved by the process of the Mécanique Céleste, which, in the present notation, corresponds to making

$$x' = x, \quad y' = y, \quad x'' = p,$$

and therefore gives

$$p = \frac{x^2 + y^2}{p}, \quad p^2 = x^2 + y^2.$$

But the demonstration above assigned for the law of the *direction* of the resultant, appears to Sir William Hamilton to be new.