

Locally Nilpotent Linear Groups

A. S. DETINKO AND D. L. FLANNERY

Dedicated to Martin L. Newell

ABSTRACT. This article examines aspects of the theory of locally nilpotent linear groups. We also present a new classification result for locally nilpotent linear groups over an arbitrary field \mathbb{F} .

1. WHY LOCALLY NILPOTENT LINEAR GROUPS?

Linear (matrix) groups are a commonly used concrete representation of groups. The first investigations of linear groups were undertaken in the second half of the 19th century, and currently linear group theory is a highly developed branch of group theory. In the past few decades interest in matrix groups has revived and increased, driven partly by the rapid development of computational group theory.

Locally nilpotent groups are a generalization of nilpotent groups. Over the years, many structural and classification results for locally nilpotent linear groups have been obtained. Further progress in the study of these groups is possible using computational techniques.

Group theoretical algorithms take as input a finite generating set for a group. The celebrated ‘Tits alternative’ states that a finitely generated linear group G either is solvable-by-finite (that is, G contains a normal solvable subgroup of finite index), or G contains a nonabelian free subgroup. For linear groups of the latter type, some basic computational problems, such as membership testing and the conjugacy problem, are undecidable in general. Nilpotent linear groups on the other hand are solvable-by-finite and so are more suitable for computation (note that the class of nilpotent-by-finite linear

2000 *Mathematics Subject Classification*. Primary 20F19, Secondary 20H20.

This publication has emanated from research conducted with the financial support of Science Foundation Ireland.

groups—which includes all locally nilpotent linear groups—forms an important subclass of

This point is underlined by Gromov’s result [5], which implies that a finitely generated group has polynomial growth if and only if it is nilpotent-by-finite: hence, as explained in [1], certain algorithmic efficiency problems can be successfully overcome for locally nilpotent linear groups.

Another motivation for further study of locally nilpotent linear groups lies in possible application to abstract group theory concerns. Here an example is recent work [11] on the Carter conjecture for finite groups. Another example is the almost crystallographic groups, which are nilpotent-by-finite and arise naturally as linear groups over \mathbb{Q} (see [4, §5.3]). Note that a finitely generated nilpotent group is polycyclic and so isomorphic to a subgroup of $\mathrm{GL}(n, \mathbb{Z})$ for some n ; hence algorithms for nilpotent subgroups of $\mathrm{GL}(n, \mathbb{Q})$ serve as a key step toward algorithms for abstract finitely generated nilpotent groups.

2. STRUCTURE OF LOCALLY NILPOTENT LINEAR GROUPS

Research into locally nilpotent linear groups relies heavily on knowledge of the structure of such groups. Systematic study of the structure of locally nilpotent linear groups was carried out by D. A. Suprunenko, beginning in the late 1940s [14]. Among other things, Suprunenko classified the maximal locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ when \mathbb{F} is algebraically closed. Various authors extended some of Suprunenko’s results to other fields. In particular, criteria for finiteness of the number of conjugacy classes of maximal locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$, as well as classification of such groups in some partial cases, have been obtained from a detailed description of the structure of locally nilpotent linear groups over an arbitrary field (see e.g. [8]).

In the rest of this section we outline some of the most important structural results for locally nilpotent linear groups. A natural point of focus is the maximals, because each locally nilpotent subgroup of $\mathrm{GL}(n, \mathbb{F})$ is contained in a maximal locally nilpotent subgroup (by way of contrast, note that a nilpotent subgroup of $\mathrm{GL}(n, \mathbb{F})$ may not be contained in a maximal nilpotent subgroup).

We proceed via a standard reduction scheme: reducible \rightarrow completely reducible \rightarrow irreducible \rightarrow absolutely irreducible \rightarrow primitive.

2.1. Reducible Locally Nilpotent Linear Groups. We use standard terminology for linear groups, as in [14, 15]. A reducible subgroup G of $\text{GL}(n, \mathbb{F})$ is conjugate to a group of block upper triangular matrices, where the diagonal blocks form the irreducible parts of G , which are irreducible representations of G over \mathbb{F} in smaller degree. If G is indecomposable locally nilpotent then a stronger statement holds: the irreducible parts of G are pairwise equivalent ([15, p.223, Theorem 2]), so that G is conjugate to a group of block upper triangular matrices

$$\begin{pmatrix} a(g) & a_{12}(g) & \cdots & a_{1k}(g) \\ 0 & a(g) & \cdots & a_{2k}(g) \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a(g) \end{pmatrix}, \quad g \in G$$

where $a(G) = \{a(g) \mid g \in G\} \leq \text{GL}(m, \mathbb{F})$ is irreducible locally nilpotent for some m dividing n , and $a_{ij}(g) \in \text{Mat}(m, \mathbb{F})$. If \mathbb{F} is a perfect field then $a_{ij}(g) = c_{ij}(g)a(g)$ where $c_{ij}(g) \in \mathbb{C}_{\text{Mat}(m, \mathbb{F})}(a(G))$; that is, G is contained in the direct product of a completely reducible group over \mathbb{F} with equivalent irreducible parts, and a unitriangular group over a division algebra. This reduces study of locally nilpotent linear groups to the irreducible case.

Another way to obtain a reduction to the completely reducible case is to use the Jordan decomposition. For each $g \in \text{GL}(n, \mathbb{F})$ there is a unique unipotent matrix $g_u \in \text{GL}(n, \overline{\mathbb{F}})$ and a unique diagonalizable matrix $g_d \in \text{GL}(n, \overline{\mathbb{F}})$ such that $g = g_d g_u = g_u g_d$ (here $\overline{\mathbb{F}}$ is the algebraic closure of \mathbb{F}); see [16, p. 91, 7.2]. If \mathbb{F} is perfect then g_u and g_d are both in $\text{GL}(n, \mathbb{F})$.

Theorem 2.1. ([16, p.97, 7.11] and [15, p.240, Theorem 6]) *Let $G \leq \text{GL}(n, \mathbb{F})$ be locally nilpotent, and define $G_u = \{g \in G \mid g \text{ unipotent}\}$, $G_d = \{g \in G \mid g \text{ diagonalizable}\}$. Then G_u and G_d are normal subgroups of G , and $\langle G_u, G_d \rangle = G_u \times G_d$.*

The theorem implies that if G is a completely reducible locally nilpotent subgroup of $\text{GL}(n, \mathbb{F})$ then every subgroup of G is completely reducible, and in particular every element of G is diagonalizable ([16, p.98, 7.12] and [15, p.239, Theorem 5]).

Corollary 2.2. ([16, p.98, 7.13]) *If $G \leq \text{GL}(n, \mathbb{F})$ is locally nilpotent, and if for each $g \in G$ we have $g_u, g_d \in G$, then $G = G_u \times G_d$.*

The hypothesis $g_u, g_d \in G$ for all $g \in G$ is satisfied if \mathbb{F} is finite. This is a partial case of the following.

Theorem 2.3. ([12, p.136, Proposition 3]) *Let \mathbb{F} be perfect and G be nilpotent. Define $\hat{G}_u = \{g_u \mid g \in G\}$ and $\hat{G}_d = \{g_d \mid g \in G\}$. Then \hat{G}_u, \hat{G}_d are subgroups of $\mathrm{GL}(n, \mathbb{F})$, and $G \leq \hat{G}_u \times \hat{G}_d$.*

The group \hat{G}_d defined in Theorem 2.3 is completely reducible over \mathbb{F} .

2.2. Irreducible locally nilpotent linear groups. Each irreducible maximal locally nilpotent subgroup of $\mathrm{GL}(n, \mathbb{F})$ can be thought of as an absolutely irreducible maximal locally nilpotent subgroup of $\mathrm{GL}(m, \mathbb{E})$ for some m dividing n and field $\mathbb{E} \supseteq \mathbb{F}$ (see [15, p.217, Theorem 4]). This affords a reduction to the absolutely irreducible case, particularly in the classification of irreducible maximal locally nilpotent linear groups. Further investigation is possible in two directions, which are not mutually exclusive: reduction to p -subgroups of $\mathrm{PGL}(n, \mathbb{F})$, and reduction to primitive groups. The former is based on the following theorem.

Theorem 2.4. ([15, pp.220–221, Theorems 8, 9])

- (i) *Let $n = p_1^{a_1} \cdots p_k^{a_k}$ be the prime factorisation of n , where the p_i are pairwise distinct primes. If G is a maximal absolutely irreducible locally nilpotent subgroup of $\mathrm{GL}(n, \mathbb{F})$ then $G = G_1 \otimes \cdots \otimes G_k$ where $G_i \leq \mathrm{GL}(p_i^{a_i}, \mathbb{F})$ is maximal absolutely irreducible locally nilpotent, $1 \leq i \leq k$.*
- (ii) *Let p be a prime. An absolutely irreducible subgroup G of $\mathrm{GL}(p^a, \mathbb{F})$ containing $\mathbb{F}^\times 1_{p^a}$ is a maximal absolutely irreducible locally nilpotent subgroup of $\mathrm{GL}(p^a, \mathbb{F})$ if and only if $G/\mathbb{F}^\times 1_{p^a}$ is a Sylow p -subgroup of $\mathrm{PGL}(p^a, \mathbb{F})$.*

Irreducible locally nilpotent linear groups are center-by-periodic; in fact, the central quotient of each is a direct product of p -groups (see [13, Corollary 3.2.4]).

Except when \mathbb{F} is finite or algebraically closed, the description of Sylow p -subgroups of $\mathrm{PGL}(n, \mathbb{F})$ is quite different to the description of Sylow p -subgroups of $\mathrm{GL}(n, \mathbb{F})$.

Sylow p -subgroups of $\mathrm{PGL}(n, \mathbb{F})$ were considered in [6], mainly for $p > 2$. Classifying the Sylow 2-subgroups of $\mathrm{PGL}(n, \mathbb{F})$ is difficult.

In [6], p -subgroups of $\mathrm{PGL}(n, \mathbb{F})$ are handled using the same techniques as for locally nilpotent linear groups, including the reduction to primitives.

The reduction to primitives is not so straightforward for locally nilpotent linear groups as it is for some other classes of linear groups, such as solvable groups. To appreciate this disparity, note that an irreducible imprimitive solvable subgroup of $\mathrm{GL}(n, \mathbb{F})$ is conjugate to a subgroup of $G \wr T$ where G is a primitive solvable linear group and T is a transitive solvable permutation group ([15, p.129, Theorem 5]); however, the wreath product of a locally nilpotent linear group and a nilpotent permutation group need not even be (locally) nilpotent.

A discussion of techniques for studying nilpotent primitive subgroups G of $\mathrm{GL}(n, \mathbb{F})$ can be found in [8, Section 2]. One technique is to use the series $G \geq H \geq K \geq 1$ where $K = [G, G]$ and $H = C_G(K)$.

A basic result here is as follows.

Theorem 2.5. ([8, Theorem 2]) *Let $G \leq \mathrm{GL}(p^a, \mathbb{F})$ be primitive absolutely irreducible locally nilpotent, $p \neq \mathrm{char} \mathbb{F}$. Then K is an abelian p -group, $\Sigma = \langle K \rangle_{\mathbb{F}}$ is a field, $G/H \cong \mathrm{Gal}(\Sigma/\mathbb{F})$, $[H, H] \leq \mathbb{F}^{\times} 1_{p^a}$, and H is an absolutely irreducible primitive class 2 nilpotent subgroup of $\mathrm{GL}(m, \Sigma)$, $m = p^a / |\Sigma : \mathbb{F}|$.*

The primitive nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ for finite fields \mathbb{F} have been completely classified, in [3].

The paper [8] also summarizes results and methods for classifying maximal locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ over an arbitrary field \mathbb{F} .

Considerable attention has been paid to the problem of determining when the number of $\mathrm{GL}(n, \mathbb{F})$ -conjugacy classes of maximal locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ is finite. Finiteness of that number depends on finiteness of the groups $\mathbb{F}^{\times} / (\mathbb{F}^{\times})^m$, for m dividing n .

Groups over an algebraically closed field have been the most intensively studied.

Theorem 2.6. ([14, Chapter III] and [15, Chapter VII]) *Let \mathbb{F} be algebraically closed.*

- (i) *Irreducible locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ exist if and only if $\mathrm{char} \mathbb{F} = 0$ or $\mathrm{char} \mathbb{F}$ does not divide n , in which cases there exists an irreducible nilpotent subgroup of $\mathrm{GL}(n, \mathbb{F})$ of nilpotency class l , for each and every nilpotency class $l \geq 1$.*

- (ii) *Irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ are monomial and pairwise conjugate.*

The matrix form of the groups in (ii) is given in [14, Chapter III, §7]. Note that (ii) implies that a completely reducible locally nilpotent linear group over an algebraically closed field is monomial.

In summary, locally nilpotent linear groups constitute a well-studied class of groups, for which a lot of structural information and efficient methods of investigation are known. However the theory still has significant gaps. Most results deal only with absolutely irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$. Those results do not readily yield analogous results for locally nilpotent linear groups that are not maximal or are not absolutely irreducible (cf. [3]). Complete classifications of locally nilpotent subgroups of $\mathrm{GL}(n, \mathbb{F})$ are feasible only by placing restrictions on the field \mathbb{F} or the degree n . In the sequel we allow arbitrary fields but restrict the degree.

3. PRIME DEGREE LOCALLY NILPOTENT LINEAR GROUPS

We now give an illustration of how techniques in the theory of locally nilpotent linear groups may be applied to obtain a full classification in that theory. Specifically, in this section we classify the irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(q, \mathbb{F})$, where q is prime and \mathbb{F} is any field. This classification is in the form of a complete list of $\mathrm{GL}(q, \mathbb{F})$ -conjugacy class representatives of the groups, with each listed group defined by a generating set of matrices. Also we provide criteria to decide conjugacy between listed groups. Restricting to prime degree has several advantages: an irreducible subgroup G of $\mathrm{GL}(q, \mathbb{F})$ is either abelian or absolutely irreducible, and is either primitive or monomial. Additionally, if G is absolutely irreducible locally nilpotent then lies in a Sylow q -subgroup of $\mathrm{PGL}(q, \mathbb{F})$, and Sylow q -subgroups of $\mathrm{PGL}(q, \mathbb{F})$ have a simpler description than do Sylow subgroups of $\mathrm{PGL}(n, \mathbb{F})$ for composite degree n .

Certainly, a partial classification of the irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(q, \mathbb{F})$ can be derived from a description of the absolutely irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(q^a, \mathbb{F})$. However, here we propose other methods and give a complete, self-contained result, which can be extended to get a complete classification in prime power degrees q^a . In particular we give an exact description of the Sylow 2-subgroups of $\mathrm{PGL}(2, \mathbb{F})$,

omitted by other authors. This is of special importance because (for example) classification of the Sylow 2-subgroups of $\mathrm{PGL}(n, \mathbb{F})$ in arbitrary degree n depends on classification in degree 2 (cf. the case of 2-subgroups of $\mathrm{GL}(n, \mathbb{F})$ in [7, 10]).

The methods used in this section were originally developed to classify the maximal irreducible periodic subgroups of $\mathrm{PGL}(q, \mathbb{F})$, in [2]. We begin by considering absolutely irreducible groups; abelian groups will be treated at the end.

By [15, Theorem 6, p. 217], $\mathrm{GL}(q, \mathbb{F})$ contains absolutely irreducible locally nilpotent subgroups if and only if \mathbb{F}^\times has an element ξ of order q (of course $\mathrm{char} \mathbb{F} \neq q$). Let $D = \{\mathrm{diag}(\beta\beta_1, \dots, \beta\beta_q) \mid \beta_i \in \mathrm{Syl}_q(\mathbb{F}^\times), \beta \in \mathbb{F}^\times\}$. For $\alpha \in \mathbb{F}^\times$, define

$$I_\alpha = \begin{pmatrix} 0 & 1_{q-1} \\ \alpha & 0 \end{pmatrix} \in \mathrm{GL}(q, \mathbb{F})$$

and write I in place of I_1 . For $H \leq \mathrm{GL}(q, \mathbb{F})$ let $\mathrm{Det}(H) = \{\det(h) \mid h \in H\}$.

Assuming $\xi \in \mathbb{F}^\times$ we define $H_\alpha = \langle D, I_\alpha \rangle$. The subgroup H_α of $\mathrm{GL}(q, \mathbb{F})$ is monomial and absolutely irreducible. Since $H_\alpha/\mathbb{F}^\times 1_q$ is a q -subgroup of $\mathrm{PGL}(q, \mathbb{F})$, H_α is locally nilpotent. If $\mathrm{Syl}_q(\mathbb{F}^\times)$ is finite then H_α is nilpotent with nilpotency class $1 + (q-1) \log_q |\mathrm{Syl}_q(\mathbb{F}^\times)|$.

Denote by π the natural homomorphism from the group of all monomial matrices in $\mathrm{GL}(q, \mathbb{F})$ onto the group $\mathrm{Sym}(q)$ of $q \times q$ permutation matrices. The kernel of π is the group $\mathrm{D}(q, \mathbb{F})$ of all diagonal matrices in $\mathrm{GL}(q, \mathbb{F})$.

Lemma 3.1. (Cf. [2, Lemma 20]) *Let $a, b \in \mathrm{D}(q, \mathbb{F})$. The following statements are equivalent.*

- (i) Ia, Ib are $\mathrm{GL}(q, \mathbb{F})$ -conjugate.
- (ii) Ia, Ib are $\mathrm{D}(q, \mathbb{F})$ -conjugate.
- (iii) $\det(a) = \det(b)$.

Lemma 3.2. *Let H be an irreducible monomial locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$. Then H is conjugate in $\mathrm{GL}(q, \mathbb{F})$ to a subgroup of H_α for some $\alpha \in \mathbb{F}^\times$.*

Proof. If H is abelian then $\pi(H) \leq \mathrm{Sym}(q)$ is a transitive abelian group i.e. a cycle of order q , and $H \cap \mathrm{D}(q, \mathbb{F}) \leq \mathbb{F}^\times 1_q$. On the other hand if H is absolutely irreducible then $H\mathbb{F}^\times/\mathbb{F}^\times 1_q$ is a q -group, so that $H \cap \mathrm{D}(q, \mathbb{F}) \leq D$, and $\pi(H)$ is again a cycle of order q . Hence up to conjugacy $H \leq \langle D, Ia \rangle$ for some $a \in \mathrm{D}(q, \mathbb{F})$. Then H is conjugate to a subgroup of $H_{\det(a)}$ by Lemma 3.1. \square

Denote the set $\text{Det}(D) = \text{Syl}_q(\mathbb{F}^\times)(\mathbb{F}^\times)^q$ by S .

Lemma 3.3. *If $\alpha \in S$ then H_α is $D(q, \mathbb{F})$ -conjugate to H_1 . If $\alpha_1, \alpha_2 \notin S$ then H_{α_1} and H_{α_2} are $\text{GL}(q, \mathbb{F})$ -conjugate if and only if $\text{Det}(H_{\alpha_1}) = \text{Det}(H_{\alpha_2})$ i.e. $\langle \alpha_1 S \rangle$ and $\langle \alpha_2 S \rangle$ are identical subgroups of \mathbb{F}^\times/S of order q .*

Proof. Suppose $\alpha = \beta_1 \beta^q$ for some $\beta_1 \in \text{Syl}_q(\mathbb{F}^\times)$ and $\beta \in \mathbb{F}^\times$. Then $\det(I_\alpha) = \det(Ib)$ where $b = \text{diag}(\beta_1 \beta, \beta, \dots, \beta) \in D$. Therefore H_α is $D(q, \mathbb{F})$ -conjugate to $\langle Ib, D \rangle = H_1$ by Lemma 3.1.

Now suppose $\alpha_1, \alpha_2 \notin S$ and $\alpha_1 \in \langle \alpha_2, S \rangle$. Then $\det(I_{\alpha_1}) = \det(I_{\alpha_2}^r c)$ for some $c \in D$ and $1 \leq r \leq q-1$. Also there exists $x \in \text{Sym}(q)$ such that $x I_{\alpha_2}^r c x^{-1} = Ib$ for some $b \in D(q, \mathbb{F})$. Hence by Lemma 3.1 once more, H_{α_1} and H_{α_2} are conjugate (this time by a monomial matrix). \square

Corollary 3.4. *Define $\mathcal{H} = \{H_\alpha \mid \alpha \in \mathbb{F}^\times \setminus S\}$. The $\text{GL}(q, \mathbb{F})$ -conjugacy classes of the groups in \mathcal{H} are in one-to-one correspondence with the distinct subgroups of \mathbb{F}^\times/S of order q . Consequently the number of such classes is finite if and only if \mathbb{F}^\times/S is finite.*

Remark 3.5. If \mathbb{F} is algebraically closed or finite then \mathcal{H} is empty: a maximal absolutely irreducible monomial locally nilpotent subgroup of $\text{GL}(q, \mathbb{F})$ is conjugate to H_1 .

We turn next to primitive groups.

Lemma 3.6. *Let H be a primitive locally nilpotent subgroup of $\text{GL}(q, \mathbb{F})$. Then H has an irreducible abelian normal subgroup.*

Proof. First we show that H has an abelian noncentral normal subgroup. As H is locally nilpotent, it is solvable, and thus has an abelian normal subgroup A of finite index (see e.g. [15, p.135, Theorem 6]). If $A \leq Z(H)$ then $H/Z(H)$ is finite and thus H is nilpotent. But a nonabelian nilpotent group certainly contains an abelian noncentral normal subgroup.

Any abelian noncentral normal subgroup A of H must be irreducible. For if A were reducible then it would be diagonalizable with inequivalent irreducible parts, which contradicts primitivity of H . \square

Lemma 3.6 implies that a primitive locally nilpotent subgroup of $\text{GL}(q, \mathbb{F})$ is contained in the $\text{GL}(q, \mathbb{F})$ -normalizer of the multiplicative group of a field extension Δ of $\mathbb{F}1_q$ of degree q . Since this

degree is prime, Δ is a cyclic extension of \mathbb{F} , with Galois group of order q . As long as \mathbb{F}^\times has an element ξ of order q , $\Delta = \langle h \rangle_{\mathbb{F}}$ for some $h \in \text{GL}(q, \mathbb{F})$ such that $h^q = \beta 1_q \in \mathbb{F}^\times 1_q$ by [9, p. 289, Theorem 6.2]. As $\beta \in (\mathbb{F}^\times)^q$ implies that h is scalar, we have $\beta = \alpha^r \gamma^q$ for some $\alpha, \gamma \in \mathbb{F}^\times$ and $1 \leq r \leq q-1$. Then $\gamma^{-1}h$ and I_α^r have the same characteristic (minimal) polynomial $X^q - \alpha^r 1_q$, and because $X^q - \alpha^r 1_q$ is \mathbb{F} -irreducible, $\gamma^{-1}h$ and I_α^r are conjugate. Hence Δ is conjugate to $\Delta_\alpha := \langle I_\alpha \rangle_{\mathbb{F}}$, $\alpha \notin (\mathbb{F}^\times)^q$, and $\text{N}_{\text{GL}(q, \mathbb{F})}(\Delta_\alpha^\times) = \langle \Delta_\alpha^\times, d \rangle$ where $d = \text{diag}(1, \xi, \dots, \xi^{q-1})$. Denote by $G(\alpha, b)$ the subgroup $\langle A_\alpha, db \rangle$ of $\langle \Delta_\alpha^\times, d \rangle$, where $A_\alpha \supseteq \mathbb{F}^\times 1_q$, $A_\alpha / \mathbb{F}^\times 1_q$ is the Sylow q -subgroup of $\Delta_\alpha^\times / \mathbb{F}^\times 1_q$, and $b \in \Delta_\alpha^\times$. Since A_α is a noncentral irreducible subgroup, $G(\alpha, b)$ is absolutely irreducible.

Lemma 3.7. *An absolutely irreducible primitive locally nilpotent subgroup H of $\text{GL}(q, \mathbb{F})$ is conjugate to a subgroup of some $G(\alpha, b)$.*

Proof. Up to conjugacy $H = \langle H \cap \Delta_\alpha^\times, db \rangle$ for some $\alpha \in \mathbb{F}^\times \setminus (\mathbb{F}^\times)^q$ and $b \in \Delta_\alpha^\times$. Then $H \cap \Delta_\alpha \leq A_\alpha$ by Theorem 2.4. \square

We use the notation ε_k to stand for an element of multiplicative order 2^k in the algebraic closure of \mathbb{F} . If $k = 2$ then we drop the subscript; that is, ε is a square root of -1 .

Lemma 3.8. *Suppose \mathbb{F} does not have characteristic 2, and $\varepsilon \notin \mathbb{F}$. Let $\mathbb{E} = \mathbb{F}(\varepsilon)$, and let σ be the \mathbb{F} -involution of \mathbb{E} . If $\text{Syl}_2(\mathbb{E}^\times) = \langle \varepsilon_m \rangle$ is cyclic then $\text{Syl}_2(\mathbb{E}^\times / \mathbb{F}^\times)$ is cyclic. Explicitly, one of the following must be true:*

- (i) $\sigma(\varepsilon_m) = -\varepsilon_m^{-1}$, and $\text{Syl}_2(\mathbb{E}^\times / \mathbb{F}^\times) = \langle \varepsilon_m \mathbb{F}^\times \rangle$ of order 2^{m-1} ;
or
- (ii) $\sigma(\varepsilon_m) = \varepsilon_m^{-1}$, and $\text{Syl}_2(\mathbb{E}^\times / \mathbb{F}^\times) = \langle (1 + \varepsilon_m) \mathbb{F}^\times \rangle$ of order 2^m .

Proof. We make some preliminary observations. First, either $\sigma(\varepsilon_m) = \varepsilon_m^{-1}$ or $\sigma(\varepsilon_m) = -\varepsilon_m^{-1}$. Suppose $\sigma(\varepsilon_k) = \varepsilon_k^{-1}$. Then

$$\sigma((1 + \varepsilon_k)^{2^k}) = \sigma(1 + \varepsilon_k)^{2^k} = (1 + \varepsilon_k^{-1})^{2^k} = \left(\frac{1 + \varepsilon_k}{\varepsilon_k} \right)^{2^k} = (1 + \varepsilon_k)^{2^k}.$$

Thus $(1 + \varepsilon_k)^{2^k} \in \mathbb{F}$ and so

$$(1 + \varepsilon_k) \mathbb{F}^\times \in \text{Syl}_2(\mathbb{E}^\times / \mathbb{F}^\times) \quad (1)$$

if $k \geq 2$. Also, if $k > 2$ then $\varepsilon_k^{-1}(\varepsilon_{k-1} + 1) = \varepsilon_k^{-1}(\varepsilon_k^2 + 1) = \varepsilon_k + \varepsilon_k^{-1} \in \mathbb{F}^\times$ and so

$$1 + \varepsilon_{k-1} \in \langle \varepsilon_k \rangle \mathbb{F}^\times. \quad (2)$$

Now let $x\mathbb{F}^\times$ be a nontrivial element of $\text{Syl}_2(\mathbb{E}^\times/\mathbb{F}^\times)$ of order 2^l , meaning that $x^{2^l} \in \mathbb{F}^\times \setminus (\mathbb{F}^\times)^2$.

Suppose $l = 1$. Write $x = a + \varepsilon b$, $a, b \in \mathbb{F}$. Then $2ab\varepsilon \in \mathbb{F}$ implies $a = 0$ i.e.

$$x \in \langle \varepsilon\mathbb{F}^\times \rangle \leq \langle \varepsilon_m\mathbb{F}^\times \rangle. \quad (3)$$

Suppose $l \geq 2$. We have $\sigma(x) = yx$ for some $y \in \text{Syl}_2(\mathbb{E}^\times)$, $y^{2^l} = 1$. Then $y \neq -1$, because $y = -1$ implies $|x\mathbb{F}^\times| = 2 < 2^l$. Further, $x = \sigma^2(x) = \sigma(yx) = \sigma(y)yx$ and so

$$\sigma(y) = y^{-1}. \quad (4)$$

As $\text{tr}(x) = (1 + y)x \in \mathbb{F}$,

$$x \in (1 + y)^{-1}\mathbb{F}^\times = (1 + \sigma(y))\mathbb{F}^\times = (1 + y^{-1})\mathbb{F}^\times. \quad (5)$$

At last we are ready to complete the proof that either (i) or (ii) must be true. Let $\sigma(\varepsilon_m) = -\varepsilon_m^{-1}$; then $m > 2$. By (3), we may take $l \geq 2$, in which event $4 < |y| \leq 2^{m-1}$ by (4). Then $(1 + y^{-1})\mathbb{F}^\times \subseteq \langle \varepsilon_m \rangle\mathbb{F}^\times$ by (2), and by (5), (i) is proved.

If $\sigma(\varepsilon_m) = \varepsilon_m^{-1}$ then by (1), $\langle (1 + \varepsilon_m)\mathbb{F}^\times \rangle \subseteq \text{Syl}_2(\mathbb{E}^\times/\mathbb{F}^\times)$. Also $\varepsilon_m\mathbb{F}^\times \in \langle (1 + \varepsilon_m)\mathbb{F}^\times \rangle^2$, since σ fixes $\varepsilon_m^{-1}(1 + \varepsilon_m)^2$. Part (ii) now follows from (2), (3), and (5). \square

Corollary 3.9. *If $\text{Syl}_2(\mathbb{E}^\times)$ is quasicyclic then $\text{Syl}_2(\mathbb{E}^\times/\mathbb{F}^\times)$ is also quasicyclic, and $\text{Syl}_2(\mathbb{E}^\times/\mathbb{F}^\times) = \{\langle \varepsilon_k\mathbb{F}^\times \rangle \mid \varepsilon_k \in \text{Syl}_2(\mathbb{E}^\times)\}$.*

Proof. For each $\varepsilon_k \in \text{Syl}_2(\mathbb{E}^\times)$, $\sigma(\varepsilon_k) = \varepsilon_k^{-1}$ i.e. $\sigma(\varepsilon_k) = -\varepsilon_k^{-1}$ is impossible. The corollary is then a consequence of Lemma 3.8 (ii) and (2) in the proof of the lemma. \square

Lemma 3.10. *Let $|\mathbb{E} : \mathbb{F}| = q$ and $\mathbb{E} = \mathbb{F}(a)$, where $a^{q^l} \in \mathbb{F}$. Suppose $\xi \in \mathbb{F}^\times$, and $\mathbb{E} \neq \mathbb{F}(\xi)$ if $q = 2$. Then $\text{Syl}_q(\mathbb{E}^\times/\mathbb{F}^\times) = \langle a\mathbb{F}^\times \rangle$.*

Proof. If $q > 2$, or $q = 2$ and $\varepsilon \in \mathbb{F}$, then the lemma follows from [8, Lemma 2].

Let $q = 2$ and $x\mathbb{F}^\times \in \mathbb{E}^\times/\mathbb{F}^\times$ be of order 2^m , $m \geq 1$, so that $x^{2^m} = \alpha$ for some $\alpha \in \mathbb{F}^\times \setminus (\mathbb{F}^\times)^2$. If $\alpha = -4\gamma^4$, $\gamma \in \mathbb{F}^\times$, then $x^{2^{m-1}}/2\gamma^2$ is a square root of -1 , contradicting $\mathbb{E} \neq \mathbb{F}(\varepsilon)$.

Suppose $\alpha \notin -4(\mathbb{F}^\times)^4$. The polynomial $X^4 - \alpha$ is \mathbb{F} -irreducible, so that if $m \geq 2$ then $|\mathbb{E} : \mathbb{F}| \geq 4$. Therefore $m = 1$, $a = \sqrt{\beta}$ for some $\beta \in \mathbb{F}^\times$ and $x = \sqrt{\alpha}$. For some $x_1, x_2 \in \mathbb{F}$ we have $\sqrt{\alpha} = x_1 + x_2\sqrt{\beta}$. Then $\alpha = x_1^2 + \beta x_2^2 + 2x_1x_2\sqrt{\beta}$ implies $x_1 = 0$ or $x_2 = 0$; as the latter is impossible we get $x \in \langle a\mathbb{F}^\times 1_2 \rangle$ as required. \square

Lemma 3.11. *If $q > 2$ or $\alpha \notin -(\mathbb{F}^\times)^2$ then A_α is the monomial group $\langle I_\alpha, \mathbb{F}^\times 1_q \rangle$; otherwise, A_α is primitive.*

Proof. In Lemma 3.10 put $\mathbb{E} = \Delta_\alpha$ and $a = I_\alpha$. Then $A_\alpha = \langle I_\alpha, \mathbb{F}^\times 1_q \rangle$ unless $q = 2$ and $\Delta_\alpha \cong \mathbb{F}(\varepsilon)$ i.e. $\alpha \in -(\mathbb{F}^\times)^2$. If $A_{-\gamma^2}$ were monomial then $A_{-\gamma^2}^2$ would be in $\mathbb{F}^\times 1_q$; however $A_{-\gamma^2}/\mathbb{F}^\times 1_q$ contains the element $(1_2 + \gamma^{-1}I_{-\gamma^2})\mathbb{F}^\times 1_q$ of order 4. \square

We refer to the set of hypotheses $q = 2$ and $\alpha \in -(\mathbb{F}^\times)^2$ as case (*). Lemma 3.8 and Corollary 3.9 give an explicit description of the A_α in case (*). Actually, a group $G(-\gamma^2, b)$ in this case is conjugate to some $G(-1, b')$, since $I_{-\gamma^2}$ is $D(q, \mathbb{F})$ -conjugate to γI_{-1} by Lemma 3.1. In all but case (*), $|G(\alpha, b)/\mathbb{F}^\times 1_q| = q^2$ (because $(db)^q = \det(b)1_q$ and $[I_\alpha, d]$ is scalar) and so $G(\alpha, b)$ is class 2 nilpotent. The group $G(-1, b)$ is locally nilpotent, and it is nilpotent only if $\text{Syl}_2(\Delta_{-1}^\times)$ is cyclic; then $G(-1, b)/\mathbb{F}^\times 1_q$ is a dihedral 2-group, and $G(-1, b)$ has nilpotency class $\log_2 |\text{Syl}_2(\Delta_{-1}^\times)/\mathbb{F}^\times 1_2|$.

Lemma 3.12. *In case (*), $G(\alpha, b)$ is primitive. In all other cases, $G(\alpha, b)$ is primitive if and only if $\det(b) \notin \langle (-1)^{q-1}\alpha, (\mathbb{F}^\times)^q \rangle = \text{Det}(A_\alpha)$.*

Proof. By Lemma 3.11, assume we are in a case other than (*). Then [8, Lemma 1] yields that $G(\alpha, b)$ is primitive if and only if all elements of $G(\alpha, b)$ of order q are scalar. Suppose $\det(b) \notin \text{Det}(A_\alpha)$ and let $h \in G(\alpha, b)$, $|h| = q$. If $h \notin A_\alpha$ i.e. $h = dbb_1$, $b_1 \in A_\alpha$, then $h^q = \det(bb_1)1_q$ implies that $\det(b) \in \text{Det}(A_\alpha)$. Thus $h \in A_\alpha$, and h is scalar by Lemma 3.11. Conversely, if $\det(b) \in \text{Det}(A_\alpha)$ then for some $x \in A_\alpha$, dbx is a nonscalar element of $G(\alpha, b)$ of order q . \square

Remark 3.13. Except in case (*), if \mathbb{F} is finite then $G(\alpha, b)$ is monomial.

Lemma 3.14. *For $i = 1, 2$, let $g_i = db_i$ where $b_i \in \Delta_\alpha$. The following statements are equivalent.*

- (i) g_1 and g_2 are $\text{GL}(q, \mathbb{F})$ -conjugate.
- (ii) g_1 and g_2 are Δ_α -conjugate.
- (iii) $\det(b_1) = \det(b_2)$.

Proof. See [2, Lemma 21]. \square

Corollary 3.15. *Apart from case (*), primitive groups $G(\alpha, b_1)$, $G(\alpha, b_2)$ are conjugate if and only if $\text{Det}(G(\alpha, b_1)) = \text{Det}(G(\alpha, b_2))$ and $\det(b_1) = \det(b_2c)$ for some $c \in A_\alpha$.*

Proof. Suppose $tG(\alpha, b_1)t^{-1} = G(\alpha, b_2)$. Since t normalizes A_α as a consequence of Lemma 3.12, so $t \in \langle d, \Delta_\alpha^\times \rangle$. Then it can be checked that $tdb_1t^{-1} \in db_2A_\alpha$. The other direction is clear by Lemma 3.14. \square

Denote by \mathcal{G} the set of all groups $G(\alpha, b)$ that are primitive, subject to the proviso that the only groups in case (*) included in \mathcal{G} are the $G(-1, b)$.

Remark 3.16. Note that \mathcal{G} is empty if \mathbb{F} is algebraically closed, for then $G(\alpha, b)$ is not defined. When \mathbb{F} is finite, \mathcal{G} is nonempty if and only if $q = 2$ and $|\mathbb{F}| \equiv 3 \pmod{4}$.

Lemma 3.17. *If nonempty, the subset $\tilde{\mathcal{G}}$ of \mathcal{G} consisting of the groups not in case (*) splits up into finitely many $\mathrm{GL}(q, \mathbb{F})$ -conjugacy classes if and only if $\mathbb{F}^\times / (\mathbb{F}^\times)^q$ is finite.*

Proof. If $\tilde{\mathcal{G}}$ has only finitely many non-conjugate elements then $\mathbb{F}^\times / (\mathbb{F}^\times)^q$ is finitely generated and so finite. Conversely, if $\mathbb{F}^\times / (\mathbb{F}^\times)^q$ is finite then there are only finitely many subsets of \mathbb{F}^\times that are possibilities for $\mathrm{Det}(G(\alpha, b))$; hence the number of $\mathrm{GL}(q, \mathbb{F})$ -conjugacy classes in $\tilde{\mathcal{G}}$ is finite by Corollary 3.15. \square

The next theorem is our main classification result.

Theorem 3.18. *Suppose $\xi \in \mathbb{F}^\times$. A subgroup G of $\mathrm{GL}(q, \mathbb{F})$ is an absolutely irreducible maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ if and only if G is conjugate to a group in $\mathcal{N} = \{H_1\} \cup \mathcal{H} \cup \mathcal{G}$, with the following exceptions when $q = 2$ and $\varepsilon \notin \mathbb{F}$:*

- (i) H_1 is a proper subgroup of $G(-1, 1) \in \mathcal{G}$;
- (ii) if $\alpha \notin -(\mathbb{F}^\times)^2$ and either $\det(b) \in -(\mathbb{F}^\times)^2$ or $\det(b) \in \alpha(\mathbb{F}^\times)^2$, then $G(\alpha, b)$ is conjugate to a proper subgroup of $G(-1, c)$ where $\det(c) = \alpha$.

Proof. We have observed previously that all of the groups in \mathcal{N} are absolutely irreducible locally nilpotent. By Lemmas 3.2, 3.3, and 3.7, and remarks after Lemma 3.11, an absolutely irreducible locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ is conjugate to a subgroup of a group in \mathcal{N} . Therefore it remains to show that the H_α and $G(\alpha, b) \in \mathcal{G}$ are really maximal locally nilpotent, apart from the stated exceptions.

Let G be a maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ containing H_α . If G is monomial then $tGt^{-1} = H_\beta$ for some H_β and $t \in \mathrm{GL}(q, \mathbb{F})$. If $tDt^{-1} \neq D$ then $tDt^{-1} \cap D$ is scalar of index q in D ,

so $|H_\beta/\mathbb{F}^\times 1_q| = q^2$; but $H_\beta/\mathbb{F}^\times 1_q$ has cardinality at least q^{q+1} . Thus $tDt^{-1} = D$, and then $q = |H_\beta : D| \geq |tH_\alpha t^{-1} : D| = |H_\alpha : D| = q$. Therefore $tH_\alpha t^{-1} = H_\beta$ i.e. $H_\alpha = G$. Now suppose G is primitive, hence conjugate to some $G(\alpha_1, b)$. In a case other than $(*)$ we have $|G/\mathbb{F}^\times 1_q| = q^2$, which is less than the cardinality of $H_\alpha/\mathbb{F}^\times 1_q$. Hence $q = 2$, $\varepsilon \notin \mathbb{F}^\times$, G is conjugate to $G(-1, b)$, and $H_\alpha = \langle d, I_\alpha, \mathbb{F}^\times 1_2 \rangle$. Either I_α or $L_\alpha = dI_\alpha$ is conjugate to nonscalar $h \in A_{-1}$ such that $h^2 \in \mathbb{F}^\times 1_2$. Now h has the form γI_{-1} , $\gamma \in \mathbb{F}^\times$, and by comparing determinants we get $\alpha = \pm\gamma^2$. Thus if $H_\alpha \in \mathcal{H}$ then H_α is maximal. However $H_1 = \langle d, I_{-1}, \mathbb{F}^\times 1_2 \rangle$ is a proper subgroup of $G(-1, 1)$.

Let G be a maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ containing $G(\alpha, b) \in \mathcal{G}$. For some $t \in \mathrm{GL}(q, \mathbb{F})$,

$$tGt^{-1} = G(\alpha_1, b_1) \in \mathcal{G}.$$

Apart from when $q = 2$, $\varepsilon \notin \mathbb{F}^\times$, and $\alpha_1 = -1$, we have

$$|G(\alpha, b)/\mathbb{F}^\times 1_q| = |G(\alpha_1, b_1)/\mathbb{F}^\times 1_q| = |G/\mathbb{F}^\times 1_q| = q^2,$$

and thus $G(\alpha, b) = G$. Suppose now that $q = 2$, $\varepsilon \notin \mathbb{F}$, and $\alpha_1 = -1$. If $\alpha \in -(\mathbb{F}^\times)^2$ then $G(\alpha, b)$ is conjugate to some $G(-1, b_2)$, so that $G(\alpha, b) = G$. Therefore if $G(\alpha, b)$ is not maximal then $\alpha \notin -(\mathbb{F}^\times)^2$. For the rest of the proof $\alpha \notin -(\mathbb{F}^\times)^2$, which means that $G(\alpha, b) = \langle I_\alpha, db, \mathbb{F}^\times 1_2 \rangle$. One of the following must occur: $tI_\alpha t^{-1} \notin A_{-1}$ and $tdbt^{-1} \in A_{-1}$, or $tI_\alpha t^{-1} \notin A_{-1}$ and $tdbt^{-1} \notin A_{-1}$.

In the first case, $\det(b) \in -(\mathbb{F}^\times)^2$. In the second case, the condition that $[I_\alpha, db]$ is scalar forces $tdbt^{-1} \in I_{-1}tI_\alpha t^{-1}\mathbb{F}^\times$, so $\det(b) \in \alpha(\mathbb{F}^\times)^2$. Suppose $\det(b) \in -(\mathbb{F}^\times)^2$. Then $\det(db) = \det(\gamma I_{-1})$ for some $\gamma \in \mathbb{F}^\times$. Since also $\mathrm{tr}(db) = 0 = \mathrm{tr}(\gamma I_{-1})$, there exists $s \in \mathrm{GL}(2, \mathbb{F})$ such that $sdb s^{-1} = \gamma I_{-1}$.

Since $I_\alpha db I_\alpha^{-1} = -db$, we have that $sI_\alpha s^{-1}$ normalizes Δ_{-1}^\times . Hence $sI_\alpha s^{-1} = dc$, where $c \in \Delta_{-1}^\times$ and $\det(c) = \alpha$. It follows that $G(\alpha, b)$ is conjugate to a proper subgroup of $G(-1, c)$. The reasoning that leads to this same conclusion in the case $\det(b) \in \alpha(\mathbb{F}^\times)^2$ is entirely similar. \square

With the next lemma we complete our classification of the irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(q, \mathbb{F})$.

Lemma 3.19. *Suppose $\mathrm{GL}(q, \mathbb{F})$ has irreducible abelian subgroups, and let H be an irreducible maximal abelian subgroup of $\mathrm{GL}(q, \mathbb{F})$.*

- (i) If $\xi \notin \mathbb{F}^\times$ then H is a maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$, and moreover any maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ is abelian.
- (ii) Let $\xi \in \mathbb{F}^\times$. Then H is a maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ unless $q = 2$ and $\varepsilon \notin \mathbb{F}$. If $q = 2$ and $\varepsilon \notin \mathbb{F}$ then H is a maximal locally nilpotent subgroup of $\mathrm{GL}(q, \mathbb{F})$ if and only if $H/\mathbb{F}^\times 1_2$ is not a 2-group.

Proof. so that if H is not maximal locally nilpotent then $q = 2$, $\varepsilon \notin \mathbb{F}$, and $H = A_\alpha \leq G(\alpha, \beta) \in \mathcal{G}$, by Lemma 3.11. \square

One situation in which all irreducible maximal locally nilpotent subgroups of $\mathrm{GL}(q, \mathbb{F})$ are nonabelian is when $q = 2$ and $|\mathbb{F}|$ is a Mersenne prime.

Acknowledgment. This article is part of research carried out in the Department of Mathematics at NUI, Galway over a number of years. We are grateful to Professor Martin Newell for his ongoing generous support and encouragement of our work during this time.

REFERENCES

- [1] R. Beals, *Algorithms for matrix groups and the Tits alternative*, J. Comput. System Sci. **58** (1999), 260–279.
- [2] A. S. Detinko and D. L. Flannery, *Periodic subgroups of projective linear groups in positive characteristic* (submitted).
- [3] A. S. Detinko and D. L. Flannery, *Classification of nilpotent primitive linear groups over finite fields*, Glasgow Math. J. **46** (2004), 585–594.
- [4] B. Eick, *Computational group theory*, Jahresbericht der DMV 107, Heft 3 (2005), 155–170.
- [5] M. L. Gromov, *Groups of polynomial growth and expanding maps*, Inst. Hautes Études Sci. Publ. Math. **53** (1981), 53–73.
- [6] V. S. Konyukh, *Sylow p -subgroups of a projective linear group*, Vestsi Akad. Navuk BSSR Ser. Fiz.-Mat. Navuk (1985), no. 6, 23–29, 124–125.
- [7] V. S. Konyukh, *On linear p -groups*, Vestsi Akad. Navuk BSSR Ser. Fiz.-Mat. Navuk (1987), no. 1, 3–8, 124.
- [8] V. S. Konyukh, *Irreducible locally nilpotent linear groups*, Fundam. Prikl. Mat. **4** (1998), 1345–1364.
- [9] Serge Lang, *Algebra*, Addison-Wesley Publishing Co., Inc., Reading, Mass., 1993.
- [10] C. R. Leedham-Green and W. Plesken, *Some remarks on Sylow subgroups of general linear groups*, Math. Z. **191** (1986), 529–535.
- [11] A. Previtali, M. C. Tamburini, and E. P. Vdovin, *The Carter subgroups of some classical groups*, Bull. London Math. Soc. **36** (2004), 145–155.
- [12] D. Segal, *Polycyclic groups*, Cambridge University Press, Cambridge, 1983.

- [13] M. Shirvani and B. A. F. Wehrfritz, *Skew linear groups*, London Mathematical Society Lecture Note Series, vol. 118, Cambridge University Press, Cambridge, 1986.
- [14] D. A. Suprunenko, *Soluble and nilpotent linear groups*, Transl. Math. Monogr., vol. 9, American Mathematical Society, Providence, RI, 1963.
- [15] D. A. Suprunenko, *Matrix groups*, Transl. Math. Monogr., vol. 45, American Mathematical Society, Providence, RI, 1976.
- [16] B. A. F. Wehrfritz, *Infinite linear groups*, Springer-Verlag, Berlin, Heidelberg, New York, 1973.

A. S. Detinko and D. L. Flannery,
Department of Mathematics,
National University of Ireland,
Galway
alla.detinko@nuigalway.ie
dane.flannery@nuigalway.ie