

**MA3484—Methods of Mathematical  
Economics  
School of Mathematics, Trinity College  
Hilary Term 2017  
Lecture 19 (March 9, 2017)**

David R. Wilkins

## 5. Duality and Convexity

### 5.1. General Linear Programming Problems

Linear programming is concerned with problems seeking to maximize or minimize a linear functional of several real variables subject to a finite collection of constraints, where each constraint either fixes the values of some linear function of the variables or else requires those values to be bounded, above or below, by some fixed quantity.

The objective of such a problem involving  $n$  real variables  $x_1, x_2, \dots, x_n$  is to maximize or minimize an *objective function* of those variables that is of the form

$$c_1x_1 + c_2x_2 + \cdots + c_nx_n,$$

subject to appropriate constraints. The coefficients  $c_1, c_2, \dots, c_n$  that determine the objective function are then fixed real numbers.

Now such an optimization problem may be presented as a minimization problem, because simply changing the signs of all the coefficients  $c_1, c_2, \dots, c_n$  converts any maximization problem into a minimization problem. We therefore suppose, without loss of generality, that the objective of the linear programming problem is to find a feasible solution satisfying appropriate constraints which minimizes the value of the objective function amongst all such feasible solutions to the problem.

Some of the constraints may simply require specific variables to be non-negative or non-positive. Now a constraint that requires a particular variable  $x_j$  to be non-positive can be reformulated as one requiring a variable to be non-negative by substituting  $x_j$  for  $-x_j$  in the statement of the problem. Thus, without loss of generality, we may suppose that all constraints that simply specify the sign of a variable  $x_j$  will require that variable to be non-negative. Then all such constraints can be specified by specifying a subset  $J^+$  of  $\{1, 2, \dots, n\}$ : the constraints then require that  $x_j \geq 0$  for all  $j \in J^+$ .

## 5. Duality and Convexity (continued)

There may be further constraints in addition to those that simply specify whether one of the variables is required to be non-positive or non-negative. Suppose that there are  $m$  such additional constraints, and let them be numbered between 1 and  $m$ . Then, for each integer  $i$  between 1 and  $m$ , there exist real numbers  $A_{i,1}, A_{i,2}, \dots, A_{i,n}$  and  $b_i$  that allow the  $i$ th constraint to be expressed either as an *inequality constraint* of the form

$$A_{i,1}x_1 + A_{i,2}x_2 + \dots + A_{i,n}x_n \geq b_i$$

or else as an *equality constraint* of the form

$$A_{i,1}x_1 + A_{i,2}x_2 + \dots + A_{i,n}x_n = b_i.$$

It follows from the previous discussion that the statement of a general linear programming problem can be transformed, by changing the signs of some of the variables and constants in the statement of the problem, so as to ensure that the statement of the problem conforms to the following restrictions:—

- the objective function is to be minimized;
- some of the variables may be required to be non-negative;
- other constraints are either inequality constraints placing a lower bound on the value of some linear function of the variables or else equality constraints fixing the value of some linear function of the variables.

## 5. Duality and Convexity (continued)

Let us describe the statement of a linear programming problem as being in *general primal form* if it conforms to the restrictions just described.

A linear programming problem is expressed in general primal form if the specification of the problem conforms to the following restrictions:—

- the objective of the problem is to find an optimal solution minimizing the objective function amongst all feasible solutions to the problem;
- any variables whose sign is prescribed are required to be non-negative, not non-positive;
- all inequality constraints are expressed by prescribing a lower bound on the value on some linear function of the variables.

## 5. Duality and Convexity (continued)

A linear programming problem in general primal form can be specified by specifying the following data: an  $m \times n$  matrix  $A$  with real coefficients, an  $m$ -dimensional vector  $\mathbf{b}$  with real components; an  $n$ -dimensional vector  $\mathbf{c}$  with real components; a subset  $I^+$  of  $\{1, 2, \dots, m\}$ ; and a subset  $J^+$  of  $\{1, 2, \dots, n\}$ . The linear programming programming problem specified by this data is the following:—

*seek  $\mathbf{x} \in \mathbb{R}^n$  that minimizes the objective function  $\mathbf{c}^T \mathbf{x}$   
subject to the following constraints:—*

- $A\mathbf{x} \geq \mathbf{b}$ ;
- $(A\mathbf{x})_i = (\mathbf{b})_i$  unless  $i \in I^+$ ;
- $(\mathbf{x})_j \geq 0$  for all  $j \in J^+$ .



We refer to the  $m \times n$  matrix  $A$ , the  $m$ -dimensional vector  $\mathbf{b}$  and the  $n$ -dimensional vector  $\mathbf{c}$  employed in specifying a linear programming problem in general primal form as the *constraint matrix*, *target vector* and *cost vector* respectively for the linear programming problem. Let us refer to the subset  $I^+$  of  $\{1, 2, \dots, m\}$  specifying those constraints that are inequality constraints as the *inequality constraint specifier* for the problem, and let us refer to the subset  $J^+$  of  $\{1, 2, \dots, n\}$  that specifies those variables that are required to be non-negative for a feasible solution as the *variable sign specifier* for the problem.

We denote by  $\text{Primal}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  the linear programming problem whose specification in general primal form is determined by a constraint matrix  $A$ , target vector  $\mathbf{b}$ , cost vector  $\mathbf{c}$ , inequality constraint specifier  $I^+$  and variable sign specifier  $J^+$ .

A linear programming problem formulated in general primal form can be reformulated as a problem in Dantzig standard form, thus enabling the use of the Simplex Method to find solutions to the problem.

Indeed consider a linear programming problem

Primal( $A, \mathbf{b}, \mathbf{c}, I^+, J^+$ ) where the constraint matrix  $A$  is an  $m \times n$  matrix with real coefficients, the target vector  $\mathbf{b}$  and the cost vector  $\mathbf{c}$  are vectors of dimension  $m$  and  $n$  respectively with real coefficients. Then the inequality constraint specifier  $I^+$  is a subset of  $\{1, 2, \dots, m\}$  and the variable sign specifier  $J^+$  is a subset of  $\{1, 2, \dots, n\}$ . The problem is already in Dantzig standard form if and only if  $I^+ = \emptyset$  and  $J^+ = \{1, 2, \dots, n\}$ .

## 5. Duality and Convexity (continued)

If the problem is not in Dantzig standard form, then each variable  $x_j$  for  $j \notin J^+$  can be replaced by a pair of variables  $x_j^+$  and  $x_j^-$  satisfying the constraints  $x_j^+ \geq 0$  and  $x_j^- \geq 0$ : the difference  $x_j^+ - x_j^-$  of these new variables is substituted for  $x_j$  in the objective function and the constraints. Also a *slack variable*  $z_i$  can be introduced for each  $i \in I^+$ , where  $z_i$  is required to satisfy the sign constraint  $z_i \geq 0$ , and the inequality constraint

$$A_{i,1}x_1 + A_{i,2}x_2 + \dots + A_{i,n}x_n \geq b_i$$

is then replaced by the corresponding equality constraint

$$A_{i,1}x_1 + A_{i,2}x_2 + \dots + A_{i,n}x_n - z_i = b_i.$$

## 5. Duality and Convexity (continued)

The linear programming problem  $\text{Primal}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  in general primal form can therefore be reformulated as a linear programming problem in Dantzig standard form as follows:—

*determine values of  $x_j$  for all  $j \in J^+$ ,  $x_j^+$  and  $x_j^-$  for all  $j \in J^0$ , where  $J^0 = \{1, 2, \dots, n\} \setminus J^+$ , and  $z_i$  for all  $i \in I^+$  so as to minimize the objective function*

$$\sum_{j \in J^+} c_j x_j + \sum_{j \in J^0} c_j x_j^+ - \sum_{j \in J^0} c_j x_j^-$$

*subject to the following constraints:—*

## 5. Duality and Convexity (continued)

- (i)  $\sum_{j \in J^+} A_{i,j} x_j + \sum_{j \in J^0} A_{i,j} x_j^+ - \sum_{j \in J^0} A_{i,j} x_j^- = b_i$  for each  $i \in \{1, 2, \dots, n\} \setminus I^+$ ;
- (ii)  $\sum_{j \in J^+} A_{i,j} x_j + \sum_{j \in J^0} A_{i,j} x_j^+ - \sum_{j \in J^0} A_{i,j} x_j^- - z_i = b_i$  for each  $i \in I^+$ ;
- (iii)  $x_j \geq 0$  for all  $j \in J^+$ ;
- (iv)  $x_j^+ \geq 0$  and  $x_j^- \geq 0$  for all  $j \in J^0$ ;
- (v)  $z_i \geq 0$  for all  $i \in I^+$ .

Once the problem has been reformulated in Dantzig standard form, techniques based on the Simplex Method can be employed in the search for solutions to the problem.

### 5.2. Duals of Linear Programming Problems

Every linear programming problem  $\text{Primal}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  in general primal form determines a corresponding linear programming problem, which we shall denote by  $\text{Dual}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$ , in *general dual form*. The second linear programming problem is referred to as the *dual* of the first, and the first linear programming problem is referred to as the *primal* of its dual.

We shall give the definition of the dual problem associated with a given linear programming problem, and investigate some important relationships between the primal linear programming problem and its dual.

## 5. Duality and Convexity (continued)

Let  $\text{Primal}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  be a linear programming problem in general primal form specified in terms of an  $m \times n$  constraint matrix  $A$ ,  $m$ -dimensional target vector  $\mathbf{b}$ ,  $n$ -dimensional cost vector  $\mathbf{c}$ , inequality constraint specifier  $I^+$  and variable sign specifier  $J^+$ . The corresponding dual problem  $\text{Dual}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  may be specified in *general dual form* as follows:

*seek  $\mathbf{p} \in \mathbb{R}^m$  that maximizes the objective function  $\mathbf{p}^T \mathbf{b}$  subject to the following constraints:—*

- $\mathbf{p}^T A \leq \mathbf{c}^T$ ;
- $(\mathbf{p})_i \geq 0$  for all  $i \in I^+$ ;
- $(\mathbf{p}^T A)_j = (\mathbf{c})_j$  unless  $j \in J^+$ .



**Lemma 5.1**

*Let  $\text{Primal}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  be a linear programming problem expressed in general primal form with constraint matrix  $A$  with  $m$  rows and  $n$  columns, target vector  $\mathbf{b}$ , cost vector  $\mathbf{c}$ , inequality constraint specifier  $I^+$  and variable sign specifier  $J^+$ . Then the feasible and optimal solutions of the corresponding dual linear programming problem  $\text{Dual}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  are those of the problem  $\text{Primal}(-A^T, -\mathbf{c}, -\mathbf{b}, J^+, I^+)$ .*

**Proof**

An  $m$ -dimensional vector  $\mathbf{p}$  satisfies the constraints of the dual linear programming problem  $\text{Dual}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  if and only if  $\mathbf{p}^T A \leq \mathbf{c}^T$ ,  $(\mathbf{p})_i \geq 0$  for all  $i \in I^+$  and  $(\mathbf{p}^T A)_j = (\mathbf{c})_j$  unless  $j \in J^+$ . On taking the transposes of the relevant matrix equations and inequalities, we see that these conditions are satisfied if and only if  $-A^T \mathbf{p} \geq -\mathbf{c}$ ,  $(\mathbf{p})_i \geq 0$  for all  $i \in I^+$  and  $(-A^T \mathbf{p})_j = (-\mathbf{c})_j$  unless  $j \in J^+$ . But these are the requirements that the vector  $\mathbf{p}$  must satisfy in order to be a feasible solution of the linear programming problem  $\text{Primal}(-A^T, -\mathbf{c}, -\mathbf{b}, J^+, I^+)$ . Moreover  $\mathbf{p}$  is an optimal solution of  $\text{Dual}(A, \mathbf{b}, \mathbf{c}, I^+, J^+)$  if and only if it maximizes the value of  $\mathbf{p}^T \mathbf{b}$ , and this is the case if and only if it minimizes the value of  $-\mathbf{b}^T \mathbf{p}$ . The result follows. ■

## 5. Duality and Convexity (continued)

A linear programming problem in Dantzig standard form is specified by specifying integers  $m$  and  $n$  a constraint matrix  $A$  which is an  $m \times n$  matrix with real coefficients, a target vector  $\mathbf{b}$  belonging to the real vector space  $\mathbb{R}^m$  and a cost vector  $\mathbf{c}$  belonging to the real vector space  $\mathbb{R}^n$ . The objective of the problem is to find a feasible solution to the problem that minimizes the quantity  $\mathbf{c}^T \mathbf{x}$  amongst all  $n$ -dimensional vectors  $\mathbf{x}$  for which  $A\mathbf{x} = \mathbf{b}$  and  $\mathbf{x} \geq \mathbf{0}$ .

The objective of the dual problem is then to find some feasible solution to the problem that maximizes the quantity  $\mathbf{p}^T \mathbf{b}$  amongst all  $m$ -dimensional vectors  $\mathbf{p}$  for which  $\mathbf{p}^T A \leq \mathbf{c}$ .