

**MA3484—Methods of Mathematical
Economics
(Mathematical Programming)
School of Mathematics, Trinity College
Hilary Term 2019
Section 2: Finite-Dimensional Vector
Spaces**

David R. Wilkins

2. Finite-Dimensional Vector Spaces

2.1. Real Vector Spaces

Definition

A *real vector space* consists of a set V on which there is defined an operation of vector addition, yielding an element $\mathbf{v} + \mathbf{w}$ of V for each pair \mathbf{v}, \mathbf{w} of elements of V , and an operation of multiplication-by-scalars that yields an element $\lambda\mathbf{v}$ of V for each $\mathbf{v} \in V$ and for each real number λ . The operation of vector addition is required to be commutative and associative. There must exist a zero element $\mathbf{0}_V$ of V that satisfies $\mathbf{v} + \mathbf{0}_V = \mathbf{v}$ for all $\mathbf{v} \in V$, and, for each $\mathbf{v} \in V$ there must exist an element $-\mathbf{v}$ of V for which $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}_V$. The following identities must also be satisfied for all $\mathbf{v}, \mathbf{w} \in V$ and for all real numbers λ and μ :

$$(\lambda + \mu)\mathbf{v} = \lambda\mathbf{v} + \mu\mathbf{v}, \quad \lambda(\mathbf{v} + \mathbf{w}) = \lambda\mathbf{v} + \lambda\mathbf{w},$$

$$\lambda(\mu\mathbf{v}) = (\lambda\mu)\mathbf{v}, \quad 1\mathbf{v} = \mathbf{v}.$$

2. Finite-Dimensional Vector Spaces (continued)

Let n be a positive integer. The set \mathbb{R}^n consisting of all n -tuples of real numbers is then a real vector space, with addition and multiplication-by-scalars defined such that

$$(x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$$

and

$$\lambda(x_1, x_2, \dots, x_n) = (\lambda x_1, \lambda x_2, \dots, \lambda x_n)$$

for all $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ and for all real numbers λ .

The set $M_{m,n}(\mathbb{R})$ of all $m \times n$ matrices is a real vector space with respect to the usual operations of matrix addition and multiplication of matrices by real numbers.

2.2. Linear Dependence and Bases

Elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$ of a real vector space V are said to be *linearly dependent* if there exist real numbers $\lambda_1, \lambda_2, \dots, \lambda_m$, not all zero, such that

$$\lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_m \mathbf{u}_m = \mathbf{0}_V.$$

If elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$ of real vector space V are not linearly dependent, then they are said to be *linearly independent*.

Elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ of a real vector space V are said to *span* V if, given any element \mathbf{v} of V , there exist real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that $\mathbf{v} = \lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_n \mathbf{u}_n$.

A vector space is said to be *finite-dimensional* if there exists a finite subset of V whose members span V .

2. Finite-Dimensional Vector Spaces (continued)

Elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ of a finite-dimensional real vector space V are said to constitute a *basis* of V if they are linearly independent and span V .

Lemma 2.1

Elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ of a real vector space V constitute a basis of V if and only if, given any element \mathbf{v} of V , there exist uniquely-determined real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that

$$\mathbf{v} = \lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_n \mathbf{u}_n.$$

Proof

Suppose that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ is a basis of V . Let \mathbf{v} be an element V . The requirement that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ span V ensures that there exist real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that

$$\mathbf{v} = \lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_n \mathbf{u}_n.$$

If $\mu_1, \mu_2, \dots, \mu_n$ are real numbers for which

$$\mathbf{v} = \mu_1 \mathbf{u}_1 + \mu_2 \mathbf{u}_2 + \cdots + \mu_n \mathbf{u}_n,$$

then

$$(\mu_1 - \lambda_1) \mathbf{u}_1 + (\mu_2 - \lambda_2) \mathbf{u}_2 + \cdots + (\mu_n - \lambda_n) \mathbf{u}_n = \mathbf{0}_V.$$

It then follows from the linear independence of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ that $\mu_i - \lambda_i = 0$ for $i = 1, 2, \dots, n$, and thus $\mu_i = \lambda_i$ for $i = 1, 2, \dots, n$. This proves that the coefficients $\lambda_1, \lambda_2, \dots, \lambda_n$ are uniquely-determined.

2. Finite-Dimensional Vector Spaces (continued)

Conversely suppose that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ is a list of elements of V with the property that, given any element \mathbf{v} of V , there exist uniquely-determined real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that

$$\mathbf{v} = \lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_n \mathbf{u}_n.$$

Then $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ span V . Moreover we can apply this criterion when $\mathbf{v} = \mathbf{0}$. The uniqueness of the coefficients $\lambda_1, \lambda_2, \dots, \lambda_n$ then ensures that if

$$\lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_n \mathbf{u}_n = \mathbf{0}_V$$

then $\lambda_i = 0$ for $i = 1, 2, \dots, n$. Thus $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ are linearly independent. This proves that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ is a basis of V , as required. ■

Proposition 2.2

Let V be a finite-dimensional real vector space, let

$$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$$

be elements of V that span V , and let K be a subset of $\{1, 2, \dots, n\}$. Suppose either that $K = \emptyset$ or else that those elements \mathbf{u}_i for which $i \in K$ are linearly independent. Then there exists a basis of V whose members belong to the list $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ which includes all the vectors \mathbf{u}_i for which $i \in K$.

Proof

We prove the result by induction on the number of elements in the list $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ of vectors that span V . The result is clearly true when $n = 1$. Thus suppose, as the induction hypothesis, that the result is true for all lists of elements of V that span V and that have fewer than n members.

2. Finite-Dimensional Vector Spaces (continued)

If the elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ are linearly independent, then they constitute the required basis. If not, then there exist real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$, not all zero, such that

$$\lambda_1 \mathbf{u}_1 + \lambda_2 \mathbf{u}_2 + \cdots + \lambda_n \mathbf{u}_n = \mathbf{0}_V.$$

Now there cannot exist real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$, not all zero, such that both $\sum_{i=1}^n \lambda_i \mathbf{u}_i = \mathbf{0}_V$ and also $\lambda_i = 0$ whenever $i \neq K$.

Indeed, in the case where $K = \emptyset$, this conclusion follows from the requirement that the real numbers λ_i cannot all be zero, and, in the case where $K \neq \emptyset$, the conclusion follows from the linear independence of those \mathbf{u}_i for which $i \in K$. Therefore there must exist some integer i satisfying $1 \leq i \leq n$ for which $\lambda_i \neq 0$ and $i \notin K$.

2. Finite-Dimensional Vector Spaces (continued)

Without loss of generality, we may suppose that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ are ordered so that $n \notin K$ and $\lambda_n \neq 0$. Then

$$\mathbf{u}_n = - \sum_{i=1}^{n-1} \frac{\lambda_i}{\lambda_n} \mathbf{u}_i.$$

Let \mathbf{v} be an element of V . Then there exist real numbers $\mu_1, \mu_2, \dots, \mu_n$ such that $\mathbf{v} = \sum_{i=1}^n \mu_i \mathbf{u}_i$, because $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ span V . But then

$$\mathbf{v} = \sum_{i=1}^{n-1} \left(\mu_i - \frac{\mu_n \lambda_i}{\lambda_n} \right) \mathbf{u}_i.$$

We conclude that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{n-1}$ span the vector space V . The induction hypothesis then ensures that there exists a basis of V consisting of members of this list that includes the linearly independent elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$, as required. ■

Corollary 2.3

Let V be a finite-dimensional real vector space, and let

$$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$$

be elements of V that span the vector space V . Then there exists a basis of V whose elements are members of the list $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$.

Proof

This result is a restatement of Proposition 2.2 in the special case where the set K in the statement of that proposition is the empty set. ■

2.3. Dual Spaces

Definition

Let V be a real vector space. A *linear functional* $\varphi: V \rightarrow \mathbb{R}$ on V is a linear transformation from the vector space V to the field \mathbb{R} of real numbers.

Given linear functionals $\varphi: V \rightarrow \mathbb{R}$ and $\psi: V \rightarrow \mathbb{R}$ on a real vector space V , and given any real number λ , we define $\varphi + \psi$ and $\lambda\varphi$ to be the linear functionals on V defined such that

$(\varphi + \psi)(\mathbf{v}) = \varphi(\mathbf{v}) + \psi(\mathbf{v})$ and $(\lambda\varphi)(\mathbf{v}) = \lambda\varphi(\mathbf{v})$ for all $\mathbf{v} \in V$.

The set V^* of linear functionals on a real vector space V is itself a real vector space with respect to the algebraic operations of addition and multiplication-by-scalars defined above.

2. Finite-Dimensional Vector Spaces (continued)

Definition

Let V be a real vector space. The *dual space* V^* of V is the vector space whose elements are the linear functionals on the vector space V .

Now suppose that the real vector space V is finite-dimensional. Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ be a basis of V , where $n = \dim V$. Given any $\mathbf{v} \in V$ there exist uniquely-determined real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that $\mathbf{v} = \sum_{j=1}^n \lambda_j \mathbf{u}_j$. It follows that there are well-defined functions $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ from V to the field \mathbb{R} defined such that

$$\varepsilon_i \left(\sum_{j=1}^n \lambda_j \mathbf{u}_j \right) = \lambda_i$$

for $i = 1, 2, \dots, n$ and for all real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$. These functions are linear transformations, and are thus linear functionals on V .

Lemma 2.4

Let V be a finite-dimensional real vector space, let

$$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$$

be a basis of V , and let $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ be the linear functionals on V defined such that

$$\varepsilon_i \left(\sum_{j=1}^n \lambda_j \mathbf{u}_j \right) = \lambda_i$$

for $i = 1, 2, \dots, n$ and for all real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$. Then $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ constitute a basis of the dual space V^ of V .*

Moreover $\varphi = \sum_{i=1}^n \varphi(\mathbf{u}_i) \varepsilon_i$ for all $\varphi \in V^$.*

2. Finite-Dimensional Vector Spaces (continued)

Proof

Let $\mu_1, \mu_2, \dots, \mu_n$ be real numbers with the property that

$\sum_{i=1}^n \mu_i \varepsilon_i = \mathbf{0}_{V^*}$. Then

$$0 = \left(\sum_{i=1}^n \mu_i \varepsilon_i \right) (\mathbf{u}_j) = \sum_{i=1}^n \mu_i \varepsilon_i(\mathbf{u}_j) = \mu_j$$

for $j = 1, 2, \dots, n$. Thus the linear functionals $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ on V are linearly independent elements of the dual space V^* .

2. Finite-Dimensional Vector Spaces (continued)

Now let $\varphi: V \rightarrow \mathbb{R}$ be a linear functional on V , and let $\mu_i = \varphi(\mathbf{u}_i)$ for $i = 1, 2, \dots, n$. Now

$$\varepsilon_i(\mathbf{u}_j) = \begin{cases} 1 & \text{if } i = j; \\ 0 & \text{if } i \neq j. \end{cases}$$

It follows that

$$\begin{aligned} \left(\sum_{i=1}^n \mu_i \varepsilon_i \right) \left(\sum_{j=1}^n \lambda_j \mathbf{u}_j \right) &= \sum_{i=1}^n \sum_{j=1}^n \mu_i \lambda_j \varepsilon_i(\mathbf{u}_j) = \sum_{j=1}^n \mu_j \lambda_j \\ &= \sum_{j=1}^n \lambda_j \varphi(\mathbf{u}_j) = \varphi \left(\sum_{j=1}^n \lambda_j \mathbf{u}_j \right) \end{aligned}$$

for all real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$.

It follows that

$$\varphi = \sum_{i=1}^n \mu_i \varepsilon_i = \sum_{i=1}^n \varphi(\mathbf{u}_i) \varepsilon_i.$$

We conclude from this that every linear functional on V can be expressed as a linear combination of $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$. Thus these linear functionals span V^* . We have previously shown that they are linearly independent. It follows that they constitute a basis of V^* . Moreover we have verified that $\varphi = \sum_{i=1}^n \varphi(\mathbf{u}_i) \varepsilon_i$ for all $\varphi \in V^*$, as required. ■

Definition

Let V be a finite-dimensional real vector space, let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ be a basis of V . The corresponding *dual basis* of the dual space V^* of V consists of the linear functionals $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ on V , where

$$\varepsilon_i \left(\sum_{j=1}^n \lambda_j \mathbf{u}_j \right) = \lambda_i$$

for $i = 1, 2, \dots, n$ and for all real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$.

Corollary 2.5

Let V be a finite-dimensional real vector space, and let V^ be the dual space of V . Then $\dim V^* = \dim V$.*

Proof

We have shown that any basis of V gives rise to a dual basis of V^* , where the dual basis of V has the same number of elements as the basis of V to which it corresponds. The result follows immediately from the fact that the dimension of a finite-dimensional real vector space is the number of elements in any basis of that vector space. ■

2. Finite-Dimensional Vector Spaces (continued)

Let V be a real-vector space, and let V^* be the dual space of V . Then V^* is itself a real vector space, and therefore has a dual space V^{**} . Now each element \mathbf{v} of V determines a corresponding linear functional $E_{\mathbf{v}}: V^* \rightarrow \mathbb{R}$ on V^* , where $E_{\mathbf{v}}(\varphi) = \varphi(\mathbf{v})$ for all $\varphi \in V^*$. It follows that there exists a function $\iota: V \rightarrow V^{**}$ defined so that $\iota(\mathbf{v}) = E_{\mathbf{v}}$ for all $\mathbf{v} \in V$. Then $\iota(\mathbf{v})(\varphi) = \varphi(\mathbf{v})$ for all $\mathbf{v} \in V$ and $\varphi \in V^*$.

Now

$$\iota(\mathbf{v} + \mathbf{w})(\varphi) = \varphi(\mathbf{v} + \mathbf{w}) = \varphi(\mathbf{v}) + \varphi(\mathbf{w}) = (\iota(\mathbf{v}) + \iota(\mathbf{w}))(\varphi)$$

and

$$\iota(\lambda \mathbf{v})(\varphi) = \varphi(\lambda \mathbf{v}) = \lambda \varphi(\mathbf{v}) = (\lambda \iota(\mathbf{v}))(\varphi)$$

for all $\mathbf{v}, \mathbf{w} \in V$ and $\varphi \in V^*$ and for all real numbers λ . It follows that $\iota(\mathbf{v} + \mathbf{w}) = \iota(\mathbf{v}) + \iota(\mathbf{w})$ and $\iota(\lambda \mathbf{v}) = \lambda \iota(\mathbf{v})$ for all $\mathbf{v}, \mathbf{w} \in V$ and for all real numbers λ . Thus $\iota: V \rightarrow V^{**}$ is a linear transformation.

Proposition 2.6

*Let V be a finite-dimensional real vector space, and let $\iota: V \rightarrow V^{**}$ be the linear transformation defined such that $\iota(\mathbf{v})(\varphi) = \varphi(\mathbf{v})$ for all $\mathbf{v} \in V$ and $\varphi \in V^*$. Then $\iota: V \rightarrow V^{**}$ is an isomorphism of real vector spaces.*

Proof

Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ be a basis of V , let $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ be the dual basis of V^* , where

$$\varepsilon_i(\mathbf{u}_j) = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j, \end{cases}$$

and let $\mathbf{v} \in V$. Then there exist real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that $\mathbf{v} = \sum_{i=1}^n \lambda_i \mathbf{u}_i$.

2. Finite-Dimensional Vector Spaces (continued)

Suppose that $\iota(\mathbf{v}) = \mathbf{0}_{V^{**}}$. Then $\varphi(\mathbf{v}) = E_{\mathbf{v}}(\varphi) = 0$ for all $\varphi \in V^*$. In particular $\lambda_i = \varepsilon_i(\mathbf{v}) = 0$ for $i = 1, 2, \dots, n$, and therefore $\mathbf{v} = \mathbf{0}_V$. We conclude that $\iota: V \rightarrow V^{**}$ is injective.

Now let $F: V^* \rightarrow \mathbb{R}$ be a linear functional on V^* , let $\lambda_i = F(\varepsilon_i)$ for $i = 1, 2, \dots, n$, let $\mathbf{v} = \sum_{i=1}^n \lambda_i \mathbf{u}_i$, and let $\varphi \in V^*$. Then

$\varphi = \sum_{i=1}^n \varphi(\mathbf{u}_i) \varepsilon_i$ (see Lemma 2.4), and therefore

$$\begin{aligned}\iota(\mathbf{v})(\varphi) &= \varphi(\mathbf{v}) = \sum_{i=1}^n \lambda_i \varphi(\mathbf{u}_i) = \sum_{i=1}^n F(\varepsilon_i) \varphi(\mathbf{u}_i) \\ &= F\left(\sum_{i=1}^n \varphi(\mathbf{u}_i) \varepsilon_i\right) = F(\varphi).\end{aligned}$$

2. Finite-Dimensional Vector Spaces (continued)

Thus $\iota(\mathbf{v}) = F$. We conclude that the linear transformation $\iota: V \rightarrow V^{**}$ is surjective. We have previously shown that this linear transformation is injective. There $\iota: V \rightarrow V^{**}$ is an isomorphism between the real vector spaces V and V^{**} as required. ■

The following corollary is an immediate consequence of Proposition 2.6.

Corollary 2.7

Let V be a finite-dimensional real vector space, and let V^ be the dual space of V . Then, given any linear functional $F: V^* \rightarrow \mathbb{R}$, there exists some $\mathbf{v} \in V$ such that $F(\varphi) = \varphi(\mathbf{v})$ for all $\varphi \in V^*$.*