1S1 Tutorial Sheet 3^1

17 November 2006

Useful facts:

• The derivative

$$f'(x) = \frac{df}{dx} = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$
(1)

• Higher derivatives:

$$f''(x) = \frac{d^2 f}{dx^2} = \frac{d}{dx} \left(\frac{df}{dx}\right)$$

$$f'''(x) = \frac{d^3 f}{dx^3} = \frac{d}{dx} \left(\frac{d^2 f}{dx^2}\right)$$
(2)

and so on, a bracket number is used after f''', for example f'''' is written $f^{(4)}$.

- Constant: If f = c a constant f' = 0.
- **x** to the n: If $f = x^n$ then $f' = nx^{n-1}$.
- Linearity:

$$\frac{d}{dx}(cf) = c\frac{df}{dx}$$
$$\frac{d}{dx}(f+g) = \frac{df}{dx} + \frac{dg}{dx}$$
(3)

• Product rule:

$$\frac{d}{dx}fg = f\frac{df}{dx} + g\frac{dg}{dx} \tag{4}$$

• Quotient rule:

$$\frac{d}{dx}\frac{f}{g} = \frac{g\frac{df}{dx} - f\frac{dg}{dx}}{g^2} \tag{5}$$

• Trigonometric functions

$$\frac{d}{dx}\cos x = -\sin x$$

$$\frac{d}{dx}\sin x = \cos x \tag{6}$$

¹Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/1S1

Questions

The numbers in brackets give the numbers of marks available for the question.

1. (2) Work out f', f'' and keep going differentiating until you get zero.

$$f(x) = x^7 + 6x^4 + 4x - 1 \tag{7}$$

2. (2) Differentiate

$$f = \frac{1}{x^2} \tag{8}$$

and

$$f = x + \frac{1}{x} \tag{9}$$

3. (2) Use the product and quotient rules to differentiate

$$f = \sqrt{x(x^2 + 1)} \tag{10}$$

and

$$f = \frac{\sqrt{x}}{x^2 + 1} \tag{11}$$

4. (2) The tan is defined as

$$\tan x = \frac{\sin x}{\cos x} \tag{12}$$

and the secant as

$$\sec x = \frac{1}{\cos x} \tag{13}$$

Show using the quotient rule that

$$\frac{d}{dx}\tan x = \sec^2 x \tag{14}$$

Extra Questions

The questions are extra; you don't need to do them in the tutorial class.

1. Differentiate

$$\frac{1}{1+x}, \quad \frac{x^2 - x - 6}{x - 3}, \quad \frac{3x^2 - 4}{x^2 - x + 16}, \quad \frac{x^2 - 4}{\sqrt{x}}, \quad \sin x \cos x, \quad x \sin x \tag{15}$$

- 2. $f(x) = \sin^2 x + \cos^2 x$, we know by the Pythagorous theorem that f(x) = 1, but check that f'(x) = 0 from the original formula with sines and cosines. If f'(x) = 0 then f(x) is a constant, check the value of the constant by working out f(0). Is the answer consistent with the Pythagorous theorem?
- 3. Differentiate $\sec x$ using the quotient rule.