1S1 Tutorial Sheet 4^1

1 December 2006

Useful facts:

- Constant: If f = c a constant f' = 0.
- **x to the n**: If $f = x^n$ then $f' = nx^{n-1}$.
- Trigonometric functions

$$\frac{d}{dx}\cos x = -\sin x$$

$$\frac{d}{dx}\sin x = \cos x \tag{1}$$

• Product rule:

$$\frac{d}{dx}fg = f\frac{df}{dx} + g\frac{dg}{dx} \tag{2}$$

• Quotient rule:

$$\frac{d}{dx}\frac{f}{g} = \frac{g\frac{df}{dx} - f\frac{dg}{dx}}{g^2} \tag{3}$$

• Chain rule: If f(u(x)) then

$$\frac{df}{dx} = \frac{df}{du}\frac{du}{dx} \tag{4}$$

• Stationary points: If f'(x) > 0 a function is increasing at x, if it is negative it is decreasing, if f'(x) = 0 then x is a critical point. If f''(x) > 0 at a critical point it is a minimum, if f''(x) < 0 at a critical point it is a maximum, if f''(x) = 0 it is undecided, it could be a point of inflection, a point joining a concave up f''(x) > 0 interval from a concave down, f''(x) < 0, interval.

Questions

The numbers in brackets give the numbers of marks available for the question.

1. (4) Work out f' for

$$f(x) = x^2 \sin 3x, \qquad \sin x^2, \qquad f(x) = \frac{\cos x}{\sin^2 x}, \qquad f(x) = x \cos \sqrt{x}$$
 (5)

2. (2) Find and classify the stationary points

$$f = 2x^4 - 4x^2 + 6 \tag{6}$$

- 3. (1) Show $f(x) = \tan x$ has a point of inflection at x = 0.
- 4. (1) Classify the stationary points of $3x^4 4x^4$

Extra Questions

The questions are extra; you don't need to do them in the tutorial class.

1. Differentiate

$$\frac{1}{\sqrt{1+x}}, \quad \sin\cos x, \quad \sin\frac{1}{1+x^2}, \quad \sqrt{\sin x}, \quad \tan 2x, \quad \sin\frac{1}{x}$$
 (7)

2. Classify the stationary points for

$$f(x) = x^3 + 3x^2 - 9x + 1,$$
 $f(x) = x^{2/3},$ $f(x) = x^{1/3}(x+4),$ $2x^3 - 6x + 7$ (8)

3. Find and classify the stationary points of $x^3 - 3x^2 + 3x - 1$. You may have to look at the sign of f'(x) to do this.

¹Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/1S1