

A note on conservative fields¹

3 November 2005

Conservative fields

A smooth vector field \mathbf{F} is *conservative* iff there exists a smooth scalar field ϕ such that

$$\mathbf{F} = \text{grad } \phi \quad (1)$$

ϕ is often called a *potential* for \mathbf{F} .

Since $\text{curl grad } \phi = 0$ for any scalar field ϕ , $\text{curl } \mathbf{F} = 0$ is a necessary condition for \mathbf{F} to be conservative. It isn't sufficient, however, this is something we will return to, but, for now, we notice that it makes it easy to spot fields that aren't conservative, for example, if $\mathbf{F} = (-y, x, 0)$ then

$$\nabla \times \mathbf{F} = \begin{pmatrix} 0 \\ 0 \\ 2 \end{pmatrix} \quad (2)$$

On the other hand, it is easy to see that $\mathbf{F} = \mathbf{r}$ is conservative because

$$\nabla \frac{r^2}{2} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (3)$$

and therefore $\phi = r^2/2$ is a potential for \mathbf{F} . Of course, $\text{curl } \mathbf{r} = 0$.

Path independent fields

A field \mathbf{F} is *path independent* if, for any two points p_1 and p_2 the line integral along any path between those points has the same value.

In other words, a field is path independent if the line integral doesn't depend on the path it is taken along. Obviously, the integral around a closed path should be zero for a path independent field and, in fact we can state a lemma:


Lemma: A smooth vector field \mathbf{F} is path independent iff

$$\oint_c \mathbf{F} \cdot d\mathbf{l} = 0 \quad (4)$$

for any closed curve c .²

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/231>

²The integral sign with a circle on it is a standard notation for an integral around a closed path.



Proof: For a path independent \mathbf{F} choose any two points on a closed curve c and label the two curves from p_1 to p_2 as c_1 and c_2 so that $c = c_1 - c_2$. Now, by path independence,

$$\int_{c_1} \mathbf{F} \cdot d\mathbf{l} = \int_{c_2} \mathbf{F} \cdot d\mathbf{l} \quad (5)$$

and so

$$\oint_{c_1 - c_2} \mathbf{F} \cdot d\mathbf{l} = \oint_c \mathbf{F} \cdot d\mathbf{l} = 0 \quad (6)$$

Conversely, two difference paths between two points p_1 and p_2 can be subtracted from each other to give a closed path $c = c_1 - c_2$ so

$$0 = \oint_c \mathbf{F} \cdot d\mathbf{l} = \oint_{c_1 - c_2} \mathbf{F} \cdot d\mathbf{l} \quad (7)$$

implies

$$\int_{c_1} \mathbf{F} \cdot d\mathbf{l} = \int_{c_2} \mathbf{F} \cdot d\mathbf{l} \quad (8)$$

□

Conservative fields are path independent

Theorem: On a connected domain, a smooth vector field \mathbf{F} is path-independent iff it is conservative.

Proof: First, given a conservative field $\mathbf{F} = \nabla\phi$ for some smooth ϕ consider the line integral along some curve c from p_1 to p_2 . Let $\mathbf{r}(t)$ be a parameterization of the curve with $p_1 = \mathbf{r}(t_1)$ and $p_2 = \mathbf{r}(t_2)$. Now,

$$\int_c \mathbf{F} \cdot d\mathbf{l} = \int_{t_1}^{t_2} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt$$

$$\begin{aligned}
&= \int_{t_1}^{t_2} \nabla \phi \cdot \frac{d\mathbf{r}}{dt} dt \\
&= \int_{t_1}^{t_2} \left(\frac{\partial \phi}{\partial x} \frac{dx}{dt} + \frac{\partial \phi}{\partial y} \frac{dy}{dt} + \frac{\partial \phi}{\partial z} \frac{dz}{dt} \right) dt
\end{aligned} \tag{9}$$

where we have used the fact that \mathbf{F} is conservative and we have written the dot product out explicitly. Now, we use the chain rule to rewrite the dot product

$$\frac{\partial \phi}{\partial x} \frac{dx}{dt} + \frac{\partial \phi}{\partial y} \frac{dy}{dt} + \frac{\partial \phi}{\partial z} \frac{dz}{dt} = \frac{d\phi}{dt} \tag{10}$$

where $\phi(x, y, z)$ is a function of t along the curve through the t dependence of x , y and z : $\phi(t) = \phi(x(t), y(t), z(t))$. This gives

$$\int_c \mathbf{F} \cdot d\mathbf{l} = \int_{t_1}^{t_2} \frac{d\phi}{dt} dt = \phi(t) \Big|_{t_1}^{t_2} = \phi(t_2) - \phi(t_1) \tag{11}$$

which depends only on the value of the ϕ at the beginning and end of the curve and is therefore path independent.


To go the other way, for give path independent \mathbf{F} and some fixed point p let

$$\phi(\mathbf{r}) = \int_c \mathbf{F} \cdot d\mathbf{l} \tag{12}$$

where c is a path from p to \mathbf{r} . Since the field is path independent ϕ is well defined. It clearly depends on the choice of the reference point p and so ϕ isn't unique, however, two choices only differ by an overall constant. We now want to prove that $\mathbf{F} = \nabla \phi$. We begin by proving

$$F_1 = \frac{\partial \phi}{\partial x} \tag{13}$$

where $\mathbf{F} = F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k}$. If this hold it should hold for all paths, so lets choose a path $c = c_1 + c_2$ where c_1 goes from p to a point (x', y, z) and c_2 runs from (x', y, z) to (x, y, z) .



Now,

$$\begin{aligned}\phi(\mathbf{r}) &= \int_c \mathbf{F} \cdot d\mathbf{l} = \int_{c_1} \mathbf{F} \cdot d\mathbf{l} + \int_{c_2} \mathbf{F} \cdot d\mathbf{l} \\ &= \int_{c_1} \mathbf{F} \cdot d\mathbf{l} + \int_{x'}^x F_1 dx\end{aligned}\tag{14}$$

where in the last line we have used that $dy = dz = 0$ along the curve c_2 . Now, noting that the first integral doesn't depend on x , we differentiate

$$\frac{\partial}{\partial x} \phi = \frac{d}{dx} \int_{x'}^x F_1 dx = F_1\tag{15}$$

using the fundamental theorem of calculus. Now, a similar argument could be used for the other component and so the theorem is proved. \square

Conservative fields and irrotational fields

All conservative fields are irrotational because

$$\nabla \times \mathbf{F} = \nabla \times \nabla \phi = 0\tag{16}$$

but, the converse isn't true in general, an irrotational field is not necessarily conservative. However, Stoke's theorem can be used to prove that it is true locally: if a field is irrotational then for any point there is a neighbourhood of the point for which it is conservative.

A more general theorem applies to simply connected domain. A domain is *simply connected* if any two paths between the same two points can be deformed into each other smoothly. The plane is simply connected but the plane minus a point, say $\mathbf{R}^2 \setminus (0,0)$ ³ is not because the missing point gets in the way. So, in the $\mathbf{R}^2 \setminus (0,0)$ example the path from $(-1,0)$ to $(1,0)$ along the upper half circle can't be deformed into the path along the lower half circle because the missing point gets in the way. Anyway, without proof, we have a theorem:

Theorem: On a simply connected domain a smooth vector field is irrotational iff it is conservative.

³The backslash is a set minus, that is

$$\mathbf{R}^2 \setminus (0,0) = \{(x,y) \in \mathbf{R}^2 | (x,y) \neq (0,0)\}\tag{17}$$