231 Tutorial Sheet 11.[1,2]

20 January 2005

Useful facts:

- A function $f(x)$ has period l if $f(x+l) = f(x)$, it is odd if $f(-x) = -f(x)$ and even if $f(-x) = f(x)$.
- A function with period l has the Fourier series expansion

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{2\pi nx}{l}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{2\pi nx}{l}\right).$$

  where

$$\begin{aligned} a_0 &= \frac{2}{l}\int_{-l/2}^{l/2} f(x)dx \\ a_n &= \frac{2}{l}\int_{-l/2}^{l/2} f(x)\cos\left(\frac{2\pi nx}{l}\right)dx \\ b_n &= \frac{2}{l}\int_{-l/2}^{l/2} f(x)\sin\left(\frac{2\pi nx}{l}\right)dx \end{aligned}$$

- A function with period l has the Fourier series expansion

$$f(x) = \sum_{n=-\infty}^{\infty} c_n \exp\frac{2\pi nx}{l}.$$

  where

$$c_n = \frac{1}{l}\int_{-l/2}^{l/2} f(x)\exp\left(\frac{-2\pi nx}{l}\right)dx$$

[1]Conor Houghton, `houghton@maths.tcd.ie`, see also `http://www.maths.tcd.ie/~houghton/231`
[2]Including material from Chris Ford, to whom many thanks.

Questions

1. Express the following periodic functions ($l = 2\pi$) as complex Fourier series

   (a)
   $$f(x) = \begin{cases} 0 & -\pi < x < -a \\ 1 & -a < x < a \\ 0 & a < x < \pi \end{cases}$$
   where $a \in (0, \pi)$ is a constant.

   (b)
   $$f(x) = \frac{1}{2 - e^{ix}}.$$

2. Show that the periodic function f defined by $f(x) = |x| - \frac{1}{2}\pi$ for $-\pi < x < \pi$ and $f(x + 2\pi) = f(x)$ has the Fourier series expansion
   $$f(x) = -\frac{4}{\pi} \sum_{n>0,\ \text{odd}} \frac{\cos nx}{n^2}.$$

3. Use the Fourier series given in question 2 to compute the following sums
   $$S_1 = 1 - \frac{1}{3^2} - \frac{1}{5^2} + \frac{1}{7^2} + \frac{1}{9^2} - \frac{1}{11^2} - \frac{1}{13^2} + ...$$
   $$S_2 = 1 + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + ...$$
   Remark: With calculations of this kind it makes sense to try a quick numerical check of your answer.