

## 231 Tutorial Sheet 2<sup>12</sup>

21 October 2005

### Useful facts:

- The Jacobian in three-dimensions:

$$dx_1 dx_2 dx_3 = J dy_1 dy_2 dy_3 \quad (1)$$

where

$$J = \frac{\partial(x_1, x_2, x_3)}{\partial(y_1, y_2, y_3)} = \left\| \begin{array}{ccc} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} & \frac{\partial x_1}{\partial y_3} \\ \frac{\partial x_2}{\partial y_1} & \frac{\partial x_2}{\partial y_2} & \frac{\partial x_2}{\partial y_3} \\ \frac{\partial x_3}{\partial y_1} & \frac{\partial x_3}{\partial y_2} & \frac{\partial x_3}{\partial y_3} \end{array} \right\| \quad (2)$$

- For a scalar field  $\phi$  the gradient is

$$\text{grad } \phi = \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k} \quad (3)$$

- If  $\mathbf{a}$  is a vector  $\hat{\mathbf{a}}$  is the corresponding unit vector

$$\hat{\mathbf{a}} = \frac{1}{|\mathbf{a}|} \mathbf{a} \quad (4)$$

- The direction derivative of a scalar field  $f$  in the  $\mathbf{a}$  direction is  $D_{\mathbf{a}} f = \hat{\mathbf{a}} \cdot \nabla f$ .

- For a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  the divergence is

$$\text{div } \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \quad (5)$$

- For a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  the curl is

$$\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} \quad (6)$$

---

<sup>1</sup>Conor Houghton, [houghton@maths.tcd.ie](mailto:houghton@maths.tcd.ie), see also <http://www.maths.tcd.ie/~houghton/231>

<sup>2</sup>Including material from Chris Ford, to whom many thanks.

## Questions

1. Compute the Jacobian of the transformation from cartesian to parabolic cylinder coordinates

$$x = \frac{1}{2}(u^2 - v^2), \quad y = uv.$$

2. Determine the volume of the region enclosed by the cylinder  $x^2 + y^2 = 4$  and the planes  $y + z = 4$  and  $z = 0$ . *Suggestion: Use Cartesian coordinates.*
3. Check that the Jacobian for the transformation from cartesian to spherical polar coordinates is

$$J = r^2 \sin \theta.$$

Consider the hemisphere defined by

$$\sqrt{x^2 + y^2 + z^2} \leq 1, \quad z \geq 0.$$

Using spherical polar coordinates compute its volume and centroid.

4. Determine the curl of the vector fields
  - (a)  $\mathbf{F} = -yz \sin x \mathbf{i} + z \cos x \mathbf{j} + y \cos x \mathbf{k}$ .
  - (b)  $\mathbf{F} = \frac{1}{2}y \mathbf{i} - \frac{1}{2}x \mathbf{j}$ .
5. Show that away from the origin the vector field

$$\mathbf{F} = \frac{\hat{\mathbf{r}}}{r^3} = \frac{\mathbf{r}}{r^3}$$

is divergenceless.