

231 Tutorial Sheet 4¹²

5 November 2005

Useful facts:

- For a vector field $\mathbf{F} = (F_1, F_2, F_3)$ the curl is

$$\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} \quad (1)$$

- To evaluate the line integral for a parameterized curve:

$$\int_c \mathbf{F} \cdot d\mathbf{l} = \int_{t_1}^{t_2} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt \quad (2)$$

where t_1 and t_2 are the parameter values corresponding to the beginning and end of the curve.

- A vector field is *solenoidal* if it has zero divergence.
- A vector field is *irrotational* if it has zero curl.
- A vector field \mathbf{F} is *conservative* if there is a scalar field ϕ such that

$$\mathbf{F} = \nabla \phi \quad (3)$$

Questions

- Compute the line integrals:

(a) $\int_C (dx \, xy + \frac{1}{2} dy \, x^2 + dz)$ where C is the line segment joining the origin and the point $(1, 1, 2)$.

(b) $\int_C (dx \, yz + dy \, xz + dz \, yx^2)$ where C is the same line as in the previous part

- Is $F = \mathbf{r}/r$ irrotational? Is it conservative. Is it conservative on a restricted domain? It is path independent?

- Consider the vector field

$$\mathbf{F} = \frac{\frac{1}{2}y}{x^2 + y^2} \mathbf{i} - \frac{\frac{1}{2}x}{x^2 + y^2} \mathbf{j}. \quad (4)$$

(a) Determine the line integral $\oint_C d\mathbf{l} \cdot \mathbf{F}$ where C is the unit circle centred at the origin in the $z = 0$ plane (taken anti-clockwise).

(b) Compute the curl of \mathbf{F} .

(c) Is \mathbf{F} a conservative vector field?

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/231>

²Including material from Chris Ford, to whom many thanks.