

231 Tutorial Sheet 5¹²

11 November 2005

Useful facts:

- To evaluate the line integral for a parameterized curve:

$$\int_c \mathbf{F} \cdot d\mathbf{l} = \int_{t_1}^{t_2} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt \quad (1)$$

where t_1 and t_2 are the parameter values corresponding to the beginning and end of the curve.

- To evaluate the surface integral for a parameterized surface:

$$\iint_S \mathbf{F} \cdot d\mathbf{A} = \iint_S \mathbf{F} \cdot \left(\frac{d\mathbf{r}}{du} \times \frac{d\mathbf{r}}{dv} \right) du dv \quad (2)$$

Questions

- (a) Compute the flux of the vector field $\mathbf{F} = x\mathbf{i} + 2y\mathbf{j} + z\mathbf{k}$ out of the cylinder defined by $x^2 + y^2 = 1$ and $0 \leq z \leq 1$.
(b) What is the flux if the vector field is replaced with $\mathbf{F} = \mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$?
(c) Find the flux of $\mathbf{F} = z^2\mathbf{k}$ upwards through the part of the sphere $x^2 + y^2 + z^2 = a^2$ in the first octant of \mathbf{R}^3 (in the first octant all three coordinates x , y and z are positive).
- Find the flux of $\mathbf{F} = 2x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ upwards through the surface with parametrization $\mathbf{r}(u, v) = u^2v\mathbf{i} + uv^2\mathbf{j} + v^3\mathbf{k}$ where u and v range from 0 to 1.

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/231>

²Including material from Chris Ford, to whom many thanks.