

231 Outline Solutions Tutorial Sheet 4, 5 and 6.¹²

25 November 2005

Problem Sheet 4

1. Compute the line integrals:

- (a) $\int_C (dx \, xy + \frac{1}{2}dy \, x^2 + dz)$ where C is the line segment joining the origin and the point $(1, 1, 2)$.
(b) $\int_C (dx \, yz + dy \, xz + dz \, yx^2)$ where C is the same line as in the previous part

Solution: A quick way here is to note that \mathbf{F} is conservative.

$$\mathbf{F} = xy\mathbf{i} + \frac{1}{2}x^2\mathbf{j} + \mathbf{k} = \nabla\phi \quad (1)$$

where $\phi = \frac{1}{2}x^2y + z$. Hence

$$\int_C \mathbf{F} \cdot d\mathbf{l} = \phi(1, 1, 2) - \phi(0, 0, 0) = \frac{5}{2}. \quad (2)$$

For the next part, use the parametrization $x(u) = u$, $y(u) = u$, $z(u) = 2u$ ($0 \leq u \leq 1$).

$$\begin{aligned} \frac{d\mathbf{r}}{du} &= \mathbf{i} + \mathbf{j} + 2\mathbf{k}, \\ \mathbf{F} \cdot \frac{d\mathbf{r}}{du} &= 2u^2 + 2u^2 + 2u^3 = 4u^2 + 2u^3 \end{aligned} \quad (3)$$

so

$$\int_C \mathbf{F} \cdot d\mathbf{l} = \int_0^1 du \, (4u^2 + 2u^3) = \frac{4}{3} + \frac{1}{2} = \frac{11}{6}. \quad (4)$$

2. Is $F = \mathbf{r}/r$ irrotational? Is it conservative. Is it conservative on a restricted domain? It is path independent?

Solution: Well

$$\nabla r = \mathbf{r}/r \quad (5)$$

So this field is conservative and path independent. Interestingly, the field is not defined at $(0, 0, 0)$ and so this only applied on the restricted domain, however, the scalar potential is defined everywhere.

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/231>

²Including material from Chris Ford, to whom many thanks.

3. Consider the vector field

$$\mathbf{F} = \frac{\frac{1}{2}y}{x^2 + y^2} \mathbf{i} - \frac{\frac{1}{2}x}{x^2 + y^2} \mathbf{j}. \quad (6)$$

- (a) Determine the line integral $\oint_C \mathbf{dl} \cdot \mathbf{F}$ where C is the unit circle centred at the origin in the $z = 0$ plane (taken anti-clockwise).
- (b) Compute the curl of \mathbf{F} .
- (c) Is \mathbf{F} a conservative vector field?

Solution: Well on the circle

$$\mathbf{r} = \cos \theta \mathbf{i} + \sin \theta \mathbf{j} \quad (7)$$

and so

$$\frac{\partial \mathbf{r}}{\partial \theta} = -\sin \theta \mathbf{i} + \cos \theta \mathbf{j} \quad (8)$$

and the integral become

$$\oint \mathbf{F} \cdot \mathbf{dl} = - \int_0^{2\pi} d\theta = -2\pi \quad (9)$$

since $\mathbf{F} = -\cos \theta \mathbf{i} + \sin \theta \mathbf{j}$ on the circle. However, the curl does vanish, for convenient write $\rho = \sqrt{x^2 + y^2}$ and using the obvious notation

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial_x & \partial_y & \partial_z \\ \frac{y}{2\rho^2} & -\frac{x}{2\rho^2} & 0 \end{vmatrix} = \begin{pmatrix} 0 \\ 0 \\ \frac{\partial}{\partial x} \frac{x}{2\rho^2} + \frac{\partial}{\partial y} \frac{y}{2\rho^2} \end{pmatrix} \quad (10)$$

Now

$$\frac{\partial}{\partial x} \frac{x}{2\rho^2} = \frac{1}{2\rho^2} - \frac{x^2}{\rho^4} \quad (11)$$

and adding this to the similar result for y gives zero. So, it is not conservative, there is no contradiction here, the domain isn't simply connected, for the field to be defined the z -axis, along which $\rho = 0$ must be excluded and so for example, the unit circle we integrated around can't be shrunk to point. We will see later, problem sheet 6, that there is locally a scalar potential, but this isn't well defined over the whole domain.

Problem Sheet 5

1. (a) Compute the flux of the vector field $\mathbf{F} = x\mathbf{i} + 2y\mathbf{j} + z\mathbf{k}$ out of the cylinder defined by $x^2 + y^2 = 1$ and $0 \leq z \leq 1$.
- (b) What is the flux if the vector field is replaced with $\mathbf{F} = \mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$?
- (c) Find the flux of $\mathbf{F} = z^2\mathbf{k}$ upwards through the part of the sphere $x^2 + y^2 + z^2 = a^2$ in the first octant of \mathbf{R}^3 (in the first octant all three coordinates x , y and z are positive).

Solution: For the first part divide the cylinder up into a curved shaft and a disk at the top and bottom. For the curved part, use cylindrical polars with $\rho = 1$: $u = \theta$, $v = z$ so that $x = \cos u$, $y = \sin u$, $z = v$ or

$$\mathbf{r}(u, v) = \cos u \mathbf{i} + \sin u \mathbf{j} + v \mathbf{k} \quad (12)$$

and

$$\frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = \cos u \mathbf{i} + \sin u \mathbf{j}. \quad (13)$$

so

$$\mathbf{F} \cdot \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = x \cos u + 2y \sin u = \cos^2 u + 2 \sin^2 u. \quad (14)$$

Hence

$$\int_{\text{shaft}} \mathbf{F} \cdot d\mathbf{A} = \int_0^1 dv \int_0^{2\pi} du (\cos^2 u + 2 \sin^2 u) = 1(\pi + 2\pi) = 3\pi. \quad (15)$$

Now, the base has $z = 0$ and $\mathbf{n} = -\mathbf{k}$ so $\mathbf{F} \cdot \mathbf{n} = 0$, the top has $z = 1$ and $\mathbf{n} = \mathbf{k}$ and hence $\mathbf{F} \cdot \mathbf{n} = 1$ so

$$\int_{\text{top}} \mathbf{F} \cdot d\mathbf{A} = \pi \quad (16)$$

because the top is a disk of radius one and hence area π . Thus

$$\int_{\text{top}} \mathbf{F} \cdot d\mathbf{A} = 4\pi \quad (17)$$

If $\mathbf{F} = \mathbf{r}$ then the integral over the shaft becomes

$$\int_{\text{shaft}} \mathbf{F} \cdot d\mathbf{A} = \int_0^1 dv \int_0^{2\pi} du = 2\pi. \quad (18)$$

because

$$\mathbf{r} \cdot \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = \cos^2 u + \sin^2 u = 1 \quad (19)$$

and the surface area of the curved part is 2π . The flux out of the top and bottom is unchanged since only the \mathbf{k} component is relevant and this is the same, hence

$$\int_{\text{top}} \mathbf{F} \cdot d\mathbf{A} = 3\pi \quad (20)$$

Finally, the octant of the sphere is parameterized by

$$\begin{aligned}x &= a \cos \phi \sin \theta \\y &= a \sin \phi \sin \theta \\z &= a \cos \theta\end{aligned}\tag{21}$$

and so

$$\begin{aligned}\frac{\partial \mathbf{r}}{\partial \theta} &= a \cos \phi \cos \theta \mathbf{i} + a \sin \phi \cos \theta \mathbf{j} + -a \sin \theta \mathbf{k} \\ \frac{\partial \mathbf{r}}{\partial \phi} &= -a \sin \phi \sin \theta \mathbf{i} + a \cos \phi \sin \theta \mathbf{j}\end{aligned}\tag{22}$$

and hence

$$\frac{\partial \mathbf{r}}{\partial \theta} \times \frac{\partial \mathbf{r}}{\partial \phi} = a \sin \theta (x \mathbf{i} + y \mathbf{j} + z \mathbf{k})\tag{23}$$

This result can also be obtained directly by noting that the area element on a sphere of radius a is $dA = a^2 \sin \theta d\theta d\phi$ and the normal is \mathbf{r}/a . Now

$$\mathbf{F} \cdot d\mathbf{A} = az^3 \sin \theta d\theta d\phi\tag{24}$$

and the integral becomes

$$\int_S \mathbf{F} \cdot d\mathbf{A} = a^4 \int_0^{\pi/2} d\phi \int_0^{\pi/2} d\theta \cos^3 \theta \sin \theta = \frac{a^4 \pi}{2} \int_0^1 \theta d \cos \theta \cos^3 = \frac{a^4 \pi}{8}\tag{25}$$

2. Find the flux of $\mathbf{F} = 2x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ upwards through the surface with parametrization $\mathbf{r}(u, v) = u^2v\mathbf{i} + uv^2\mathbf{j} + v^3\mathbf{k}$ where u and v range from 0 to 1.

Solution: Here the parametrization of the surface is given $\mathbf{r} = u^2v\mathbf{i} + uv^2\mathbf{j} + v^3\mathbf{k}$.

$$\frac{\partial \mathbf{r}}{\partial u} = 2uv\mathbf{i} + v^2\mathbf{j}, \quad \frac{\partial \mathbf{r}}{\partial v} = u^2\mathbf{i} + 2uv\mathbf{j} + 3v^2\mathbf{k}.\tag{26}$$

A short calculation gives

$$\frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = 3v^4\mathbf{i} - 6uv^3\mathbf{j} + 3u^2v^2\mathbf{k}.\tag{27}$$

Now

$$\mathbf{F} \cdot \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = 3v^4 \cdot 2x - 6uv^3y + 3u^2v^2z = 3u^2v^5,\tag{28}$$

which gives

$$\int_S \mathbf{F} \cdot d\mathbf{A} = 3 \int_0^1 du \int_0^1 dv u^2v^5 = \frac{1}{6}.\tag{29}$$

Problem Sheet 6

- For each of the following vector fields compute the line integral $\oint_C \mathbf{F} \cdot d\mathbf{l}$ where C is the unit circle in the xy -plane taken anti-clockwise.

(a) $\mathbf{F} = x\mathbf{i} + y\mathbf{j}$

(b) $\mathbf{F} = y\mathbf{i} - x^2y\mathbf{j}$.

Solution: In the first part $\mathbf{F} = \nabla \frac{1}{2}(x^2 + y^2)$ so that \mathbf{F} is conservative giving $\oint_C \mathbf{F} \cdot d\mathbf{l}$. In the second part parametrize curve:

$$\begin{aligned} x(u) &= \cos u \\ y(u) &= \sin u \\ z(u) &= 0 \end{aligned} \quad (30)$$

where $0 \leq u \leq 2\pi$ or $\mathbf{r}(u) = \cos u\mathbf{i} + \sin u\mathbf{j}$. Now

$$\frac{d\mathbf{r}(u)}{du} = -\sin u\mathbf{i} + \cos u\mathbf{j}. \quad (31)$$

and

$$\mathbf{F} \cdot \frac{d\mathbf{r}}{du} = -y \sin u - x^2 y \cos u = -\sin^2 u - \cos^3 u \sin u. \quad (32)$$

Thus

$$\oint_C \mathbf{F} \cdot d\mathbf{l} = \int_0^{2\pi} du (-\sin^2 u - \cos^3 u \sin u) = -\pi, \quad (33)$$

since the average value of $\sin^2 u$ is $\frac{1}{2}$ and $\int_0^{2\pi} du \cos^3 u \sin u = 0$ by symmetry.

- Compute the flux of the vector field $\mathbf{F} = (x + x^2)\mathbf{i} + y\mathbf{j}$ out of the cylinder defined by $x^2 + y^2 = 1$ and $0 \leq z \leq 1$.

Solution: As problem sheet 5, $\mathbf{r}(u, v) = \cos u\mathbf{i} + \sin u\mathbf{j} + v\mathbf{k}$ giving

$$\frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = \cos u\mathbf{i} + \sin u\mathbf{j}. \quad (34)$$

and

$$\mathbf{F} \cdot \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} = (x + x^2) \cos u + y \sin u = \cos^2 u + \cos^3 u + \sin^2 u = 1 + \cos^3 u. \quad (35)$$

and so, noting that the \mathbf{F} is perpendicular to the normal at both ends and so we need only include the curved surface

$$\int_S \mathbf{F} \cdot d\mathbf{A} = \int_0^1 dv \int_0^{2\pi} du (1 + \cos^3 u) = 2\pi, \quad (36)$$

since the $\cos^3 u$ integral is zero by symmetry.

Note: This problem can be solved by noting that $x^2\mathbf{i}$ makes no contribution (by symmetry) and $\mathbf{F} = x\mathbf{i} + y\mathbf{j}$ has flux 2π since $\mathbf{F} \cdot \mathbf{n} = 1$.

3. Find the flux of $\mathbf{F} = z^3 \mathbf{k}$ upwards through the part of the sphere $x^2 + y^2 + z^2 = a^2$ above the $z = 0$ plane.

Solution: To parametrize the sphere choose $(u, v) = (\theta, \phi)$, that is, spherical polar angles. Since the radius $r = a$ this gives $x(u, v) = a \sin u \cos v$, $y(u, v) = a \sin u \sin v$, $z(u, v) = a \cos u$ with $0 \leq \theta \leq \pi/2$ and $0 \leq \phi < 2\pi$. Since \mathbf{F} only has a non-zero z -component we just need

$$\begin{aligned} \left(\frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} \right)_3 &= \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial x}{\partial v} \\ &= a^2 (\cos u \cos v \sin u \cos v + \cos u \sin v \sin u \sin v) \\ &= a^2 \cos u \sin u, \end{aligned} \quad (37)$$

which is positive. So the orientation is upwards. Now $F_z = z^3 = a^3 \cos^3 u$, and so

$$\begin{aligned} \int_S \mathbf{F} \cdot d\mathbf{A} &= a^5 \int_0^{\frac{1}{2}\pi} du \int_0^{2\pi} dv \cos^4 u \sin u \\ &= 2\pi a^5 \int_0^{\frac{1}{2}\pi} du \cos^4 u \sin u \\ &= 2\pi a^5 \left[-\frac{\cos^5 u}{5} \right]_0^{\frac{1}{2}\pi} = \frac{2\pi a^5}{5}. \end{aligned} \quad (38)$$

4. Consider the ‘point vortex’ vector field

$$\mathbf{F} = \frac{y}{x^2 + y^2} \mathbf{i} - \frac{x}{x^2 + y^2} \mathbf{j}.$$

Show that $\text{curl } \mathbf{F} = 0$ away from the z -axis. Establish that \mathbf{F} is *not* conservative in the (non simply-connected) domain $x^2 + y^2 \geq \frac{1}{2}$. Is \mathbf{F} conservative in the domain defined by $x^2 + y^2 \geq \frac{1}{2}$, $y \geq 0$? If so obtain a scalar potential for \mathbf{F} .

Solution:

$$\begin{aligned} \nabla \times \mathbf{F} &= \frac{1}{2} \mathbf{k} \left[\partial_x \left(\frac{-x}{x^2 + y^2} \right) + \partial_y \left(\frac{y}{x^2 + y^2} \right) \right] \\ &= \frac{1}{2} \mathbf{k} \left[-\frac{1}{x^2 + y^2} + \frac{2x^2}{(x^2 + y^2)^2} - \frac{1}{x^2 + y^2} + \frac{2y^2}{(x^2 + y^2)^2} \right] \\ &= 0. \end{aligned} \quad (39)$$

To show that \mathbf{F} is not conservative consider $\oint_C \mathbf{F} \cdot d\mathbf{l}$ where C is the unit circle. Using the obvious parametrization

$$\begin{aligned} \oint_C \mathbf{F} \cdot d\mathbf{l} &= \int_0^{2\pi} du (-\sin^2 u - \cos^2 u) \\ &= -2\pi \neq 0, \end{aligned} \quad (40)$$

therefore \mathbf{F} is not conservative.

The domain $x^2 + y^2 \geq \frac{1}{2}$, $y \geq 0$ is simply connected and \mathbf{F} is irrotational and smooth is the domain. Thus \mathbf{F} is conservative.

Write $\mathbf{F} = \nabla\phi$. Seek a $\phi(x, y)$ such that

$$\frac{\partial\phi}{\partial x} = \frac{y}{x^2 + y^2}, \quad \frac{\partial\phi}{\partial y} = -\frac{x}{x^2 + y^2}. \quad (41)$$

Integrate first equation by treating y as a constant

$$\phi(x, y) = y \int \frac{dx}{x^2 + y^2} = \tan^{-1} \frac{x}{y} + C(y). \quad (42)$$

Assume that x and y are non-negative, then

$$\tan^{-1} \frac{x}{y} + \tan^{-1} \frac{y}{x} = \frac{\pi}{2},$$

so that $\phi(x, y) = -\tan^{-1} \frac{y}{x} +$ a possibly y -dependent constant. However it is easy to check that $\phi = -\tan^{-1} \frac{y}{x}$ satisfies $\frac{\partial\phi}{\partial y} = -\frac{x}{x^2 + y^2}$. Clearly, $\tan^{-1} \frac{y}{x}$ is the usual polar angle θ , that is $\phi = -\theta$.

Can try to extend this back to the original domain $x^2 + y^2 \geq \frac{1}{2}$, but ϕ will suffer a branch cut discontinuity at, say $\theta = \frac{3}{2}\pi$.

5. Let D be a plane region with area A whose boundary is a piecewise smooth closed curve C . Use Green's theorem to prove that the centroid (\bar{x}, \bar{y}) of D is

$$\begin{aligned} \bar{x} &= \frac{1}{2A} \oint_C dy x^2 \\ \bar{y} &= -\frac{1}{2A} \oint_C dx y^2. \end{aligned} \quad (43)$$

Use this result to compute the centroid of a semi-circle (this was determined in the lectures using the more standard formula).

Solution: The centroid (\bar{x}, \bar{y}) of a plane region D is given by

$$\bar{x} = \frac{\int_D dA x}{A} \quad \bar{y} = \frac{\int_D dA y}{A}. \quad (44)$$

If the boundary of D is a piecewise smooth closed curve C , Green's theorem reads

$$\int_D dA \left(\frac{\partial g(x, y)}{\partial x} - \frac{\partial f(x, y)}{\partial y} \right) = \oint_C (dx f(x, y) + dy g(x, y)), \quad (45)$$

where $f(x, y)$ and $g(x, y)$ are function with continuous first derivatives and the curve is oriented anti-clockwise. Now taking $g(x, y) = \frac{1}{2}x^2$ and $f(x, y) = 0$ yields

$$\int_D dA x = \frac{1}{2} \int_C dy x^2. \quad (46)$$

Inserting this into the formula for \bar{x} gives

$$\bar{x} = \frac{1}{2A} \oint_C dy x^2 \quad (47)$$

Similarly the choice $f(x, y) = \frac{1}{2}y^2$, $g(x, y) = 0$ gives

$$\bar{y} = -\frac{1}{2A} \oint_C dx y^2 \quad (48)$$

Here C comprises the semi-circular arc plus the line segment joining $(-1, 0)$ and $(1, 0)$. The integral $\int_C dy x^2$ is zero since the positive x part of the arc integral cancels the negative x part. Also the integral along the line segment is zero. This implies that $\bar{x} = 0$.

For the other integral, $\int_C dx y^2$, only the arc contributes since $y = 0$ along the line segment. Now

$$\int_C dx y^2 = - \int_{C, \text{clockwise}} dx y^2 = - \int_{-1}^1 dx (1 - x^2) = -\frac{4}{3}. \quad (49)$$

Since $A = \frac{1}{2}\pi$ it follows that

$$\bar{y} = \frac{\frac{4}{3}}{2A} = \frac{4}{3\pi}. \quad (50)$$