

## 231 Tutorial Sheet 7<sup>12</sup>

24 November 2006

### Useful facts:

- To evaluate the line integral for a parameterized curve:

$$\int_c \mathbf{F} \cdot d\mathbf{l} = \int_{t_1}^{t_2} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt \quad (1)$$

where  $t_1$  and  $t_2$  are the parameter values corresponding to the beginning and end of the curve.

- To evaluate the surface integral for a parameterized surface:

$$\int \int_S \mathbf{F} \cdot d\mathbf{A} = \int \int_S \mathbf{F} \cdot \left( \frac{d\mathbf{r}}{du} \times \frac{d\mathbf{r}}{dv} \right) du dv \quad (2)$$

- Stokes' theorem: for an orientable piecewise smooth surface  $S$  with an orientable piecewise smooth boundary  $C$  oriented so that  $\mathbf{N} \times d\mathbf{l}$  points into  $S$  with  $\mathbf{n}$  the normal and  $d\mathbf{l}$  a tangent to  $C$  and  $\mathbf{F}$  a vector field defined in a region containing  $S$ , then

$$\int_S \text{curl } \mathbf{F} \cdot d\mathbf{A} = \int_C \mathbf{F} \cdot d\mathbf{l} \quad (3)$$

- If  $S$  is a surface  $\partial S$  is its bounding curve with the orientation describe above.
- Green's theorem on the plane: let  $D$  be a region in the  $xy$ -plane bounded by a piecewise continuous curve  $C$ , if  $f(x, y)$  and  $g(x, y)$  have continuous first derivatives

$$\int_D \left( \frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dx dy = \int_C (f dx + g dy) \quad (4)$$

---

<sup>1</sup>Conor Houghton, [houghton@maths.tcd.ie](mailto:houghton@maths.tcd.ie), see also <http://www.maths.tcd.ie/~houghton/231>

<sup>2</sup>Including material from Chris Ford, to whom many thanks.

## Questions

1. Which of the following vector fields are conservative?
  - (a)  $\mathbf{F} = -yz \sin x \mathbf{i} + z \cos x \mathbf{j} + y \cos x \mathbf{k}$ .
  - (b)  $\mathbf{F} = \frac{1}{2}y \mathbf{i} - \frac{1}{2}x \mathbf{j}$ .
  - (c)  $\mathbf{F} = \frac{1}{2}(\mathbf{B} \times \mathbf{r})$  where  $\mathbf{B}$  is a constant vector.
2. Compute the flux of the vector field  $\mathbf{F} = (x + x^2)\mathbf{i} + y\mathbf{j}$  out of the cylinder defined by  $x^2 + y^2 = 1$  and  $0 \leq z \leq 1$ .
3. Find the flux of  $\mathbf{F} = z^3\mathbf{k}$  upwards through the part of the sphere  $x^2 + y^2 + z^2 = a^2$  above the  $z = 0$  plane.
4. Let  $D$  be a plane region with area  $A$  whose boundary is a piecewise smooth closed curve  $C$ . Use Green's theorem to prove that the centroid  $(\bar{x}, \bar{y})$  of  $D$  is

$$\begin{aligned}\bar{x} &= \frac{1}{2A} \oint_C dy \, x^2 \\ \bar{y} &= -\frac{1}{2A} \oint_C dx \, y^2.\end{aligned}\tag{5}$$

Use this result to compute the centroid of a semi-circle.