

## 231 Tutorial Sheet 17<sup>12</sup>

3 April 2008

### Useful facts:

- Series solution: assume there is a solution of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

and, by substituting into the equation find a recursion relation: an equation relating higher terms in  $a_n$  to lower one.

- By expanding out the sum it is easy to see  $y(0) = a_0$  and  $y'(0) = a_1$
- The method of Frobenius: assume there is a solution of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+s}$$

under the assumption that  $a_0 \neq 0$ , this assumption will give an equation for  $s$  called the indicial equation.

---

<sup>1</sup>Conor Houghton, [houghton@maths.tcd.ie](mailto:houghton@maths.tcd.ie), see also <http://www.maths.tcd.ie/~houghton/231>

<sup>2</sup>Including material from Chris Ford, to whom many thanks.

## Questions

1. Use the recursion relation

$$a_{n+2} = \frac{2(n - \alpha)a_n}{(n + 1)(n + 2)}$$

or the generating function

$$\Phi(x, h) = e^{2xh - h^2} = \sum_{n=0}^{\infty} \frac{h^n}{n!} H_n(x)$$

to obtain polynomial solutions of Hermite's equation  $y'' - 2xy' + 2\alpha y = 0$  for  $\alpha = 3, 4$  and  $5$ .

2. Legendre's equation can be written

$$(1 - x^2)y'' - 2xy' + \alpha y = 0,$$

where  $\alpha$  is a constant. Consider a series solution of the form

$$y(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Determine a recursion relation for the  $a_n$  coefficients. For what values of  $\alpha$  does Legendre's equation have polynomial solutions?

3. (Frobenius training exercise) For each of the following equations obtain the indicial equation for a Frobenius series of the form

$$y(x) = \sum_{n=0}^{\infty} a_n x^{n+s}$$

- (a)  $y'' + y = 0$ .
- (b)  $x^2 y'' + 3xy' + y = 0$
- (c)  $4xy'' + 2y' + y = 0$ .

In case a) use the method of Frobenius to obtain the general solution. In case b) use the method of Frobenius to find one solution (the method fails to give the other solution).

4. Use direct substitution to show that the functions  $H_n$  defined through the generating function

$$\Phi(x, h) = e^{2xh - h^2} = \sum_{n=0}^{\infty} \frac{h^n}{n!} H_n(x)$$

satisfy Hermite's equation

$$y'' - 2xy' + 2ny = 0.$$