## **231** Tutorial Sheet $5^{12}$

## 9 November 2007

## Useful facts:

• For a scalar field  $\phi$  the gradiant is

grad 
$$\phi = \frac{\partial \phi}{\partial x}\mathbf{i} + \frac{\partial \phi}{\partial y}\mathbf{j} + \frac{\partial \phi}{\partial z}\mathbf{k}$$
 (1)

• For a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  the divergence is

div 
$$\mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$
 (2)

• For a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  the curl is

$$\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix}$$
(3)

• The Laplacian:

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \tag{4}$$

• To evaluate the line integral for a parameterized curve:

$$\int_{c} \mathbf{F} \cdot \mathbf{dl} = \int_{t_1}^{t_2} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} dt$$
(5)

where  $t_1$  and  $t_2$  are the parameter values corresponding to the beginning and end of the curve.

- A vector field is *solinoidal* if it has zero divergence.
- A vector field is *irrotational* if it has zero curl.

<sup>&</sup>lt;sup>1</sup>Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/231 <sup>2</sup>Including material from Chris Ford, to whom many thanks.

## Questions

- 1. Calculate curl  $\mathbf{r}/r$  and div  $\mathbf{r}/r$  away from the origin. What is  $\Delta r$ ?
- 2. Prove the identity

$$\nabla \cdot (\nabla \times \mathbf{F}) = 0 \tag{6}$$

3. Prove the identity

$$\nabla \times (\nabla \times \mathbf{F}) = \nabla (\nabla \cdot \mathbf{F}) - \triangle \mathbf{F}.$$
(7)

- 4. Compute the line integrals:
  - (a)  $\int_C (dx xy + \frac{1}{2}dy x^2 + dz)$  where C is the line segment joining the origin and the point (1, 1, 2).
  - (b)  $\int_C (dx \ yz \ + dy \ xz \ + dz \ yx^2)$  where C is the same line as in the previous part
- 5. For each of the following vector fields compute the line integral  $\oint_C \mathbf{F} \cdot \mathbf{dl}$  where C is the unit circle in the xy-plane taken anti-clockwise.

(a) 
$$\mathbf{F} = x\mathbf{i} + y\mathbf{j}$$

(b)  $\mathbf{F} = y\mathbf{i} - x^2 y\mathbf{j}$ .