

Gated channels¹

22 October 2008

So the persistent Potassium channel has gating probability n^4 , the transient sodium channel, m^3h . The n , m and h satisfy

$$\frac{d\ell}{dt} = \alpha_\ell(V)(1 - \ell) - \beta_\ell(V)\ell \quad (1)$$

which can be rewritten as

$$\tau_\ell \frac{d\ell}{dt} = \ell_\infty - \ell \quad (2)$$

where

$$\tau_\ell = \frac{1}{\alpha_\ell + \beta_\ell} \quad (3)$$

and

$$\ell_\infty = \frac{\alpha_\ell}{\alpha_\ell + \beta_\ell} \quad (4)$$

A standard example set of functional forms for the alpha and beta are given in Dayan and Abbott:

$$\begin{aligned} \alpha_n &= \frac{.01(V + 55)}{1 - \exp(-.1(V + 55))} \\ \beta_n &= .125 \exp(-.0125(V + 65)) \end{aligned} \quad (5)$$

for n , for m

$$\begin{aligned} \alpha_m &= \frac{.1(V + 40)}{1 - \exp(-.1(V + 40))} \\ \beta_m &= 4 \exp(-.0556(V + 65)) \end{aligned} \quad (6)$$

and for h

$$\begin{aligned} \alpha_h &= .07 \exp(-.05(V + 65)) \\ \beta_h &= \frac{1}{1 + \exp(-.1(V + 35))} \end{aligned} \quad (7)$$

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/231>