## MA22S3 Tutorial Sheet 8.<sup>1</sup>

## 3 December 2009

## Useful facts:

- To solve the equation  $a\ddot{y} + b\dot{y} + cy = 0$ , with a, b and c constants, use an exponential substitution  $y = \exp(\lambda t)$  to get the auxiliary equation  $a\lambda^2 + b\lambda + c = 0$  and solve for  $\lambda$ . Usually this will give two solutions.
- If there is only one  $\lambda$  then  $y = t \exp(\lambda t)$  will also be a soln.
- The solution of the equation  $a\ddot{y} + b\dot{y} + cy = f(t)$  is  $y = y_c + y_p$  where  $y_c = C_1y_1 + C_2y_2$  is the solution of  $a\ddot{y} + b\dot{y} + cy = 0$ .
- If  $f(t) = \exp(\mu t)$  and  $\mu$  isn't a solution of the auxiliary equation, then substitute  $y = C \exp(\mu t)$  and solve for C. If  $\mu$  is a solution to the auxiliary equation, use  $y = Ct \exp(\mu t)$  or even  $y = Ct^2 \exp(\mu t)$  if the auxiliary equation has two repeated roots.
- If  $f(t) = f_1(t) + f_2(t)$  then solve  $a\ddot{y} + b\dot{y} + cy = f_1(t)$  and  $a\ddot{y} + b\dot{y} + cy = f_2(t)$  and add the solutions.
- $\cosh t = [\exp(t) + \exp(-t)]/2$

## Questions

1. (2) Obtain the general solution to

$$\ddot{y} + 8\dot{y} + 16y = 0 \tag{1}$$

2. (2) Obtain the general solution to

$$\ddot{y} + 16y = 0 \tag{2}$$

3. (2) Obtain the solution to

$$\ddot{y} + \dot{y} - 2y = e^{5t} \tag{3}$$

with  $y(0) = \dot{y}(0) = 0$ .

4. (2) Obtain the general solution to

$$\ddot{y} + 8\dot{y} + 16y = 4\cosh t \tag{4}$$

<sup>&</sup>lt;sup>1</sup>Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/MA22S3