

MA22S3 Tutorial Sheet 3.¹²

27 October 2010

Useful facts:

- **Exponential with imaginary argument:**

$$e^{i\theta} = \cos \theta + i \sin \theta \quad (1)$$

- **Cosine and sine in terms of exponentials:**

$$\begin{aligned} \cos \theta &= \frac{e^{i\theta} + e^{-i\theta}}{2} \\ \sin \theta &= \frac{e^{i\theta} - e^{-i\theta}}{2i} \end{aligned} \quad (2)$$

- The **Fourier integral** or Fourier transform:

$$\begin{aligned} f(t) &= \int_{-\infty}^{\infty} dk \widetilde{f(k)} e^{ikt} \\ \widetilde{f(k)} &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dt f(t) e^{-ikt} \end{aligned}$$

Questions

1. (4) Compute the Fourier transform of $f(t) = e^{-a|t|}$ where a is a positive constant.
2. (4) Express the following function as a Fourier integral:

$$f(x) = \begin{cases} \cos t & |t| < \frac{\pi}{2} \\ 0 & |t| > \frac{\pi}{2} \end{cases}$$

One way to do the required integral is to split the cosine into exponentials.

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/MA22S3>

²Including material from Chris Ford, to whom many thanks.