

MA22S3 Outline Solutions for Tutorial Sheet 8.¹²

16 December 2010.

Questions

1. (2) Solve $t^2\ddot{y} + 4t\dot{y} + 2y = 0$.

Solution: The standard substitution $x = e^z$ gives

$$\frac{d^2y}{dz^2} + (4-1)\frac{dy}{dz} + 2y = 0. \quad (1)$$

Auxiliary equation is $\lambda^2 + 3\lambda + 2 = 0$ with roots which has roots $\lambda_1 = -2$ and $\lambda_2 = -1$:

$$y = C_1e^{-2z} + C_2e^{-z} = C_1t^{-2} + C_2t^{-1} \quad (2)$$

2. (2) Solve $x^2\ddot{y} + 4t\dot{y} + y = t^5$. In this example there is a forcing term, but substituting for t^5 in terms of z will just give a forced damped harmonic oscillator which can be solved then in the usual way.

Solution: So the same substitution gives

$$\frac{d^2y}{dz^2} + 3\frac{dy}{dz} + 2y = e^{5z} \quad (3)$$

So, we already have the complementary function for this, we just need the particular integral, substitute $y = C \exp(5z)$ giving

$$25C + 15C + 2C = 1 \quad (4)$$

giving

$$y = C_1t^{-2} + C_2t^{-1} + \frac{1}{42}t^5. \quad (5)$$

3. (4) Assuming the solution of

$$(1-t)\dot{y} + y = 0 \quad (6)$$

has a series expansion about $t = 0$ work out the recursion relation. Write out the first few terms and notice that the series $a_2 = 0$ so the series actually terminates to give $y = a_0(1-t)$. What is the solution with $y(0) = 2$.

Solution: Well we begin by writing

$$y = \sum_{n=0}^{\infty} a_n t^n \quad (7)$$

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/231>

²Including material from Chris Ford, to whom many thanks.

and so by differentiation we get

$$\dot{y} = \sum_{n=0}^{\infty} a_n n t^{n-1} \quad (8)$$

and hence

$$t\dot{y} = \sum_{n=0}^{\infty} a_n n t^n. \quad (9)$$

Thus, substituting the differential equation we get

$$\sum_{n=0}^{\infty} a_n n t^{n-1} - \sum_{n=0}^{\infty} a_n n t^n + \sum_{n=0}^{\infty} a_n t^n = 0 \quad (10)$$

In order to make progress we need to rewrite the first of these three series so that it is in the form

$$\sum_{n=0}^{\infty} \text{stuff}_n t^n \quad (11)$$

so that all three bits in the equation match. Well, let $m = n - 1$ in the expression for \dot{y} , (8), to get

$$\dot{y} = \sum_{m=0}^{\infty} a_{m+1} (m+1) t^m. \quad (12)$$

In fact, this looks at first like it gives

$$\dot{y} = \sum_{m=-1}^{\infty} a_{m+1} (m+1) t^m \quad (13)$$

but the $m = -1$ term is zero, so that's fine. Now m is just an index so we can rename it n , don't get confused, this isn't the original n , we just want all parts of the equation to look the same.

In fact, we now have

$$\sum_{n=0}^{\infty} a_{n+1} (n+1) t^n - \sum_{n=0}^{\infty} a_n n t^n + \sum_{n=0}^{\infty} a_n t^n = 0 \quad (14)$$

and we can group this all together to give

$$\sum_{n=0}^{\infty} [a_{n+1} (n+1) + (1-n) a_n] t^n = 0. \quad (15)$$

The recursion relation is

$$a_{n+1} = - \left(\frac{1-n}{1+n} \right) a_n \quad (16)$$

and this applies to n from zero upwards since that is what appears in the sum sign.

Starting at $n = 0$ we have

$$a_1 = -a_0. \tag{17}$$

For $n = 1$ we get

$$a_2 = 0 \tag{18}$$

and the series terminates here because every term is something multiplied by the one before and so if a_2 is zero the rest of the series is zero. Thus $y = a_0(1 - t)$ for arbitrary a_0 . If $y(0) = 2$ then $a_0 = 2$ and $y = 2(1 - t)$.