MA22S3 ODE work sheet - outline solutions questions 8-9.¹

15 May 2011

8 Find the general solution for $\ddot{y} - 5\dot{y} + 6y = e^{3t}$, what is the solution is y(0) = 4 and $\dot{y}(0) = 1$.

Solution: So in this case we already know the complementary function, it is

$$y_c = C_1 e^{3t} + C_2 e^{2t} \tag{1}$$

To find the particular function we first notice that the exponential on the right hand side, $\exp 3t$, already appears in the complementary function, so the appropriate ansatz is $y = Ct \exp 3t$. Now

$$\dot{y} = Ce^{3t} + 3Cte^{3t} \ddot{y} = 6Ce^{3t} + 9Cte^{3t}$$
(2)

Substituting back into the differential equation and cancelling the $\exp 3t$ we get

$$6C + 9Ct - 5C - 15Ct + 6Ct = 1 \tag{3}$$

of C = 1, hence the general solution is

$$y = te^{3t} + C_1 e^{3t} + C_2 e^{2t} (4)$$

Now

$$y(0) = C_1 + C_2 = 4 \tag{5}$$

and

$$\dot{y} = 1 + 3C_1 + 2C_2 = 1 \tag{6}$$

so $C_1 = -8$ and $C_2 = 12$ and

$$y = te^{3t} - 8e^{3t} + 12e^{2t} \tag{7}$$

9 Find the general solution for $\ddot{y} - 6\dot{y} + 9y = e^{3t}$, what is the solution is y(0) = 4 and $\dot{y}(0) = 1$.

Solution: This time we do need to calculate the complementary function, the auxiliary equation is

$$\lambda^{2} - 6\lambda + 9 = (\lambda - 3)^{2} = 0$$
(8)

so the complementary function is

$$y_c = C_1 e^{3t} + C_2 t e^{3t} (9)$$

¹Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/MA22S3

Now, the trouble is that this means that the ansatz for the particular solution has to be $y = Ct^2 \exp 3t$ so

$$\dot{y} = 2Cte^{3t} + 3Ct^2e^{3t} \tag{10}$$

and

$$\ddot{y} = 12Cte^{3t} + 2Ce^{3t} + 9Ct^2e^{3t} \tag{11}$$

and so, substituting in and cancelling the $\exp 3t$

$$12Ct + 2C + 9Ct^{2} - 12Ct - 18Ct^{2} + 9Ct^{2} = 1$$
(12)

soC = 1/2. Now

$$y = \frac{1}{2}t^2e^{3t} + C_1e^{3t} + C_2te^{3t}$$
(13)

SO

$$y(0) = C_1 = 4 \tag{14}$$

and

$$\dot{y}(0) = C_1 + C_2 = 1 \tag{15}$$

so $C_2 = -3$ and

$$y = \frac{1}{2}t^2e^{3t} + 4e^{3t} - 3te^{3t} \tag{16}$$