

442 Tutorial Sheet 1¹

28 October 2004

1. (2) Find an expression for

$$\nabla_c \nabla_d T^a_b - \nabla_d \nabla_c T^a_b \quad (1)$$

in terms of the Riemann tensor.

2. (2) Calculate the usual metric on the surface of a sphere by considering a radius r sphere $x^2 + y^2 + z^2 = r^2$ embedded in three-dimensional flat space $ds^2 = dx^2 + dy^2 + dz^2$. To do this, change to spherical polar coördinates:

$$\begin{aligned} x &= r \cos \phi \sin \theta \\ y &= r \sin \phi \sin \theta \\ z &= r \cos \theta \end{aligned} \quad (2)$$

and then set $dr = 0$ to restrict to surface of the sphere.

3. (4) Find the curvature on a two-dimensional hyperboloid:

$$t^2 - x^2 - y^2 = r^2 \quad (3)$$

embedded in Minkowski space:

$$ds^2 = -dt^2 + dx^2 + dy^2 \quad (4)$$

In other words, change to hyperbolic coördinates

$$\begin{aligned} x &= r \cos \phi \sinh \eta \\ y &= r \sin \phi \sinh \eta \\ t &= r \cosh \eta \end{aligned} \quad (5)$$

and then restrict to the surface of the hyperboloid by setting $dr = 0$. This gives the metric on the surface of the embedded hyperboloid, now calculate its Ricci scalar.

¹Conor Houghton, houghton@maths.tcd.ie, see also <http://www.maths.tcd.ie/~houghton/442.html>