442 Tutorial Sheet 5 Solutions¹

24 January 2005

1. (2) This question is revision of Lagrangian mechanics. By varying the field derive the Euler-Lagrance equations from $\delta S = 0$ where

$$S = \int L(t, \mathbf{q}, \dot{\mathbf{q}}) dt \tag{1}$$

Solution: Well vary q to get

$$\delta L(t, \mathbf{q}, \dot{\mathbf{q}}) = \sum_{i} \frac{\partial L}{\partial q_i} \delta q_i + \sum_{i} \frac{\partial L}{\partial \dot{q_i}} \delta \dot{q_i}$$
(2)

or

$$\delta S = \int \left(\sum_{i} \frac{\partial L}{\partial q_{i}} \delta q_{i} + \sum_{i} \frac{\partial L}{\partial \dot{q}_{i}} \delta \dot{q}_{i} \right) dt \tag{3}$$

and then use integration by part on the second term, assuming that there are fixed boundary conditions there is no boundary term and

$$\delta S = \int \left(\sum_{i} \frac{\partial L}{\partial q_i} - \sum_{i} \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} \right) \delta q_i dt \tag{4}$$

and if this is true for all variations satisfying suitable boundary conditions, we get the Euler-Lagrange equation

$$\sum_{i} \frac{\partial L}{\partial q_i} - \sum_{i} \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} = 0$$
(5)

2. (2) Derive the Euler-Largrange equation

$$\partial_{\mu} \frac{\partial \mathcal{L}}{\partial(\partial_{\mu}\phi)} - \frac{\partial \mathcal{L}}{\partial\phi} = 0 \tag{6}$$

from $\delta S = 0$ where

$$S = \int \mathcal{L}(t,\phi,\partial_{\mu}\phi)d^{4}x$$
(7)

¹Conor Houghton, houghton@maths.tcd.ie, see also http://www.maths.tcd.ie/~houghton/442.html

Solution: Well this is almost the same, vary ϕ and we get, by Taylor expanding $\mathcal L$

$$\delta S = \int \left(\frac{\partial \mathcal{L}}{\partial \phi} \delta \phi + \frac{\partial \mathcal{L}}{\partial (\partial_{\mu} \phi)} \partial_{\mu} \delta \phi \right) d^{4}x \tag{8}$$

and then use a Gauss law and zero boundary conditions to give

$$\delta S = \int \left(\frac{\partial \mathcal{L}}{\partial \phi} - \frac{\partial}{\partial x^{\mu}} \frac{\partial \mathcal{L}}{\partial (\partial_{\mu} \phi)} \right) \delta \phi d^4 x \tag{9}$$

and arbitrarity of $\delta\phi$ gives the Euler-Lagrange equations.

3. (3) Derive the Euler-Largange equation

$$\nabla_{\mu} \frac{\partial \mathcal{L}}{\partial (\nabla_{\mu} \phi)} - \frac{\partial \mathcal{L}}{\partial \phi} = 0 \tag{10}$$

from $\delta S = 0$ where

$$S = \int \mathcal{L}(t,\phi,\nabla_{\mu}\phi)\sqrt{g}d^{4}x$$
(11)

Recall that $\nabla_{\mu}\phi = \partial_{\mu}\phi$ and that $\sqrt{g}\nabla_{\mu}X^{\mu} = \partial_{\mu}\sqrt{g}X^{\mu}$.

Solution: This one isn't really so different, bearing in mind $\nabla_{\mu}\phi = \partial_{\mu}\phi$ nothing changes until we use the Gauss law, then we get

$$\delta S = \int \left(\frac{\partial \mathcal{L}}{\partial \phi} \sqrt{g} - \frac{\partial}{\partial x^{\mu}} \sqrt{g} \frac{\partial \mathcal{L}}{\partial (\partial_{\mu} \phi)} \right) \delta \phi d^4 x \tag{12}$$

and then commute the \sqrt{g} through to give

$$\delta S = \int \left(\frac{\partial \mathcal{L}}{\partial \phi} \sqrt{g} - \sqrt{g} \nabla_{\mu} \frac{\partial \mathcal{L}}{\partial (\partial_{\mu} \phi)} \right) \delta \phi d^4 x \tag{13}$$

and rewrite the $\partial_{\mu}\phi$ s as $\nabla_{\mu}\phi$ to get the answer.

4. (4) Calculate the energy-momentum tensor

$$T_{\mu\nu} = F_{\lambda\mu}F^{\lambda}_{\ \nu} - \frac{1}{4}g_{\mu\nu}F_{\lambda\rho}F^{\lambda\rho}$$
(14)

for the Maxwell field:

$$S = -\frac{1}{4} \int F^{\mu\nu} F_{\mu\nu} \sqrt{g} d^4 x \tag{15}$$

where

$$F_{\mu\nu} = \nabla_{\mu}A_{\nu} - \nabla_{\nu}A_{\mu} \tag{16}$$

Solution: So, we need to write the action in a way that exposes the dependence on the metric,

$$S = -\frac{1}{4} \int g^{\lambda\mu} g^{d\nu} F_{\lambda\rho} F_{\mu\nu} \sqrt{g} d^4 x \tag{17}$$

and then use

$$\begin{aligned}
\delta g_{\mu\nu} &= h_{\mu\nu} \\
\delta g^{\mu\nu} &= -h^{\mu\nu} \\
\delta \sqrt{g} &= \frac{1}{2}\sqrt{g}h
\end{aligned}$$
(18)

where, as usual, we are using the notation $h = g^{\mu\nu} h_{\mu\nu}$. Now,

$$S = \int \mathcal{L} d^4 x \tag{19}$$

so that

$$\delta S = -\int \left(-\frac{1}{2} h^{\lambda\mu} g^{\rho\nu} F_{\lambda\rho} F_{\mu\nu} + \frac{1}{8} g^{\lambda\mu} g^{\rho\nu} F_{\lambda\rho} F_{\mu\nu} h \right) \sqrt{g} d^4 x \tag{20}$$

$$= \int \left(\frac{1}{2}F_{\lambda\rho}F_{\mu}^{\ \rho} - \frac{1}{8}F^{\rho\nu}F_{\rho\nu}g_{\lambda\mu}\right)h^{\lambda\mu}\sqrt{g}d^{4}x \tag{21}$$

hence

$$T_{\lambda\mu} = 2 \frac{\delta S}{\delta h^{\lambda\mu}} = F_{\lambda\rho} F_{\mu}{}^{\rho} - \frac{1}{4} F^{\rho\nu} F_{\rho\nu} g_{\lambda\mu}$$
(22)