The Kinematics and Morphology of Solar Coronal Mass Ejections



Jason P. Byrne, B.A.(Mod.)
School of Physics
University of Dublin, Trinity College

A thesis submitted for the degree of PhilosophiæDoctor~(PhD) September 2010

Declaration

I, Jason P. Byrne, hereby certify that I am the sole author of this thesis and that all the work presented in it, unless otherwise referenced, is entirely my own. I also declare that this work has not been submitted, in whole or in part, to any other university or college for any degree or other qualification.

in whole or in part, to any other university or college for any degree or other qualification.
Name: Jason P. Byrne
Signature: Date:
The thesis work was conducted from September 2006 to August 2016 under the supervision of Dr. Peter T. Gallagher at Trinity College University of Dublin.
In submitting this thesis to the University of Dublin I agree that the University Library may lend or copy the thesis upon request.
Name: Jason P. Byrne
Signature: Date:

Abstract

Solar coronal mass ejections (CMEs) are large-scale eruptions of plasma and magnetic field from the Sun into the corona and interplanetary space. They are the most significant drivers of adverse space weather at Earth and other locations in the heliosphere, so it is important to understand the physics governing their eruption and propagation. However the diffuse morphology and transient nature of CMEs makes them difficult to identify and track using traditional image processing techniques. Furthermore, the true three-dimensional geometry of CMEs has remained elusive due to the limitations of coronagraph plane-of-sky images with restricted fields-of-view. For these reasons the Solar Terrestrial Relations Observatory (STEREO) was launched as a twin-spacecraft mission to fly in orbits ahead and behind the Earth in order to triangulate independent observations of CME structure. It is the first time CMEs have been observed from vantage points off the Sun-Earth line and each spacecraft carries an instrument suite designed to image from the low solar corona out to the orbit of Earth in order to observe and study CME propagation toward Earth, important for space weather.

In this thesis the implementation of multiscale image processing techniques to identify and track the CME front through coronagraph images is detailed. An ellipse characterisation of the CME front is used to determine the CME kinematics and morphology with increased precision as compared to techniques used in current CME catalogues, and efforts are underway to automate this procedure for applying to a large number of CME observations for future analysis. It was found that CMEs do not simply undergo constant acceleration, but rather

tend to show a higher acceleration early in their propagation. The angular width of CMEs was also found to change as they propagate, normally increasing with height from the Sun. However these results were derived from plane-of-sky measurements with no correction for how the true CME geometry and direction affect the kinematics and morphology observed.

With the advent of the unique dual perspectives of the STEREO spacecraft, the multiscale methods were extended to an elliptical tie-pointing technique in order reconstruct the front of a CME in three-dimensions. Applying this technique to the Earth-directed CME of 12 December 2008 allowed an accurate determination of its true kinematics and morhpology, and the CME was found to undergo early acceleration, non-radial motion, angular width expansion, and aero-dynamic drag in the solar wind as it propagated towards Earth. This study and its conclusions are of vital importance to the fields of space weather monitoring and forecasting.



${\bf Acknowledgements}$

 ${\bf Acknowledgements.}$

Contents

Li	ist of	Publi	cations		vii
$\mathbf{L}_{\mathbf{i}}$	ist of	Figur	es		ix
Li	ist of	Table	\mathbf{s}		x
1	Intr	oduct	ion		1
	1.1	The S	olar Inter	ior	2
	1.2	The S	olar Atm	osphere	6
		1.2.1	Photosp	bhere	6
		1.2.2	Chromo	sphere	8
		1.2.3	Corona		10
		1.2.4	Solar W	ind	13
	1.3	Coron	al Mass I	Ejections	18
		1.3.1	Magneto	ohydrodynamic Theory	20
			1.3.1.1	Maxwell's Equations	21
			1.3.1.2	Fluid Equations	21
			1.3.1.3	The Induction Equation	22
			1.3.1.4	The Magnetic Reynolds Number	23
			1.3.1.5	Magnetic Reconnection	25
		1.3.2	Theoret	ical CME Models	28
			1.3.2.1	Catastrophe Model	30
			1.3.2.2	Toroidal Instability	34
			1.3.2.3	Breakout Model	36
		1.3.3	CME O	bservations	38

CONTENTS

			1.3.3.1	Γhomson Scattering	38
			1.3.3.2	CME Kinematics & Morphology	42
	1.4	Outlin	e of Thesis		49
2	Inst	rumen	tation		51
	2.1	SOHC	/LASCO		52
	2.2	STER	EO/SECCI	HI	56
		2.2.1	EUVI .		58
		2.2.2	COR1 .		60
		2.2.3	COR2 .		61
		2.2.4	Heliospher	ric Imagers	61
	2.3	CCD :	Detectors		64
	2.4	Coord	inate Syste	ms	68
		2.4.1	Heliograpl	nic Coordinates	68
		2.4.2	Heliocentr	ic Coordinates	69
			2.4.2.1 I	Heliocentric Earth Equatorial (HEEQ)	70
			2.4.2.2 I	Heliocentric Earth Ecliptic (HEE)	70
			2.4.2.3 I	Heliocentric Aries Ecliptic (HAE)	70
		2.4.3	Helioproje	ective Coordinates	70
3	Det	ecting	and Tracl	king CMEs	72
	3.1	CME	Detection (Catalogues	73
		3.1.1	CDAW .		74
		3.1.2	CACTus		75
		3.1.3	SEEDS.		78
		3.1.4	ARTEMIS	8	80
	3.2	Multis	cale Filteri	ng	83
	3.3	Auton	nated Multi	scale CME Detection	88
		3.3.1	Normalisii	ng Radial Graded Filter	89
		3.3.2	Threshold	ing	91
		3.3.3		Es and Streamer Interactions/Deflections	94
	3.4	Chara	cterising th	e CME Front	97

4	The	Kiner	natics and Morphology of CMEs using Multiscale Met	th-
	ods			101
	4.1	Introd	luction	. 102
	4.2	Error	Analysis	. 104
		4.2.1	3-Point Lagrangian Interpolation	. 104
	4.3	Result	ts	. 108
		4.3.1	Arcade Eruption: 2 January 2000	. 108
		4.3.2	Gradual/Expanding CME: 18 April 2000	. 110
		4.3.3	Impulsive CME: 23 April 2000	. 112
		4.3.4	Faint CME: 23 April 2001	. 115
		4.3.5	Fast CME: 21 April 2002	. 115
		4.3.6	Flux-Rope/Slow CME: 1 April 2004	. 117
		4.3.7	STEREO-B Event: 8 October 2007	. 117
		4.3.8	STEREO-A Event: 16 November 2007	. 120
	4.4	Discus	ssion & Conclusions	. 120
5	Dro	nagati	on of an Earth-Directed CME in Three-Dimensions	126
J	5.1	- 0	luction	
	5.2		TEREO Era	
	5.3		ical Tie-Pointing	
	5.5	5.3.1	-	
	5.4		-Directed CME	
	5.5		ts	
	5.5	5.5.1	3D Error Propagation	
		5.5.2	Prominence & CME Acceleration	
			Non-radial Prominence & CME Motion	
		5.5.4	CME Angular Width Expansion	
		5.5.5	-	
		5.5.6	CME Drag in the Inner Heliosphere	
	5.6	5.5.7	CME 'Pancaking'	. 161

CONTENTS

6	Con	clusio	ns & Future Work	167
	6.1	Princip	pal Results	167
	6.2	Future	e Work	171
		6.2.1	Automation	172
		6.2.2	Ridgelets/Curvelets	175
		6.2.3	3D CME Reconstructions	178
		6.2.4	Deriving CME Kinematics	180
R	ofere	nces		183

List of Publications

Refereed

- Byrne, J. P., Maloney, S. A., McAteer, R. T. J., Refojo, J. M. & Gallagher, P. T.
 - "Propagation of an Earth-directed coronal mass ejection in three-dimensions" *Nature Communications*, 1:74 doi: 10.1038/ncomms1077 (2010).
- Byrne, J. P., Gallagher, P. T., McAteer, R. T. J. & Young, C. A. "The kinematics of coronal mass ejections using multiscale methods" Astronomy & Astrophysics, 495, 325–334 (2009).
- 3. Gallagher, P. T., Young, C. A., **Byrne**, **J. P.** & McAteer, R. T. J. "Coronal mass ejection detection using wavelets, curvelets and ridgelets: Applications for space weather monitoring" *Advances in Space Research*, in press.
- Pérez-Suárez, D., Higgins, P. A., Bloomfield, D. S., McAteer, R. T. J., Krista, L. D., Byrne, J. P. & Gallagher, P. T.
 "Automated Solar Feature Detection for Space Weather Applications" in Qahwaji, R., Green, R. & Hines, E. [Eds] Applied Signal and Image Processing: Multidisciplinary Advancements, IGI, in press.
- Mierla, M., Inhester, B., Antunes, A., Boursier, Y., Byrne, J. P., Colaninno, R., Davila, J., de Koning, C.A., Gallagher, P.T., Gissot, S., Howard, R.A., Howard, T.A., Kramar, M., Lamy, P., Liewer, P.C., Maloney, S., Marqué, C., McAteer, R.T.J., Moran, T., Rodriguez, L., Srivastava, N., St. Cyr, O.C., Stenborg, G., Temmer, M., Thernisien, A., Vourlidas, A., West, M.J.,

Wood, B.E. & Zhukov, A.N.

"On the 3-D reconstruction of coronal mass ejections using coronagraph data" $\,$

Annales Geophysicae, 28, 203–215 (2010).

Contributed

1. Byrne, J. P., Young, C. A., Gallagher, P. T. & McAteer, R. T. J.

"Multiscale Characterization of Eruptive Events"

In S. A. Matthews, J. M. Davis & L. K. Harra, eds., First Results from Hinode, vol. 397 of Astronomical Society of the Pacific Conference Series, 162–163 (2008).

List of Figures

1.1	Illustration of the structure of the Sun. The core is the source of	
	energy, where fusion heats the plasma to ~ 15 MK. Energy is trans-	
	ported from the core by radiative processes in the radiation zone.	
	The convection zone is heated from the base at the tachocline, al-	
	lowing convective currents to flow to the photosphere. Locations	
	of strong magnetic fields inhibit convection and appear as dark	
	sunspots on the photosphere. These strong magnetic fields ex-	
	tend into the upper atmosphere of the Sun, responsible for coronal	
	loops, prominences and streamers. Image credit: eu.spaceref.com.	;
1.2	Illustration of the $\alpha\Omega$ effect of winding-up magnetic field due to the	
	differential rotation of the Sun, reproduced from Babcock (1961).	
	Sunspots visible on the disk are as a result of protruding field with	
	positive p and negative f polarity as shown	Ę
1.3	A 1D static model of electron density N_e [cm ⁻³] and temperature	
	T_e [K] profiles in the solar atmosphere, reproduced from Gabriel &	
	Mason (1982). In the chromosphere, the plasma is only partially	
	ionized. The plasma becomes fully ionized at the sharp transition	
	from chromospheric to coronal temperatures	7
1.4	Plasma β as a function of height for a regime of magnetic field	
	strengths between ~ 100 and $\sim 2{,}500$ G, reproduced from Gary	
	(2001). The dotted lines segregate the layers of photosphere (β >	
	1), chromosphere and corona ($\beta < 1$), and the solar wind ($\beta > 1$).	8
1.5	Soft X-ray image of the solar corona recorded by Yohkoh on 26	
	August 1992, reproduced from Aschwanden (2005)	10

1.7	Schematic of the Parker Spiral in the heliosphere. The streamlines	
	of the solar wind act to drag out the magnetic field lines of the	
	Sun, which become wound up in an Archimedean spiral as a result	
	of the Sun's rotation. Image credit: Steve Suess, $NASA/MSFC$	16
1.8	Observation of a CME from the LASCO/C2 coronagraph on board	
	the SOHO spacecraft. The field-of-view extends from the inner	
	mask at $\sim 2~R_{\odot}$ to the edges at $\sim 6~R_{\odot}$. The white circle repre-	
	sents the relative size and location of the Sun. The complexity of	
	the magnetic field driving the eruption is clearly indicated by the	
	twisted geometry of the bright ejecta	19
1.9	Geometry of the Sweet-Parker (top) and Petschek (bottom) recon-	
	nection models, reproduced from Aschwanden (2005)	26
1.10	Illustrations of the different mechanical analogues of CME erup-	
	tions, reproduced from Klimchuk (2001)	29
1.11	Theoretical evolution of a 2D flux rope, reproduced from Forbes	
	& Priest (1995). The flux rope foot-point separation λ decreases,	
	increasing the magnetic pressure until the flux rope becomes unsta-	
	ble and erupts away from the surface $(b-d)$. The resulting height	
	evolution of the flux rope is illustrated in (a)	31
1.12	A schematic of the 3D flux rope model, reproduced from Chen $\&$	
	Krall (2003), an extension of the 2D flux rope in Figure 1.11 when	
	viewed end-on as indicated by the arrow from the right. The flux	
	rope is rooted below the photosphere, and surrounded by the ambi-	
	ent coronal magnetic field $\mathbf{B_c}$ and plasma density ρ_c . Components	
	of the current density ${f J}$ and magnetic field ${f B}$ are shown, where	
	subscripts 't' and 'p' refer to the toroidal and poloidal directions	
	respectively. The flux rope has a radius of curvature R , radius of	
	cross-section a , apex height Z , footpoint separation s_f , and the	
	radial force outward is F_R	33

1.13	Schematic, and corresponding simulation snapshots, of the main	
	stages of the axisymmetric 2.5D breakout model, reproduced from	
	Lynch et al. (2008). (a) shows the initial multipolar topology of	
	the system, (b) shows the shearing phase which distorts the x-line	
	and causes breakout reconnection to begin, (c) shows the onset	
	of flare reconnection behind the eruption that disconnects the flux	
	rope, and (d) shows the system restoring itself following the eruption.	37
1.14	The Thomson scattering geometry for a single electron, reproduced	
	from Howard & Tappin (2009). a) shows the electron and incident	
	light with different observer positions indicated. \mathbf{b}), \mathbf{c}), \mathbf{d}) show	
	the resultant scattering of light seen by each observer O_1, O_2, O_3	
	respectively	39
1.15	Schematic of the Thomson surface, being the sphere of all points	
	which are located at an angle of 90° between the Sun and the	
	observer, reproduced from Vourlidas & Howard (2006). An exam-	
	ple line-of-sight is shown for an electron at point P , with radial	
	distance r from the Sun, at longitude ϕ relative to the solar limb.	41
1.16		44
1.17	(a) Height-time, (b) velocity and (c) acceleration profiles for the	
	CME on 21 April 2002, and (d) the GOES-10 soft X-ray flux for	
	the associated X1.5 flare during the interval 00:47-03:20 UT, re-	
	produced from Gallagher et al. (2003). A three-point difference	
	scheme is used on the data points, and a first-difference scheme	
	is plotted with filled circles. The solid line is the best fit of the	
	exponentially increasing and decreasing acceleration of equation 1.53.	45

1.18	The kinematics of the CME leading edge ($circles$), cavity ($pluses$)	
	and prominence (triangles) of the 15 May 2001 event, reproduced	
	from Maričić et al. (2004). (a) shows the height-time profiles. (b)	
	shows the distance between the leading edge and the prominence	
	(thick line), the cavity and the prominence (thin line), and the	
	leading edge and the cavity (dashed line). (c) shows the veloci-	
	ties determined by forward-difference technique upon the smoothed	
	height-times. (d) shows the onset of acceleration against height,	
	with a straight line fit. The horizontal bar between (a) and (c)	
	indicates the period of the soft X-ray burst ($dotted$ - precursor, $full$	
	- main rise)	47
2.1	A schematic of the SOHO spacecraft and onboard instrument suites,	
	reproduced from Domingo et al. (1995)	53
2.2	The externally occulted Lyot coronagraph, reproduced from Brueck-	
	ner $et\ al.\ (1995),$ showing: front aperture A0 and external occulter	
	D1; entrance aperture A1 and objective lens O1; the field stop;	
	inner occulter D2 and field lens O2; Lyot stop A3 and relay lens	
	O3 with Lyot spot; filter and polariser wheels F/P; and the focal	
	plane F	54
2.3	A LASCO/C3 image of the solar corona out to $\sim\!30~R_{\odot}.$	55
2.4	Payload diagram of one of the STEREO spacecraft, indicating the	
	positions of the four instrument suites onboard: Sun-Earth Con-	
	nection Coronal and Heliospheric Imagers (SECCHI); In-situ Mea-	
	surements of Particles and CME Transients (IMPACT); Plasma	
	and SupraThermal Ion Composition (PLASTIC); STEREO/WAVES	
	radio burst tracker (SWAVES). Image credit: stereo.gsfc.nasa.gov.	57
2.5	Diagram of the EUVI in the STEREO/SECCHI suite, reproduced	
	from Howard <i>et al.</i> (2000)	58
2.6	${\it Diagram\ of\ the\ COR1\ coronagraph\ in\ the\ STEREO/SECCHI\ suite},$	
	reproduced from Howard $et\ al.\ (2000).$ The measurements are	
	specified in millimetres	59

2.7	Diagram of the COR2 coronagraph in the STEREO/SECCHI suite,	
	reproduced from Howard et al. (2000). The measurements are	
	specified in millimetres	62
2.8	Diagram of the heliospheric imagers HI-1/2 in the STEREO/SECCHI	
	suite, reproduced from Howard et al. (2000)	63
2.9	The intensity profile of a CME compared to the K & F coronae observed at elongations up to 90°, and the corresponding fields-of-	
	view of the Heliospheric Imagers (HI1/2), reproduced from Howard	
	et al. (2000)	64
2.10	Illustration of a thick front-side illuminated CCD. Image credit:	
	www.ing.iac.es.	65
2.11	Illustration of a thinned back-side illuminated CCD. Image credit:	
	www.ing.iac.es.	66
2.12	Schematics of two Sun-centred coordinate systems, reproduced	
	from Thompson (2006). (a) Stonyhurst heliographic coordinates	
	commonly used to locate features on-disk. (b) Heliocentric-cartesian	
	coordinates commonly used for spatially localising features in the	
0.10	v	69
2.13	Schematic of the celestial sphere and ecliptic plane. The celestial	
	equator is a projection of the Earth's equator, and the ecliptic is	 -1
	a projection of the Earth's orbit about the Sun	71
3.1	Raw (left) and pre-processed image (right) of a CME observed by	
	LASCO/C2 on 1 April 2004. The pre-processing includes normal-	
	ising the image statistics, taking a background subtraction, and	
	masking the occulter disk. The white circle (right) indicates the	
	relative size and position of the Sun behind the occulter	74
3.2	The top image (a) shows the detection of ridges in the (t, r) stacks	
	of the CACTus catalogue, through the use of the Hough transform	
	detailed in the bottom image (b), reproduced from Robbrecht &	
	Berghmans (2004)	76

3.3	Example of the SEEDS CME detection and height determination,	
	reproduced from Olmedo et al. (2008). (a) shows the running-	
	difference image unwrapped into Sun-centre polar coordinates, show-	
	ing a CME observed by LASCO/C2 on 12 September 2002. The	
	black line distribution across the image represents the positive	
	value intensity count along each angle, and the two vertical black	
	lines mark the angular span at one standard deviation above the	
	mean intensity. (b) shows the new angular span following the re-	
	gion growing technique. (c) shows the intensity within the angular	
	span averaged across heights, and the 'Half-Max-Lead' is taken as	
	the CME height in the image	79
3.4	An example of how the synoptic maps are generated for the ARTEMIS	
	catalogue, reproduced from Boursier et al. (2009b). At a chosen	
	height in the coronagraph image an annulus is unwrapped (indi-	
	cated with the dashed line and blue square, circle and triangle)	
	and these are then stacked together to illustrate how the intensity	
	at that height changes through time. Vertical streaks represent	
	transient events occurring on smaller time-scales than the more	
	persistant streamers in the images	82
3.5	Top left, the horizontal detail, and top right, the vertical detail	
	from the high-pass filtering at one scale of the multiscale decom-	
	position (called the rows and columns respectively). Bottom left,	
	the corresponding magnitude (edge strength) and bottom right,	
	the angle information $(0-360^{\circ})$ taken from the gradient space, for	
	a CME observed in LASCO/C2 on 1 April 2004 (Byrne $\it et~\it al.,$ 2009).	86
3.6	The vectors plotted represent the magnitude and angle determined	
	from the gradient space of the high-pass filtering at scale 3. The	
	CME of 2004 April 1 shown here is highlighted very effectively by	
	this method (Byrne $et~al., 2009$)	87
3.7	A normalised, background subtracted, LASCO/C2 image (left) of a	
	CME on 1 April 2004, and the resulting NRGF image (right). The	
	image radial intensity is scaled such that structure along streamers	
	and the CME becomes visible across the field-of-view	QΩ

3.8	A chosen scale of the decomposed NRGF image provides a mag-	
	nitude image of the edge strengths displayed on the left, which	
	is thresholded at one standard deviation of the mean intensity to	
	obtain contoured regions of interest that could contain a CME	
	(sample contours indicated in red). As shown, the streamers have	
	edges which appear on the same scale as the CME edges in this	
	image. The angular information from the decomposition is dis-	
	played on the right, and the contoured regions of interest overlaid	
	for comparison. The grey scale indicates angles from $0-360^{\circ}$ and	
	it is clear that streamers tend to have a linear colour scale while	
	the CME has a gradient of colours across the scale	91
3.9	Left: four contoured regions (at one standard deviation of the	
	mean image intensity) highlighted on the magnitude information	
	from the multiscale decomposition of the 1 April 2004 CME. Right:	
	the corresponding angular distribution of each region, normalised	
	and folded into the $0-180^\circ$ range (centred on 90°). The angular	
	distribution may be thresholded with respect to its median value	
	to distinguish regions corresponding to CMEs from those along	
	streamers	92
3.10	The resulting CME detection mask from combining the thresh-	
	olded regions of strongest magnitude and angular distribution at	
	four scales of the decomposition	93
3.11	The NRGF (left) and resulting detection masks (middle and right)	
	for different frames of a CME observed by LASCO/C2 at times	
	of 00:00 UT (top), 00:40 UT (middle) and 01:20 UT (bottom)	
	on 2 April 2004. The location of the CME front is highlighted	
	very efficiently by this method, although the detection masks may	
	contain artefacts of the chosen thresholds which must be discarded	
	when characterising the CME front	95
3.12	Example of the difficulty in detecting the faint CME observed by	
	LASCO on 23 April 2001. If the CME is far-sided and/or of low	
	intensity it becomes difficult to threshold its edges in the image	
	compared to the coronal streamers	96

3.13	Ellipse inclined at angle γ , with semimajor axis a , semiminor axis	
	$b,$ and radial line ρ inclined at angle ω to the semimajor axis	98
3.14	A depiction of three frames sampled from the ellipse characterisa-	
	tion of a CME observed by LASCO/C2 on 24 January 2007. The	
	white dots along the CME front indicate the multiscale edge de-	
	tection to which the ellipse is fit. The black cross on the ellipse	
	indicates the furthest point measured from Sun centre which pro-	
	vides the height-time profile for determining the kinematics. The	
	opening angle from Sun centre measures the CME width	100
4.1	A sample of ellipse fits to the multiscale edge detection of the	
	events studied. For each event the upper and lower image show	
	LASCO/C2 and C3, except for the 2007 events which show SEC-	
	CHI/COR1 and COR2	107
4.2	Kinematic and morphological profiles for the ellipse fit to the mul-	
	tiscale edge detection of the 2 January 2000 CME observed by	
	LASCO/C2 and C3. The plots from top to bottom are height,	
	velocity, angular width, and ellipse tilt. The CDAW heights are	
	over-plotted with a dashed line. The height and velocity fits are	
	based upon the constant acceleration model	111
4.3	Kinematic and morphological profiles for the ellipse fit to the multi-	
	scale edge detection of the 18 April 2000 CME observed by LASCO/C	22
	and C3. The plots from top to bottom are height, velocity, angular	
	width, and ellipse tilt. The CDAW heights are over-plotted with	
	a dashed line. The height and velocity fits are based upon the	
	constant acceleration model	113
4.4	Kinematic and morphological profiles for the ellipse fit to the multi-	
	scale edge detection of the 23 April 2000 CME observed by LASCO/O	22
	and C3. The plots from top to bottom are height, velocity, angular	
	width, and ellipse tilt. The CDAW heights are over-plotted with	
	a dashed line. The height and velocity fits are based upon the	
	constant acceleration model	114

4.5	Kinematic and morphological profiles for the ellipse fit to the multi-	
	scale edge detection of the 23 April 2001 CME observed by LASCO/C2	
	and C3. The plots from top to bottom are height, velocity, angular	
	width, and ellipse tilt. The CDAW heights are over-plotted with	
	a dashed line. The height and velocity fits are based upon the	
	constant acceleration model	16
4.6	Kinematic and morphological profiles for the ellipse fit to the multi-	
	scale edge detection of the 21 April 2002 CME observed by LASCO/C2	
	and C3. The plots from top to bottom are height, velocity, angular	
	width, and ellipse tilt. The CDAW heights are over-plotted with	
	a dashed line. The height and velocity fits are based upon the	
	constant acceleration model	18
4.7	Kinematic and morphological profiles for the ellipse fit to the multi-	
	scale edge detection of the 1 April 2004 CME observed by LASCO/C2 $$	
	and C3. The plots from top to bottom are height, velocity, angular	
	width, and ellipse tilt. The CDAW heights are over-plotted with	
	a dashed line. The height and velocity fits are based upon the	
	constant acceleration model	19
4.8	Kinematic and morphological profiles for the ellipse fit to the mul-	
	tiscale edge detection of the 8 October 2007 CME observed by	
	SECCHI/COR1 and COR2. The plots from top to bottom are	
	height, velocity, angular width, and ellipse tilt. The height and	
	velocity fits are based upon the constant acceleration model 1	21
4.9	Kinematic and morphological profiles for the ellipse fit to the mul-	
	tiscale edge detection of the 16 November 2007 CME observed by	
	SECCHI/COR1 and COR2. The plots from top to bottom are	
	height, velocity, angular width, and ellipse tilt. The height and	
	velocity fits are based upon the constant acceleration model 1	22
5.1	Schematic of the epipolar geometry used to relate the observa-	
J.1	tions from the two STEREO spacecraft, reproduced from Inhester	
	(2006). This geometry enables us to localise features in 3D space	
		29

5.2	Schematic of how tie-pointing a curved surface within an epipolar
	geometry is limited in its ability to resolve the true feature, since
	lines-of-sight will be tangent to different edges of the surface and
	not necessarily intersect upon it. Reproduced from Inhester (2006). 131
5.3	Schematic of the Graduated Cylindrical Shell (GCS) model of a
	flux rope CME, reproduced from Thernisien et al. (2009). Indi-
	cated are the model parameters of front height \mathbf{h}_{front} , leg height
	h , angle between the legs 2α , cross-sectional radius a , and distance
	from Sun centre ${\bf O}$ to a point on the edge of the shell ${\bf r}$. Two views
	of the GCS model are shown; (a) 'face-on', and (b) 'end-on'. The
	positional parameters of longitude ϕ and latitude θ are illustrated
	in (c)
5.4	The elliptical tie-pointing technique developed to reconstruct the
	3D CME front, shown here for the 12 December 2008 event. One of
	any number of epipolar planes will intersect the ellipse characteri-
	sation of the CME at two points in each image from STEREO-A
	and B. (a) illustrates how the resulting four sight-lines intersect in
	3D space to define a quadrilateral that constrains the CME front
	in that plane. Inscribing an ellipse within the quadrilateral such
	that it is tangent to each sight-line provides a slice through the
	CME that matches the observations from each spacecraft. (\mathbf{b}) il-
	lustrates how a full reconstruction is achieved by stacking multiple
	ellipses from the epipolar slices to create a model CME front that
	is an optimum reconstruction of the true CME front. (c) illustrates
	how this is repeated for every frame of the eruption to build the
	reconstruction as a function of time and view the changes to the
	CME front as it propagates in 3D. While the ellipse characterisa-
	tion applies to both the leading edges and, when observable, the
	flanks of the CME, only the outermost part of the reconstructed
	front is shown here for clarity

5.5	An ellipse inscribed within a convex quadrilateral. An isometry of	
	the plane is applied such that the quadrilateral has vertices $(0,0)$,	
	(A,B),(0,C),(s,t). The ellipse has center $(h,k),$ semimajor axis	
	a , semiminor axis b , tilt angle δ , and is tangent to each side of the	
	quadrilateral	135
5.6	The back-projection of a STEREO 3D CME front reconstruction	
	onto the SOHO/LASCO plane-of-sky, from the observations by	
	STEREO-A (red) and STEREO-B (green) at 16:22 UT on 26 April	
	2008	138
5.7	The back-projection of a STEREO 3D CME front reconstruction	
	onto the SOHO/LASCO plane-of-sky, at 16:22 UT on 26 April	
	2008. The left panel shows the multiscale edge detection and ellipse	
	characterisation of the CME front observed off the east limb from	
	STEREO-A. The right panel shows the running-difference point-	
	and-click ellipse characterisation of the CME front observed as	
	a halo event from STEREO-B The middle panel shows the	
	LASCO/C2 observation of the CME at 16:30 UT (8 minutes later	
	than the STEREO observations) with the back-projected 3D CME $$	
	front reconstruction overplotted for comparison	140
5.8	Composite of STEREO-Ahead and Behind images from EUVI,	
	COR1, COR2, and HI1 (Byrne et al., 2010). (a) indicates the	
	STEREO spacecraft locations, separated by an angle of 86.7° at	
	the time of the event. (\mathbf{b}) shows the prominence eruption observed	
	in EUVI-B off the NW limb from approximately $03:00$ UT which	
	is considered to be the inner material of the CME. The multi-	
	scale edge detection and corresponding ellipse characterisation are	
	overplotted in COR1. (c) shows that the CME is Earth-directed,	
	being observed off the east limb in STEREO-A and the west limb	
	in STEREO-B	141

5.9	STEREO-Ahead and Behind COR1 images at 07:35 UT on the 12	
	December 2008. Overplotted are: the multiscale edge detections of	
	the CME front (red); the ellipse characterisations (blue); and the	
	resulting 3D reconstructions back-projected onto the plane-of-sky	
	(white)	142
5.10	STEREO-Ahead and Behind COR2 images at 14:52 UT on the 12	
	December 2008. Overplotted are: the multiscale edge detections of	
	the CME front (red); the ellipse characterisations (blue); and the	
	resulting 3D reconstructions back-projected onto the plane-of-sky	
	(white)	143
5.11	STEREO-Ahead and Behind HI1 images at 01:29 UT on the 13	
	December 2008. Overplotted are: the running difference edge	
	detections of the CME front (red); the ellipse characterisations	
	(blue); and the resulting 3D reconstructions back-projected onto	
	the plane-of-sky (white)	144
5.12	2 The 3D CME front reconstruction as it propagates along the Sun-	
	Earth line into interplanetary space. This particular frame of the	
	reconstruction is from the observations of COR1 at $07:35$ UT on	
	12 December 2008	145
5.13	3 The 3D CME front reconstruction as it propagates along the Sun-	
	Earth line into interplanetary space. This particular frame of the	
	reconstruction is from the observations of COR2 at 14:52 UT on	
	12 December 2008	146
5.14	The 3D CME front reconstruction as it propagates along the Sun-	
	Earth line into interplanetary space. This particular frame of the	
	reconstruction is from the observations of HI1 at $01:29$ UT on 13	
	December 2008	147

5.15	The 3D CME front reconstruction from the COR2 Ahead and Be-	
	hind frames at 14:52 UT on 12 December 2008. The lines drawn	
	from Sun-centre indicate the 'Midpoint of Front' (solid blue), the	
	'Northern/Southern Flanks' (solid red/brown), and the 'Midtop/Midl	bottom
	of Front' at angles in between (dashed red/brown). By taking these	
	measurements across all frames we may determine the kinematics	
	and morphology of the CME as plotted in Figures 5.17 and 5.18	149
5.16	The error trapezoid on the tie-pointing of two lines-of-sight in 3D	
	space, reproduced from Inhester (2006). The error w in localising	
	a point on each plane-of-sky results in a trapezoid with a diagonal	
	measuring $w/\sin(\alpha/2)$ as shown	151
5.17	The kinematics of the prominence and 3D reconstructed CME	
	front of 12 December 2008. The prominence is observed as the in-	
	ner material of the CME, with both undergoing acceleration from	
	$\sim\!06:\!00-07:\!00$ UT, peaking at $\sim\!40~\mathrm{m~s^{-2}}$ and $\sim\!94~\mathrm{m~s^{-2}}$ re-	
	spectively, before reducing to scatter about zero. Measurement	
	uncertainties are indicated by one standard deviation error bars	153
5.18	Kinematic and morphological properties of the 3D reconstruction	
	of the 12 December 2008 CME front. The top panel shows the	
	velocity of the middle of the CME front with corresponding drag	
	model and, inset, the early acceleration peak. Measurement uncer-	
	tainties are indicated by one standard deviation error-bars. The	
	middle panel shows the declinations from the ecliptic (0°) of an	
	angular spread across the front between the CME flanks with a	
	power-law fit indicative of non-radial propagation. The bottom	
	panel shows the angular width of the CME with a power-law ex-	
	pansion.	154

5.19	The 3D CME front parameters are used as initial conditions for an	
	ENLIL with Cone Model MHD simulation (Xie et al., 2004) and	
	the output density (top) and velocity (bottom) profiles of the inner	
	heliosphere are illustrated here for the time-stamp of $06:00~\mathrm{UT}$	
	on 14 December 2008. Beyond distances of $\sim\!50~R_{\odot}$ the CME is	
	slowed by its interaction with the upstream, slow-speed, solar wind	
	flow along its trajectory towards Earth, and this accounts for its	
	arrival time as detected in-situ by the ACE and WIND spacecraft	
	at the L1 point near Earth	159
5.20	The in-situ solar wind plasma and magnetic field data observed	
	by the WIND spacecraft. From top to bottom the panels show	
	proton density, bulk flow speed, proton temperature, and magnetic	
	field strength and components. The red dashed lines indicate the	
	arrival time of the density enhancement predicted from our ENLIL	
	with Cone Model run providing 08:09 UT on 16 December 2008,	
	with a potential offset error between our reconstruction and the	
	derived model height profiles accounting for an arrival time up to	
	13:20 UT. We observe a magnetic cloud signature behind the front,	
	as highlighted with blue dash-dotted lines	160
5.21	Top: Ellipses characterising the 3D CME front reconstruction in	
	the out-of-ecliptic plane along the Sun-Earth line. Bottom: The	
	ellipse tilt angle is indicative of the initial effects of 'pancaking', as	
	the curvature of the CME front decreases with increasing height	
	due to its changing morphology in the solar wind	163
6.2	An example interface which could be developed for the poten-	
	tially automated multiscale filtering and ellipse characterisation	
	for tracking CMEs observed by STEREO-A and B. Images from	
	the respective instruments appear on the left, and the resulting	
	parameters from the characterisation appear on the right	174

6.3	An example ridgelet, reproduced from Gallagher $\operatorname{\it et\ al.}$ (2010). The	
	first graph shows a typical ridgelet, and the second to fourth graphs	
	are obtained from simple geometrical manipulations, namely rota-	
	tion, rescaling and shifting	176
6.4	The curvelet filtered image of a CME observed by LASCO/C2 on	
	18 April 2000, reproduced from Gallagher $\operatorname{\it et}$ al. (2010). The detail	
	along the curved CME front is enhanced as a result of the curvelet	
	technique following the removal of certain coefficients probably due	
	to noise	177
6.6	A theoretical model for a CME with constant acceleration 2 m $\rm s^{-2}$	
	and initial velocity $300~\rm km~s^{-1},$ and two simulations of how the	
	resulting profiles for a noisy sample of datapoints behave using	
	3-point Lagrangian interpolation	180
6.7	The resulting velocity profile for the 3D reconstructed CME front	
	of the 12 December 2008 using the inversion technique of Kontar	
	& MacKinnon (2005)	182

List of Tables

2.1	Summary of EUVI wavelengths	60
4.1	Summary of CME velocities as measured by CACTus, CDAW,	
	SEEDS and multiscale methods. Asterisks indicate analysis of	
	SECCHI data rather than LASCO	109
4.2	Summary of CME accelerations as measured by CACTus, CDAW,	
	SEEDS and multiscale methods. Asterisks indicate analysis of	
	SECCHI data rather than LASCO	109
4.3	Summary of CME angular widths as measured by CACTus, CDAW,	
	SEEDS and multiscale methods. Asterisks indicate analysis of	
	SECCHI data rather than LASCO	110

Chapter 1

Introduction

The Sun, as provider of light and heat to all life on Earth, has been a constant source of mystery and wonder to humankind. History recounts numerous tales inspired by our connection with the Sun: from its worship as a deity in the earliest civilisations, to the appreciation of its seasonal influence marked by structures like Newgrange, and the eventual observance of its complex behaviour with the development of telescopes and scientific intrigue. As our nearest star, astronomers have increasingly taken interest in the complexities of the Sun, and now in the modern age of space exploration numerous observatories have been built specifically to monitor solar activity and further our understanding of its dynamic behaviour.

1.1 The Solar Interior

The Sun is a G2V main sequence star of luminosity $L_{\odot}=3.85\times 10^{26}~\mathrm{J~s^{-1}}$, mass $M_{\odot}=1.99\times 10^{30}~\mathrm{kg}$ and radius $R_{\odot}=6.96\times 10^8~\mathrm{m}$ (Prialnik, 2000). It was born from the gravitational collapse of a molecular cloud approximately 4.57 billion years ago, is currently in a state of hydrostatic equilibrium ($\nabla P=-\rho g$), and predicted to enter a red giant phase in another ~ 5 billion years before ending its life as a white dwarf (Phillips, 1999). Since we cannot directly observe the interior of the Sun, its structure and evolution are fundamentally realised with the use of the 'standard solar model' (SSM; Bahcall, 1989) which is a mathematical treatment of stellar structure described by several differential equations derived from basic physical principles. The SSM is constrained by the well-determined boundary conditions of the Sun's luminosity, radius, age and composition, and thus provides a basis for understanding the mechanisms of energy transport in the solar interior.

The Sun may be described as a large nuclear reactor. The fundamental energy process driving it is nuclear fusion in the core through the proton-proton chain at temperatures of ~ 15 MK:

$$e^{-} + {}^{1}H + {}^{1}H \rightarrow {}^{2}H + \nu + 1.44 \text{ MeV}$$
 (1.1)

$$^{2}\text{H} + ^{1}\text{H} \rightarrow ^{3}\text{He} + \gamma + 5.49 \text{ MeV}$$
 (1.2)

$${}^{3}\text{He} + {}^{3}\text{He} \rightarrow {}^{4}\text{He} + {}^{1}\text{H} + {}^{1}\text{H} + 12.86 \text{ MeV},$$
 (1.3)

where $^1{\rm H}$ is a proton, $^3{\rm He}$ and $^4{\rm He}$ are helium isotopes, e⁻ an electron, ν a neutrino and γ a gamma ray. The resulting energy release is approximately $4.3 \times 10^{-12} \ {\rm J}$.




Figure 1.1: Illustration of the structure of the Sun. The core is the source of energy, where fusion heats the plasma to ~ 15 MK. Energy is transported from the core by radiative processes in the radiation zone. The convection zone is heated from the base at the tachocline, allowing convective currents to flow to the photosphere. Locations of strong magnetic fields inhibit convection and appear as dark sunspots on the photosphere. These strong magnetic fields extend into the upper atmosphere of the Sun, responsible for coronal loops, prominences and streamers. Image credit: eu.spaceref.com.

The core extends from the centre out to $\sim 0.25~R_{\odot}$, followed by the radiation zone out to $\sim 0.75~R_{\odot}$, then the convection zone out to the solar surface at 1 R_{\odot} (Figure 1.1). The temperature across the radiation zone drops to $\sim 5~MK$ with radiation being the most efficient method of energy transport. This radiation field is closely approximated by black body theory, for which the spectral radiance at all wavelengths may be described by the Planck equation:

$$B_{\lambda}(T) = \frac{2hc^2\mu^2}{\lambda^5 \left[\exp\left(hc/\lambda kT\right) - 1\right]}$$
(1.4)

where $B_{\lambda}(T)$ is the intensity of radiation per unit wavelength interval (at temperature T), h is the Plank constant, c is the speed of light, μ is the refractive index of the medium, and k is the Boltzmann constant. By Wien's law $\lambda_{max}T=2.8979\times 10^{-3}$ m K we determine that the radiation is in the form of X-rays, and these high-energy photons undergo random walks in the optically thin plasma, escaping into the convection zone on time-scales of 10^6 years. The optically thick convection zone then transports energy by fluid motion across the temperature gradient between its base ($\sim 1-2$ MK) and the solar surface ($\sim 5,800$ K). Plasma elements move sufficiently rapidly for the energy interchange with their surroundings to be negligible, i.e., they change adiabatically. A useful measure of when convection is likely to occur is given by the Schwarzschild criterion:

$$\left. \frac{d \log T}{d \log P} \right|_{star} > \frac{\gamma - 1}{\gamma} \tag{1.5}$$

where $\gamma = C_P/C_V$ is the ratio of specific heats, equal to 5/3 for a perfect monatomic gas. The rising and falling parcels of plasma create the granulation effects observed on the surface, with granules ranging in size from hundreds




Figure 1.2: Illustration of the $\alpha\Omega$ effect of winding-up magnetic field due to the differential rotation of the Sun, reproduced from Babcock (1961). Sunspots visible on the disk are as a result of protruding field with positive p and negative f polarity as shown.

to thousands of kilometres and dissipating over tens of minutes. (Details on the above radiative and convective processes are found in, e.g., Kitchin (1987)).

Between the radiation and convection zones is a relatively thin interface called the tachocline, where the solid body rotation of the radiative interior meets the differentially rotating outer convection zone. It thus has a very large shear profile which could account for the formation of large scale magnetic fields in the solar dynamo. The differential rotation of the Sun's convection zone causes a large-scale winding up of the magnetic field within, named the Ω -effect, while the effects of the coriolis force and smaller scale motions of the plasma can give twist and writhe to the field, named the α -effect (Figure 1.2). The Sun's magnetic field is observed to undergo a 22-year periodicity, with the magnetic poles flipping every 11 years to produce the solar cycle, giving rise to periods of increased and decreased solar activity manifested by the frequency of phenomena such as active regions, flares and transients in the solar atmosphere (Schrijver & Zwaan, 2000).

1.2 The Solar Atmosphere

The Sun's atmosphere is composed of all regions extending from the surface of the convection zone out into the heliosphere. It can be separated into distinct regimes dependent on the density and temperature profiles. These are plotted in Figure 1.3 for a 1D static model of the solar atmosphere. The layers are stratified into photosphere, chromosphere, transition region, and corona; having a decreasing density with increasing height from the photosphere, but from the chromosphere up the temperature increases with a dramatic jump in the transition region giving rise to the so-called 'coronal heating problem'. The interplay between the magnetic and gas pressure represents an important determining factor in the behaviour of structures throughout the solar atmosphere, quantified by the plasma- β term:

$$\beta = \frac{p_{gas}}{p_{mag}} = \frac{nkT}{(B^2/8\pi)} \tag{1.6}$$

This is illustrated in Figure 1.4 for the different layers of the atmosphere. At the photospheric level the plasma- β is large, and plasma motions dominate the over the magnetic field forces. Through the chromosphere and corona the plasma- β decreases to low values where the magnetic field structures are seen to suspend plasma in loops and filaments. Finally in the extended upper atmosphere the plasma- β rises again, and the magnetic field is advected out with the solar wind plasma flow to ultimately form the Parker spiral.

1.2.1 Photosphere

The surface of the Sun is the photosphere defined as the point where the optical depth equals 2/3 for wavelengths of visible light, centred on 5,000 Å ($\tau_{5000} \sim 2/3$




Figure 1.3: A 1D static model of electron density N_e [cm⁻³] and temperature T_e [K] profiles in the solar atmosphere, reproduced from Gabriel & Mason (1982). In the chromosphere, the plasma is only partially ionized. The plasma becomes fully ionized at the sharp transition from chromospheric to coronal temperatures.

for $I/I_0 = e^{-\tau}$). The spectrum of light emitted has a profile like that of a black body with an effective temperature of 5,800 K interspersed with the Fraunhofer absorption lines due to the tenuous layers above the photosphere. It has a particle density of $\sim 10^{23}$ m⁻³ and a thickness of less than 500 km. Cooler regions called sunspots have temperatures of 4,000–4,500 K and are due to intense magnetic field activity that acts to suppress convective plasma motion. Granulation of the photosphere is observed as the manifestation of plasma motion in the convection zone below, with typical cell sizes on the order of 1,000 km in diameter. They occur when hot plasma rises to the surface and is transported along it to the granule edges, which appear darker as the plasma cools and descends. The gas pressure dominates the magnetic pressure ($\beta > 1$), and the magnetic field is




Figure 1.4: Plasma β as a function of height for a regime of magnetic field strengths between ~ 100 and $\sim 2,500$ G, reproduced from Gary (2001). The dotted lines segregate the layers of photosphere ($\beta > 1$), chromosphere and corona ($\beta < 1$), and the solar wind ($\beta > 1$).

effectively coupled to the plasma motion which sweeps it into the inter-granular network.

1.2.2 Chromosphere

Above the photosphere lies the chromosphere where the temperature initially drops to a minimum of $\sim 4,500$ K before increasing to $\sim 20,000$ K with increasing height from the Sun (Figure 1.3). It is approximately 2,000 km thick and the density falls by a factor of almost a million from bottom to top, so the magnetic field begins to dominate the chromospheric structure ($\beta < 1$). The second law of thermodynamics does not permit heating of the chromosphere with the thermal

energy of the cooler photosphere below. Biermann (1948), Schwarzschild (1948) and Schatzman (1949) put forward ideas on the acoustic wave heating of the chromosphere as a result of the convective plasma motions in the photosphere and convection zone beneath. Referred to as the BSS model, the hypothesis is that acoustic waves transport energy upward with little dissipation once the velocity is below the sound speed. As the density drops and the velocity reaches the sound speed, the waves steepen into shocks and rapidly dissipate the energy, consequently heating the material (Zirin, 1998). However, acoustic wave heating does not apply in regions of strong magnetic field where motions, and therefore heating, are suppressed. This has led to work on Alfvén wave heating theories, first introduced by Osterbrock (1961), in which the magnetic field itself is responsible for depositing energy from the subsurface into the chromosphere and above. These theories better sit with observations of vigorous heating above plages and emerging flux regions, since they imply the amount of heating is proportional to the rate of magnetic change.

While the brightness of the photosphere overwhelms that of the chromosphere in the optical continuum, the hotter chromospheric temperatures lead to the hydrogen being ionised, resulting in strong H α emission. Filaments are observed as dark channels on-disk in H α images (called prominences when seen on the limb). Numerous plasma columns called spicules are also observed on the limb, that typically reach heights of $\sim 3,000-10,000$ km above the Sun's surface and are very short-lived (rising and falling over $\sim 5-15$ minutes).

Between the chromosphere and corona lies the transition region where the temperature jumps rapidly to over 1 MK. It is only about 100 km thick and it marks the point where magnetic forces dominate completely over gravity, gas




Figure 1.5: Soft X-ray image of the solar corona recorded by Yohkoh on 26 August 1992, reproduced from Aschwanden (2005).

pressure and fluid motion ($\beta \ll 1$). The extreme temperatures mean hydrogen is fully ionised and light is emitted in UV and EUV from carbon, oxygen and silicon ions (ref?).

1.2.3 Corona

The outermost part of the solar atmosphere is the corona, with electron densities ranging from $\sim (1-2) \times 10^{11}$ m⁻³ at its base height of $\sim 2,500$ km above the photosphere, to $\lesssim 10^9$ m⁻³ for heights $\gtrsim 1$ R_{\odot} above the photosphere (Aschwanden, 2005). The density varies across coronal holes which can have a base

density of $\sim (0.5-1) \times 10^{11}$ m⁻³, or across streamer regions with higher densities of $\sim (3-5) \times 10^{11} \text{ m}^{-3}$. Active regions that suspend and confine plasma in strong over-arching magnetic fields usually have the highest coronal densities of $\sim 2 \times 10^{11} - 2 \times 10^{12} \text{ m}^{-3}$. The temperature of the corona is generally $\gtrsim 1 \text{ MK}$, as indicated by emission from highly ionised iron lines for example, which again appears to contradict the second law of thermodynamics given the much cooler layers of the chromosphere and photosphere below (the 'coronal heating problem'). Its temperature structure is far from homogeneous, revealed in images such as that of Figure 1.5 from the Yohkoh Soft X-ray Telescope (SXT). Loop structures are observed at temperatures of 2-6 MK across regions of increased magnetic field density (such as above active regions/sunspots), and closed field regions are observed at temperatures of 1-2 MK across the quiet Sun, while open field regions of coronal holes have temperatures $\lesssim 1$ MK. These high temperatures lead to EUV and X-ray emission due to electron scattering processes off the highly ionised ions, although it is possible to image the corona in white light during a solar eclipse or with the use of a coronagraph that occults the solar disk, which is six orders of magnitude brighter in optical wavelengths. The corona we observe comprises several parts:

• The K-corona is a strongly polarised continuous emission spectrum due to Thomson scattering of photospheric light by the free electrons of the coronal gas, and it dominates up to ~2 R_⊙(ref?). It produces a polarised whitelight continuum without the Fraunhofer lines which are broadened out by doppler shifts due to the fast electron motions at such high temperatures. The intensity of the K-corona gives the coronal electron density (Koutchmy et al., 1991).

- The F-corona is due to scattering of sunlight by interplanetary dust particles, and contains the Fraunhofer lines. It is roughly equal in intensity to the K-corona at $\sim 4~\rm R_{\odot}$, and dominates at greater distances.
- The E-corona is due to emission from highly ionized coronal atoms such as iron and calcium.
- The T-corona is caused by thermal (infrared) emission of the interplanetary dust. It is an unpolarised continuum, insignificant in the visible part of the spectrum.

In contrast to the chromosphere, solar interior, and indeed the heliosphere, the magnetic pressure in the corona dominates over the gas pressure and so governs the coronal plasma dynamics ($\beta < 1$). The coronal structure we observe is thus shaped by the magnetic fields of the Sun, resulting in extended polar regions where there is mainly open magnetic field, and 'helmet-streamers' spanning the equatorial latitudes where, except for coronal holes, the field is mostly closed. Since these features are magnetically governed, the shape of the corona varies greatly over the solar activity cycle: it appears rounder at solar maximum, when multiple streamers emerge at various latitudes distributed across the Sun; and it appears more elliptical at solar minimum, when only a few streamers are present, lying closer to the equator.

Following Chapman & Zirin (1957) the description of a static corona leads to an unreasonable pressure value at large distances from the Sun. This is outlined below, beginning with the assumption that the corona is in hydrostatic equilibrium:

$$\frac{dP}{dr} = -\rho \frac{GM_{\odot}}{r^2} \tag{1.7}$$

The plasma density is $\rho = nm_p$, the pressure contribution from the protons and electrons is $P = 2nk_BT$, and the coronal heat flux is $q = \kappa \nabla T$ with thermal conductivity $\kappa = \kappa_0 T^{5/2}$. In the absence of heat sources or sinks $\nabla \cdot q = 0$ so in a spherically symmetric system we can write:

$$\frac{1}{r^2}\frac{d}{dr}\left(r^2\kappa_0 T^{5/2}\frac{dT}{dr}\right) = 0 (1.8)$$

Applying the boundary condition that the temperature tends to zero at large distances from the Sun, we obtain:

$$T = T_0 \left(\frac{r_0}{r}\right)^{2/7} \tag{1.9}$$

where $T_0=2$ MK is the temperature of the low corona at height $r_0=1.05~{\rm R}_{\odot}$ from Sun centre. This would mean $T\approx 4\times 10^5$ K at Earth (1 AU $\approx 215~{\rm R}_{\odot}$), close to measured values. Rewriting in terms of pressure and integrating, results in:

$$P(r) = P_0 \exp\left(\frac{7}{5} \frac{GM_{\odot}m_p}{2kT_0r_0} \left[\left(\frac{r_o}{r}\right)^{5/7} - 1 \right] \right)$$
 (1.10)

which implies that as $r \to \infty$ the coronal pressure tends towards a finite constant value $P \gg P_{ISM}$. This means the static coronal model is unphysical, and a dynamic model in which the material flows outward from the Sun must be considered, leading to a description of the solar wind.

1.2.4 Solar Wind

The solar wind is the constant out-stream of charged particles of plasma from the Sun's atmosphere due to the persistent expansion of the solar corona. The




Figure 1.6:

wind consists mostly of electrons and protons at energies of ~ 1 keV, traveling at speeds of 400 km s⁻¹ (slow solar wind) up to 800 km s⁻¹ (fast solar wind, found in regions of open field lines such as coronal holes). Thermal velocities of the particles are calculated at ~ 260 km s⁻¹ for coronal temperatures on the order of 3×10^6 K, while the escape velocity in the Sun's gravitational field in the low corona can be ~ 500 km s⁻¹. The additional energy to accelerate the solar wind is imparted by the pressure gradient $P_{Sun} \gg P_{ISM}$ to attain the measured solar wind speeds (Parker, 1958). The Parker model assumes the outflow is steady, spherically symmetric and isothermal. The momentum conservation equation of

the corona takes the form:

$$\rho v \frac{dv}{dr} = -\frac{dP}{dr} - \rho \frac{GM_{\odot}}{r^2} \tag{1.11}$$

Considering mass conservation $\dot{m} = 4\pi r^2 \rho v = constant$, we obtain:

$$\frac{\partial}{\partial r} \left(r^2 \rho v \right) = 0 \quad \Rightarrow \quad \frac{1}{\rho} \frac{\partial \rho}{\partial r} = -\frac{1}{v} \frac{\partial v}{\partial r} - \frac{2}{r} \tag{1.12}$$

So for a perfect gas $P=R\rho T$ equation 1.11 can be written:

$$\left(v - \frac{RT}{v}\right)\frac{\partial v}{\partial r} - \frac{2RT}{r} + \frac{GM_{\odot}}{r^2} = 0$$
(1.13)

A critical point occurs when $\partial_r v \to 0$ so we define:

$$r_c = \frac{GM_{\odot}}{2v_c^2}$$
 where $v_c = \sqrt{RT}$ (1.14)

and rewrite equation 1.13 as:

$$\left(v^2 - v_c^2\right) \frac{1}{v} \frac{\partial v}{\partial r} = 2 \frac{v_c^2}{r^2} (r - r_c) \tag{1.15}$$

Integrating equation 1.15 gives Parker's 'solar wind solutions':

$$\left(\frac{v}{v_c}\right)^2 - \ln\left(\frac{v}{v_c}\right)^2 = 4\ln\left(\frac{r}{r_c}\right) + 4\frac{r_c}{r} + C \tag{1.16}$$

where C is a constant of integration, leading to five potential solutions as plotted in Figure 1.6. Solutions I and II are double-valued, with II being disconnected from the surface. Solution III is too large (supersonic) close to the Sun. So-




Figure 1.7: Schematic of the Parker Spiral in the heliosphere. The streamlines of the solar wind act to drag out the magnetic field lines of the Sun, which become wound up in an Archimedean spiral as a result of the Sun's rotation. *Image credit:* Steve Suess, NASA/MSFC.

lution IV is called the 'solar breeze' as it remains subsonic. Solution V is the standard solar wind solution, although the assumptions of radial expansion and isothermality are not completely true in reality, so it is only an approximate characterisation of the observed solar wind. Nonetheless it is sufficient to convey the dynamic expansion of the corona and ultimate supersonic regime of outflow, often described akin to a de Laval nozzle which is used to accelerate flows from subsonic to supersonic speeds (as detailed in Goossens, 2003, for example).

Since the gas pressure dominates over the magnetic pressure in the solar atmosphere ($\beta > 1$), the solar wind acts to drag out the magnetic field lines of the Sun which become wound up as a result of solar rotation to form the Parker Spiral (Figure 1.7). This is an Archimedan spiral drawn by the magnetic field lines as they are advected outward by the solar wind, described by the equation:

$$r - r_0 = \frac{v}{\Omega} \left(\theta - \theta_0 \right) \tag{1.17}$$

where θ is the polar angle, $\Omega = 2.7 \times 10^{-6}$ rad s⁻¹ is the angular rotation rate of the Sun, r is the distance, and v is the solar wind speed (Parks, 2004; Zirin, 1998). The different speed streams can also lead to the formation of co-rotating interaction regions (CIRs) where the fast wind encounters the slow wind ahead of it in the Parker spiral, and can form shocks in the solar wind.

The solar wind does not extend infinitely, but eventually terminates when it reaches the edge of the heliosphere. The point where the solar wind slows from supersonic to subsonic speeds is called the termination shock, beyond which the wind comes into pressure balance with the interstellar medium (ISM) to form the heliosheath, whose outer boundary is called the heliopause (Figure 1.7). In the heliosheath the continually slowing wind is compressed and becomes turbulent through its interaction with the ISM. As the heliosphere moves through interstellar space, a bow shock is thought to form ahead of the heliopause as it encounters the ISM.

1.3 Coronal Mass Ejections

"We define a coronal mass ejection (CME) to be an observable change in coronal structure that 1) occurs on a time scale of a few minutes and several hours and 2) involves the appearance (and outward motion) of a new, discrete, bright, white light feature in the coronagraph field of view."

- (Hundhausen *et al.*, 1984)

Through the association of early observed flares and the detections of geomagnetic storms at Earth, the theory was put forward that plasma transients may be ejected from the Sun and possibly impact the Earth's magnetic field several days later (Lindemann, 1919). Observations of prominence disappearances provided evidence that the material may be rising through the corona with increasing velocity to eventually exceed the escape velocity of the Sun and erupt in to space (Kiepenheuer, 1953). Theories were developed by scientists who postulated that the magnetic field should be affected by these ejections, as they might act to drag out field lines or sever completely from the Sun through magnetic reconnection and potentially drive shock disturbances in the interplanetary gas (Gold, 1962; Piddington, 1958). It was only with the advent of space-borne coronagraphs were these transients realised to be a common occurrence on the Sun and their potential geomagnetic effects on Earth led them to become a topic of great interest. An example of such observations is shown in Figure 1.8.

These coronal mass ejections (CMEs), as they became known, are the largest manifestation of the shedding of solar magnetic field during the Sun's 22 year cycle. Every 11 years the magnetic axis of the Sun flips, giving rise to periodic patterns in the activity called solar minimum and maximum. At solar minimum




Figure 1.8: Observation of a CME from the LASCO/C2 coronagraph on board the SOHO spacecraft. The field-of-view extends from the inner mask at $\sim 2~R_{\odot}$ to the edges at $\sim 6~R_{\odot}$. The white circle represents the relative size and location of the Sun. The complexity of the magnetic field driving the eruption is clearly indicated by the twisted geometry of the bright ejecta.

a CME may occur up to once a week but during solar maximum they can be as frequent as three a day. In cases of halo CMEs coming toward the Earth, the particles densities and energies involved can cause geomagnetic storms, especially if the magnetic orientation of the CME is oppositely directed to that of Earth's magnetosphere. This is referred to as space weather, and understanding this interaction is of considerable practical importance because technological systems, such as communications and navigation satellites, can suffer interruptions or damage. To this end, missions such as the Solar and Heliospheric Observatory (SOHO; Domingo et al., 1995) and Solar Terrestrial Relations Observatory (Kaiser et al., 2008) have been launched to study CMEs.

CMEs are observed as a typical three-part plasma structure of a bright leading front, dark cavity, and bright core (Illing & Hundhausen, 1985). This configuration is indicative of a magnetic loop system erupting off the Sun as a flux rope or magnetic bubble with plasma embedded within and coronal material being swept up ahead of it. Thus the forces acting on CMEs are described within the context of magnetohydrodynamics as outlined below.

1.3.1 Magnetohydrodynamic Theory

The interplay between the plasma and magnetic fields of the Sun, notably in phenomena such as flares and CMEs, may be described through the coupling of the equations of electromagnetism with the theory of fluid motions. MHD attempts to combine Maxwell's equations with the fluid equations through the relative dependence on the electron motion in the currents set up in the plasma and the effects of the magnetic fields. Thus we obtain the induction equation for

magnetised plasma and describe how the field may undergo non-ideal (resistive) MHD processes such as magnetic reconnection - an important basis of many CME models.

1.3.1.1 Maxwell's Equations

Maxwell's equations describe the interaction of magnetic field ${\bf B}$ and electric field ${\bf E}$ according to:

$$\nabla \times \mathbf{B} = \mu \mathbf{j} + \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$$
 (1.18)

$$\nabla \cdot \mathbf{B} = 0 \tag{1.19}$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \tag{1.20}$$

$$\nabla \cdot \mathbf{E} = \frac{1}{\epsilon} \rho \tag{1.21}$$

where \mathbf{j} is the current density, ρ is the charge density, μ is the magnetic permeability of a vacuum, ϵ is the permittivity of free space, and c is the speed of light. The second term of Ampére's law (equation 1.18) may be neglected if the typical plasma velocities are much less than the speed of light:

$$\nabla \times \mathbf{B} = \mu \mathbf{j} \tag{1.22}$$

1.3.1.2 Fluid Equations

The mass continuity equation states that matter is neither created nor destroyed:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \rho \mathbf{v} = 0 \tag{1.23}$$

where ρ is the plasma density, and ${\bf v}$ the plasma velocity. This can be expanded to give:

$$\frac{\partial \rho}{\partial t} + (\mathbf{v} \cdot \nabla)\rho + \rho \nabla \cdot \mathbf{v} = 0 \tag{1.24}$$

where for an incompressible fluid the convective time derivative is zero. (This is the derivative $D/Dt = \partial/\partial t + \mathbf{v} \cdot \nabla$ taken along a path moving with velocity \mathbf{v} .) So the mass continuity equation reduces to:

$$\nabla \cdot \mathbf{v} = 0 \tag{1.25}$$

The equation of motion ($\mathbf{F} = m\mathbf{a}$) for a CME may be written:

$$\rho \frac{D\mathbf{v}}{Dt} = -\nabla p + \mathbf{j} \times \mathbf{B} + \rho \mathbf{g} - \frac{1}{2}\rho \mathbf{v}^2 A_{cme} C_D$$
 (1.26)

where p is the pressure, $\mathbf{j} \times \mathbf{B}$ is the Lorentz force, \mathbf{g} is gravity, and the drag force depends on the cross-sectional area A_{cme} and drag coefficient C_D . We neglect viscous forces.

In addition, Ohm's law couples the plasma velocity to the electromagnetic fields by:

$$\mathbf{j} = \sigma(\mathbf{E} + \mathbf{v} \times \mathbf{B}) \tag{1.27}$$

1.3.1.3 The Induction Equation

It is possible to eliminate the electric field **E** by combining Ampére's law (equation 1.18) and Ohm's law (equation 1.27):

$$\mathbf{E} = -\mathbf{v} \times \mathbf{B} + \frac{1}{\mu \sigma} \nabla \times \mathbf{B} \tag{1.28}$$

and substituting into Faraday's law (equation 1.21) to obtain:

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}) - \nabla \times (\eta \nabla \times \mathbf{B}) \tag{1.29}$$

$$= \nabla \times (\mathbf{v} \times \mathbf{B}) - \eta \nabla \times (\nabla \times \mathbf{B}) \tag{1.30}$$

$$= \nabla \times (\mathbf{v} \times \mathbf{B}) + \eta \left[\nabla^2 \mathbf{B} - \nabla (\nabla \cdot \mathbf{B}) \right]$$
 (1.31)

where $\eta = 1/\mu\sigma$ is the magnetic diffusivity. Using the solenoidal constraint (equation 1.19) provides the induction equation:

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}) + \eta \nabla^2 \mathbf{B}$$

$$O\left(\frac{B}{t}\right) \sim O\left(\frac{vB}{l}\right) + O\left(\frac{\eta B}{l^2}\right)$$
(1.32)

This equation forms the basis of any model that considers magnetised plasma motion on a variety of length scales, e.g., from magnetic confinement devices on Earth, to the dynamo action of the Sun's magnetic field. How a magnetic field topology will respond to the forces of plasma motion, and vice versa, is governed by the ratio of the terms in the induction equation.

1.3.1.4 The Magnetic Reynolds Number

The magnetic Reynolds number is the ratio of the advection and diffusion terms in the induction equation:

$$R_m = \frac{\nabla \times (\mathbf{v} \times \mathbf{B})}{\eta \nabla^2 \mathbf{B}} \approx \frac{v_0 l_0}{\eta}$$
 (1.33)

for plasma speed v_0 and length scale l_0 .

If $R_m \gg 1$ the induction equation is approximated by:

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}) \tag{1.34}$$

and the coupling of the magnetic field to the plasma motion is strong and the topology of the field changes on the timescales of the plasma motion:

$$\frac{B}{\tau_{motion}} \approx \frac{v_0}{l_0} B \quad \Rightarrow \quad \tau_{motion} \approx \frac{l_0}{v_0}$$
 (1.35)

This is known as the 'frozen-in' condition, whereby field lines are carried with the plasma motion. For example, in the corona the length scales are $l_0 \sim 1,000$ km, velocities are $v_0 \sim 1,000$ m s⁻¹, and the magnetic diffusivity $\eta \sim 1$ m² s⁻¹, resulting in a magnetic Reynold's number of $R_M \sim 10^8$.

If $R_m \ll 1$ the induction equation is approximated by:

$$\frac{\partial \mathbf{B}}{\partial t} = \eta \nabla^2 \mathbf{B} \tag{1.36}$$

and the magnetic field can diffuse through the plasma and change its topology:

$$\frac{B}{\tau_{diffusion}} \approx \eta \frac{B_0}{l_0^2} \quad \Rightarrow \quad \tau_{diffusion} \approx \frac{l_0^2}{\eta}$$
 (1.37)

Magnetic diffusion is an important condition for magnetic reconnection to occur such as in a current sheet where the length scales are small enough (on the order of metres) to account for the observed restructuring of magnetic field (which is on the order of seconds in flaring active regions for example).

1.3.1.5 Magnetic Reconnection

Magnetic reconnection is generally defined as a change in connectivity of field lines in time. In an ideal plasma the 'frozen-in' condition is met and the magnetic field is coupled to the plasma motion. However, when regions of opposite polarity flux come together, a boundary layer will form to separate the two regimes of magnetic field in a form of pressure balance. The high resistivity of such a system counteracts the currents within and allows the occurrence of non-ideal MHD processes and the formation of structures having small spatial scales, e.g., a thin current sheet. This leads to a low magnetic Reynolds number and allows diffusion to occur (illustrated by the shaded regions in Figure 1.9). The connectivity of field lines then changes to a more energetically favourable configuration, and in doing so will eject plasma along resulting outflows as the reconnected field lines relax to a new equilibrium. The outflows create a low pressure in the diffusion region which in turn allows continued inflow of plasma and magnetic field forming a runaway process of reconnection until the system comes to rest in a new topology.

A description of how magnetic reconnection occurs was put forward by Sweet (1958) and Parker (1957) as a two-dimensional incompressible MHD approximation. They estimate the rate of reconnection from a boundary layer analysis (top of Figure 1.9). For a reconnection layer of length Δ and thickness δ , the outflow must balance the inflow:

$$v_{in}\Delta = v_{out}\delta \tag{1.38}$$

where v_{in} is the inflow reconnection velocity, and v_{out} is the outflow velocity, which by conservation of energy $(B_x^2/2\mu = \rho v_{out}^2/2)$ is equal to the Alfvén velocity $v_A = B_0/\sqrt{4\pi\rho}$. From Ohm's law (equation 1.27) the configuration of the straight




Figure 1.9: Geometry of the Sweet-Parker (top) and Petschek (bottom) reconnection models, reproduced from Aschwanden (2005).

field lines $(\nabla \times B = 0)$ outside the layer is $E_z + v_R B_x = 0$, and inside the layer (where there is a large current) is $E_z = \eta J_z$. Integrating Ampere's law (equation 1.22) around the layer gives $B_x = \mu J_z \delta$. Combining the results of these two laws gives:

$$v_{in} = \frac{E_z}{B_x} = \frac{\eta J_z}{\mu J_z \delta} = \frac{\eta}{\mu \delta} \tag{1.39}$$

This can be written in terms of the outflow velocity (or interchangeably the Alfvén velocity) by:

$$v_{in}^{2} = \left(v_{out}\frac{\delta}{\Delta}\right)\left(\frac{\eta}{\mu\delta}\right) = v_{out}^{2}\left(\frac{\eta}{v_{A}\mu\Delta}\right)$$
 (1.40)

The rate of reconnection is then written in terms of the Lunquist number S, which is the dimensionless ratio of an Alfvén wave crossing timescale to a resistive diffusion timescale:

$$\frac{v_{in}}{v_{out}} = \frac{1}{\sqrt{S}} \quad \text{where} \quad S = \frac{\mu \Delta v_A}{\eta}$$
 (1.41)

This is the Sweet-Parker result. The problem with the theory is that it predicts reconnection to take place on far too slow timescales to reconcile with observations. Consider solar flares, for example, with $v_A \sim 1000$ km s⁻¹, and $l \sim 10^3$ km resulting in Sweet-Parker reconnection of tens of days when flare energy release is actually observed over minutes to hours.

An extension to the Sweet-Parker model was put forward by Petschek (1964) in which the field lines do not have to reconnect along the entire length of the boundary, but could merge over a shorter length $\Delta' < \Delta$ (bottom of Figure 1.9). The remaining length of the boundary is occupied by slow shocks, where the magnetic field tension accelerates the plasma to the Alfvén velocity. The reconnection velocity in the Petschek model may be written:

$$v_{in}^2 = v_{out}^2 \left(\frac{\eta}{v_A \mu \Delta'}\right) = \frac{v_{out}}{\sqrt{S}} \sqrt{\frac{\Delta}{\Delta'}}$$
 (1.42)

a factor of $\sqrt{\Delta/\Delta'}$ faster than the Sweet-Parker reconnection velocity. Issues with the description of the magnetic field and shock formation place limits on the plausibility of Petschek's formalism that Δ' is a free parameter which may be minimised to obtain the maximum reconnection velocity (Kulsrud, 2001). Numerical simulations, such as that of Biskamp (1986), are in favour of the Sweet-Parker

result, unless anomalous resistivity, for example, is considered.

1.3.2 Theoretical CME Models

It is well known that CMEs are associated with filament eruptions and solar flares (Moon et al., 2002; Zhang & Wang, 2002) but the driver mechanism remains elusive. Several theoretical models have been developed in order to describe the forces responsible for CME initiation and propagation, all of which are based on the idea that some form of instability must trigger the eruption. These models may be explained in terms of the following mechanical analogues, illustrated in Figure 1.10.

The Thermal Blast Model proposes that the increased thermal pressure produced from a flare overcomes the magnetic field tension and blows it open to cause a CME. Observations, however, have shown that not all CMEs are preceded by a flare, nor even necessarily associated with a flare at all.

The Dynamo Model introduces the idea of magnetic flux injection or stressing of the field on a time-scale that is too fast for the system to dissipate the magnetic energy before it builds to a critical point and erupts.

The Mass Loading Model is concerned with the amount of material included in the eruption. Prominences, or regions of relatively higher electron density in the corona, overlaying a volume of lower density will erupt due to the Rayleigh-Taylor instability.




Figure 1.10: Illustrations of the different mechanical analogues of CME eruptions, reproduced from Klimchuk (2001).

The Tether Release Model is based on the restraining of the outward magnetic pressure by the magnetic tension of the overlying field. As 'tethers' are removed a loss-of-equilibrium occurs due to the magnetic pressure/tension imbalance and the system erupts.

The Tether Straining Model is a variant on the tether release model whereby an increase in magnetic pressure due to flux injection or field shearing eventually overcomes the tension forces and the 'tethers' break and release the CME.

The tether straining and release models are generally accepted as the most likely scenarios for CME initiation, being able to reproduce numerous observations of CMEs through the development of detailed 2D and 3D flux rope models, as discussed below. Within the context of MHD outlined in Section 1.3.1, we can describe the solar plasma as a fluid with the assumption that there is negligible viscosity, so the motion of the flux rope is governed by the forces of Equation 1.26. The Lorentz force is thought to be the dominant driver force in modeling CME eruptions, certainly during the early stages of propagation, before the drag force takes over and the CME propagates with the ambient solar wind through interplanetary space.

1.3.2.1 Catastrophe Model

The 2D flux rope model is driven by a catastrophic loss of mechanical equilibrium as a result of footpoint motions in the photosphere (Forbes & Isenberg, 1991; Forbes & Priest, 1995; Isenberg *et al.*, 1993; Priest & Forbes, 2000, 1990a, 2002).




Figure 1.11: Theoretical evolution of a 2D flux rope, reproduced from Forbes & Priest (1995). The flux rope foot-point separation λ decreases, increasing the magnetic pressure until the flux rope becomes unstable and erupts away from the surface (b-d). The resulting height evolution of the flux rope is illustrated in (a).

The model is illustrated in Figure 1.11 as a coronal current filament channel and overlying magnetic field lines, in equilibrium due to the balance between the magnetic pressure and tension forces acting on the system. The description of the model's evolution in time may be split into a storage phase and an eruption phase. During the storage phase the footpoints of the system are slowly moved together such that the magnetic energy of the flux rope increases. The magnetic tension thus increases and causes the flux rope to move downwards, which builds up magnetic pressure until a critical footpoint distance is reached where equilibrium is lost. In the eruption phase the flux rope is accelerated upwards, stretching the magnetic field lines such that a current sheet forms behind it. If reconnection occurs in the current sheet then all the energy is released and the upward motion of the flux rope is unbounded. Otherwise it will come to equilibrium again at a greater height, or oscillate about this equilibrium height if it still has energy in excess of that required for the initial eruption. The configuration of this model outlined in Priest & Forbes (2000) predicts kinematics of the flux rope (provided it is 'thin' so that its radius is less than the scale-length λ_0) prior to the formation of the current sheet (i.e. $h/\lambda_0 \leq 2$) according to:

$$\dot{h} \approx \sqrt{\frac{8}{\pi}} v_{A0} \left[\ln \left(\frac{h}{\lambda_0} \right) + \frac{\pi}{2} - 2 \tan^{-1} \left(\frac{h}{\lambda_0} \right) \right]^{1/2} + \dot{h}_0$$
 (1.43)

where \dot{h} is the velocity of the flux rope, \dot{h}_0 is the initial perturbation velocity, λ_0 is the source separation at the critical point, and v_{A0} is the Alfvén speed at $h = \lambda_0$. The kinematics may be further separated into an 'early' phase when the

time scale is less than the Alfvén time scale $(t \ll \lambda_0/v_{A0})$, which gives:

$$\dot{h} \approx \dot{h}_0 + \frac{2v_{A0}}{\sqrt{3\pi}} \left(\frac{\dot{h}_0}{\lambda_0}\right)^{3/2} t^{3/2}$$
 (1.44)

and a 'late' phase when $h/\lambda_0 \gg 1$, but $|\ln h|$ is still much smaller than $|\ln a|$, which gives:

$$\dot{h} \approx \sqrt{\frac{8}{\pi}} v_{A0} \left[\ln \left(\frac{h}{\lambda_0} \right) - \frac{\pi}{2} \right]^{1/2} \tag{1.45}$$

After the formation of the current sheet the system becomes too complicated for the kinematics of the CME's continued propagation to be analytically derived.

1.3.2.2 Toroidal Instability

An extension of the flux rope model to three-dimensions is illustrated in Figure 1.12 (Chen, 1996; Chen & Krall, 2003; Kliem & Török, 2006). The eruption of the flux rope is triggered by an increase in the poloidal magnetic flux of the structure. The 3D flux rope consists of a current channel **J** and magnetic field **B**, and has major radius R and minor radius a such that for r < a the magnetic field lines are helical and can be described by their toroidal and poloidal components, but for r > a the field is purely poloidal ($J_t = 0$). The major radius R is fixed, and the minor radius increases from a_f at the footpoints to a_a at the apex. The footpoints are assumed to be immobile because of the high density photosphere ($\sim 10^{23} \text{ m}^{-3}$) relative to the corona ($\lesssim 10^{16} \text{ m}^{-3}$). The poloidal field B_p is also highly non-uniform in the photosphere since $\beta \gg 1$.

The model may be directly compared to coronagraph observations as in Krall et al. (2001), where the leading edge of the CME front is located at Z + 2a with




Figure 1.12: A schematic of the 3D flux rope model, reproduced from Chen & Krall (2003), an extension of the 2D flux rope in Figure 1.11 when viewed end-on as indicated by the arrow from the right. The flux rope is rooted below the photosphere, and surrounded by the ambient coronal magnetic field $\mathbf{B_c}$ and plasma density ρ_c . Components of the current density \mathbf{J} and magnetic field \mathbf{B} are shown, where subscripts 't' and 'p' refer to the toroidal and poloidal directions respectively. The flux rope has a radius of curvature R, radius of cross-section a, apex height Z, footpoint separation s_f , and the radial force outward is F_R .

a width of 4a when viewed end-on, or a width of 2R + 4a when viewed sideon. This definition arises from the fact that the poloidal field B_p at r = 2a has decreased to about half the value of B_{pa} at r = a and is then comparable to the ambient coronal field B_c . This model sits well with observations, where the CME front corresponds to a plasma pileup ahead of the flux rope which appears as a darker cavity, and any erupting prominence material is suspended at the base of the flux rope and appears as the bright core of the CME. Background parameters such as coronal density and solar wind speeds are also specified in the model. The eruption is initiated by a poloidal flux injection that increases the toroidal current for a short period of time, increasing B_{pa} while R does not change significantly, such that the radial force $\mathbf{F_R}$ becomes more positive and exerts an upward net force on the structure. The eruption then proceeds through the corona as the external poloidal field decreases sufficiently rapidly in the direction of motion. The equation of motion (c.f. equation 1.26) may be written in terms of the forces acting to cause the apex motion:

$$M\frac{d^2Z}{dt^2} = F_R + F_g + F_d (1.46)$$

where the radial force F_R results from the Lorentz magnetic force and pressure gradient of the system, and may be written:

$$F_R = \frac{I_t^2}{c^2 R} f_R \tag{1.47}$$

showing how the toroidal current increase will add to the upward force on the structure. The change in current affects the inductance of the flux rope ($F_R \propto I_t^2 \propto L^{-2}$) and so, neglecting the gravity and drag terms, the acceleration may be expressed in terms of the geometrical size of the flux rope:

$$\frac{d^2 Z}{dt^2} \sim \frac{\Phi_p^2}{\left[R \ln (8R/a_f)\right]^2} f_R$$
(1.48)

Kliem & Török (2006) show how the height of the flux rope during the very initial stages of the eruption may be approximated as a hyperbolic function:

$$h(\tau) = \frac{P_0}{P_1} \sinh(P_1 \tau), \qquad h \equiv H/H_0 - 1 \ll 1$$
 (1.49)

where H is the height, and H_0 the initial height, of the flux rope; τ is the time normalised by the Alfvén time; P_0 comprises initial parameters on the flux rope dynamics; and P_1 associates the external magnetic field profile. Their simulations show a fast rise and gradual decay phase of the CME accelerations due to the toroidal instability. However, Schrijver *et al.* (2008) demonstrate that tuning the initial parameters changes the acceleration profile from a fast initial rise to a more gradual rise phase.

1.3.2.3 Breakout Model

In the magnetic breakout model the CME eruption is triggered by reconnection between the overlying field and a neighbouring flux system through the shearing of a multipolar topology, illustrated in Figure 1.13 (Antiochos et al., 1999; Lynch et al., 2004; MacNeice et al., 2004). It starts by shearing a potential field configuration consisting of a central arcade which will become the CME, two side arcades, and an overlying arcade, with a magnetic x-line separating the different topologies. This shearing adds magnetic pressure to the inner flux system and causes it to expand and distort the overlying field at the x-line, eventually forming a current sheet. As the current sheet grows, reconnection begins, transferring flux to the neighbouring arcades and creating a passage for the CME release as the central arcade erupts. A current sheet also forms beneath the erupting sheared field, creating a disconnected flux rope that escapes, and this is associated with flare reconnection. An increase in the rate of outward expansion drives a faster rate of breakout reconnection, yielding the positive feedback required for an explosive eruption. At given distances from the Sun, the simulation, which is intrinsically 2.5D but has recently been extended to 3D by Lynch et al. (2008),




Figure 1.13: Schematic, and corresponding simulation snapshots, of the main stages of the axisymmetric 2.5D breakout model, reproduced from Lynch *et al.* (2008). (a) shows the initial multipolar topology of the system, (b) shows the shearing phase which distorts the x-line and causes breakout reconnection to begin, (c) shows the onset of flare reconnection behind the eruption that disconnects the flux rope, and (d) shows the system restoring itself following the eruption.

produces a number of key observational properties to test against data (van der Holst et al., 2007). Simulations run by Lynch et al. (2004) produced kinematics of the CME front which showed constant acceleration, and they fit quadratics to the height-time data of the form:

$$h(t) = h_0 + v_0(t - t_0) + \frac{1}{2}a(t - t_0)^2$$
(1.50)

However, the height-time profiles from the 3D simulations by Lynch et al. (2008) showed a more complex kinematic profile with separate rising and breakout phases of acceleration, proving a better match with observations. The velocity increases linearly from zero during the initial shearing and breakout phase, followed by an acceleration peak during the second stage of reconnection behind the CME. DeVore & Antiochos (2008) find a three-phase acceleration profile with a similar initial slow and fast acceleration profile during the breakout and flare reconnections, followed by a short interval of fast deceleration as the magnetic field configuration is reformed after the eruption.

1.3.3 CME Observations

In order to test the validity of the theoretical CME models, and subsequently understand the forces governing their eruption through the solar atmosphere, comparisons must be made with observations. Generally the kinematics of events are determined from white-light coronagraph images, obtained through Thomson scattered emission of the plasma, by tracking the structure as it moves through the field-of-view.




Figure 1.14: The Thomson scattering geometry for a single electron, reproduced from Howard & Tappin (2009). **a**) shows the electron and incident light with different observer positions indicated. **b**), **c**), **d**) show the resultant scattering of light seen by each observer O_1 , O_2 , O_3 respectively.

1.3.3.1 Thomson Scattering

The low density, optically thin, plasma of the corona and solar wind is observable in white light through the process of Thomson scattering, whereby photons from the Sun are scattered by the free electrons of the coronal plasma (Billings, 1966; Minnaert, 1930; van de Hulst, 1950). Essentially, when light is incident on an electron, the electric field of the light waves will cause the electron to accelerate

(absorb a photon) and re-radiate (emit a photon) in the plane perpendicular to the incident wave (illustrated in Figure 1.14). Depending on the angle χ to the observer, the scattered light may be unpolarised (Figure 1.14b), partially polarised (Figure 1.14c), or polarised (Figure 1.14d). The tangential component of the scattered light intensity is isotropic, while the radial component varies as $\cos^2 \chi$, resulting in the following expression for the differential cross-section (Jackson, 1975):

$$\frac{d\sigma}{d\omega} = \frac{1}{2} \left(\frac{e^2}{4\pi\epsilon_0 m_e c^2} \right)^2 \left(1 + \cos^2 \chi \right) \tag{1.51}$$

where $d\omega$ is an element of solid angle at scattering angle χ , and ϵ_0 is the permittivity of free space. Since the Sun is neither a point-source, nor is the intensity of light uniform across it, it is necessary to consider the three-dimensional geometry of the Thomson scattering process in detail to appreciate the effects it will have on CME observations, outlined in detail in Billings (1966) and Howard & Tappin (2009). In a full description of the corona the density profile must be considered, as well as the effect of limb darkening on the Sun whereby the light intensity from the photosphere decreases across the disk according to:

$$I = I_0(1 - u + u\cos\psi) \tag{1.52}$$

where I is the intensity observed at angle ψ from the radial vector, I_0 is the radial intensity, and u is the limb-darkening coefficient.

Describing the Thomson sphere as the locus of all points that make an angle of $\chi = 90^{\circ}$ between the line-of-sight and the vector from the Sun to the scattering point P (Figure 1.15), the total intensity of scattered light is governed by three terms:




Figure 1.15: Schematic of the Thomson surface, being the sphere of all points which are located at an angle of 90° between the Sun and the observer, reproduced from Vourlidas & Howard (2006). An example line-of-sight is shown for an electron at point P, with radial distance r from the Sun, at longitude ϕ relative to the solar limb.

- 1. The scattering efficiency which is minimised on the Thomson sphere.
- 2. The incident intensity which is maximised on the Thomson sphere since that is where the line-of-sight is closest to the Sun.
- 3. The electron density in the scattering region which is maximised on the Thomson sphere since the solar wind density drops off with radial distance from the Sun.

The combination of these effects makes the Thomson sphere an important consideration when interpreting CME observations, especially out through the wide-angle fields-of-view of the heliospheric imagers on the STEREO spacecraft. Vourlidas & Howard (2006) note the following:

- CMEs that propagate along the solar limb and appear bright in near-Sun coronagraphs are unlikely to be detectable further out in the heliosphere.
- Frontside events are always brighter than their backside counterparts, and
 ones at intermediate angles will exhibit approximately constant levels of
 brightness over a wide range of heliocentric distances.
- The sky-plane assumption holds well for brightness observations out to at least $\sim 70~{\rm R}_{\odot}$.

If an expanding CME front moves along or crosses the Thomson sphere during its propagation, its observed brightness can change with regard to the changing location of its intersection with the sphere, as a result of the discussion above. This has implications for how the observed CME kinematics and morphology may be affected, certainly at large elongations from the Sun, and how CMEs may look quite distinct from different observers' points of view.

1.3.3.2 CME Kinematics & Morphology

A large number of CMEs have been studied since the advent of space-borne coronagraph observations, and they show speeds varying from tens up to a few thousand kilometres per second. Many of these exhibit a general multiphased kinematic evolution: CMEs tend to have an initial rise phase with possible high acceleration, and a subsequent constant-velocity cruise phase with another possible low acceleration or deceleration in their continued propagation through the heliosphere. This initially led Sheeley et al. (1999) to distinguish CMEs as either 'gradual' if their initial acceleration is low, or 'impulsive' if it is high. However, statistics on a large sample of events do not show such a clear distinction but do indicate that slow CMEs tend to result from prominence lift-offs or streamer blowouts and speed up to the solar wind speed, while fast CMEs tend to result from flares and active regions and slow down to the solar wind speed (González-Esparza et al., 2003; Gopalswamy et al., 2000; Moon et al., 2002). Statistical analyses can provide a general indication of CME properties, for example Zhang & Dere (2006) study 50 CMEs and find an average acceleration of 330.9 m $\rm s^{-2}$ with an average duration time of 180 minutes, and Gopalswamy et al. (2000) study 28 CMEs and derive a formula for their acceleration a related to their initial speed u by a = 1.41 - 0.0035u. However, plane-of-sky projection effects mean the measured kinematics are not representative of the true CME motion (Figure 1.16), with Wen et al. (2007) deducing that the error in CME leadingedge measurements grows roughly with the square of the distance from Sun centre within the first few solar radii and then varies approximately with the square root of the distance past $\sim 5 \text{ R}_{\odot}$. In an effort to overcome plane-of-sky effects, dal




Figure 1.16: .

Lago et al. (2003) use a sample of 57 limb CMEs to derive an empirical relation between their radial and expansion speeds as $V_{rad} = 0.88V_{exp}$, and Schwenn et al. (2005) similarly use 75 events to derive a formula for their transit time $T_{tr} = 203 - 20.77 \ln{(V_{exp})}$. However, Vršnak et al. (2007) show the inherent difficulties in performing and trusting such corrections for CME projection effects. This means individual CMEs must be studied with rigour in order to satisfactorily derive the kinematics and morphology to be compared with theoretical models.

An example of such rigourous analysis is seen in the kinematic study of Gallagher *et al.* (2003). In order to overcome the lower limit of coronagraph observations $\lesssim 3~\rm R_{\odot}$ they include the use of low corona EUV observations for a CME on the 21 April 2002, plotted in Figure 1.17. They track the CME front through




Figure 1.17: (a) Height-time, (b) velocity and (c) acceleration profiles for the CME on 21 April 2002, and (d) the GOES-10 soft X-ray flux for the associated X1.5 flare during the interval 00:47-03:20 UT, reproduced from Gallagher *et al.* (2003). A three-point difference scheme is used on the data points, and a first-difference scheme is plotted with filled circles. The solid line is the best fit of the exponentially increasing and decreasing acceleration of equation 1.53.

running-difference images and measure its height-time profile from ~ 20 Mm to over 10^4 Mm above the Sun's surface (almost $30~R_{\odot}$). An early high-acceleration phase is revealed across the initial $1-2~R_{\odot}$ height range of the event, which they model with a combined function of exponentially increasing and decreasing acceleration:

$$a(t) = \left[\frac{1}{a_r \exp(t/\tau_r)} + \frac{1}{a_d \exp(-t/\tau_d)}\right]^{-1}$$
 (1.53)

where a_r and a_d are the initial accelerations, and τ_r and τ_d the e-folding times, for the rise and decay phases. They reveal an early acceleration peak of $\sim 1,500 \text{ km s}^{-1}$ and show it to correspond with the duration of the soft X-ray rise phase of the associated X1.5 flare, having implications for either the thermal blast model or a magnetically-dominated process such as reconnection causing the eruption.

Maričić et al. (2004) present a study of the initiation and development of a limb CME on 15 May 2001, tracking its leading edge, cavity and associated prominence from 0.32 R_☉ out to ~30 R_☉, plotted in Figure 1.18. They distinguish a preacceleration characterised by the slow rising motion of the prominence, suggestive of an evolution of the system through a series of quasi-equilibrium states. They offer numerous possible explanations for this: a slow shearing or merging motion of the arcade footpoints (e.g. Lin et al., 2004; Priest & Forbes, 1990b), twisting of the embedded flux rope (Vrsnak, 1990), emerging azimuthal flux (Chen & Krall, 2003), mass loss from the prominence body (Vrsnak, 1990), etc. This is followed by a rapid acceleration onset of the prominence, 380 ± 50 m s⁻², simultaneous with the CME leading edge acceleration, 600 ± 150 m s⁻². This simultaneity rules out the scenarios in which the prominence motion is merely a consequence of the disruption of the overlying magnetic structure, or that the prominence




Figure 1.18: The kinematics of the CME leading edge (circles), cavity (pluses) and prominence (triangles) of the 15 May 2001 event, reproduced from Maričić et al. (2004). (a) shows the height-time profiles. (b) shows the distance between the leading edge and the prominence (thick line), the cavity and the prominence (thin line), and the leading edge and the cavity (dashed line). (c) shows the velocities determined by forward-difference technique upon the smoothed height-times. (d) shows the onset of acceleration against height, with a straight line fit. The horizontal bar between (a) and (c) indicates the period of the soft X-ray burst (dotted - precursor, full - main rise).

eruption itself drives the upper parts of the system (at least for this event, and the authors suggest a careful inspection on a larger sample of events). There is also a simultaneous soft X-ray burst which they deem an 'acceleration precursor' as postulated in the catastrophic evolution of a flux rope/arcade system (Lin & Forbes, 2000). A possible gradual deceleration of the CME in its later stages is also noted $(-23\pm1~{\rm m~s^{-2}}$ for the leading edge) and attributed to the possible effects of aerodynamic drag in the solar wind discussed in Cargill *et al.* (1996), Vršnak (2001) and Vršnak *et al.* (2004). Maričić *et al.* (2004) also measure the relative height difference between the CME leading edge, cavity and prominence, and report a kind of self-similar expansion of the event.

Temmer et al. (2008) study the CME dynamics and associated flare flux profiles for two fast events of 17 January 2005 and 6 July 2006. Using running-difference techniques they track the rising loops of the flares and consider them as the early height-time profile of the subsequent CME eruption. This results in a clear early acceleration peak for each event, of $\sim 4,400 \text{ m s}^{-2}$ and $\sim 1,100 \text{ m s}^{-2}$ respectively. They find that the peaks of the hard X-rays correspond with the peak accelerations of the CMEs, indicating a strong feedback relationship indicative of magnetic reconnection occurring in a possible current sheet behind the CME to drive the eruption under the dominant Lorentz force.

More recently Lin *et al.* (2010a) performed the same type of analysis on two CMEs in order to specifically test the kinematics of each against the predictions of theoretical models (catastrophe, breakout and toroidal instability). Tracking the CME front through running-difference images, the events of 17 December 2006 and 31 December 2007 were found to have peak accelerations of $\sim 60 \text{ m s}^{-2}$ and $\sim 1,500 \text{ m s}^{-2}$ within the first $\sim 3 \text{ R}_{\odot}$ height range, simultaneous with the soft

X-ray flux. The authors show the difficulty in distinguishing any single model as a basis for the eruption, since the predicted fits all fare relatively well in producing a profile within the scatter and error of the measured kinematics. The authors also acknowledge that the data is a two-dimensional projection of actual motion, and three-dimensional kinematics of CMEs are necessary for improved accuracy.

The conclusions of such studies rely heavily on the errors involved in the derived kinematics if they are to provide a distinction between different models. The above discussion serves to highlight the importance of the image resolution, cadence and projection effects when performing CME analyses. It is also apparent that user-specific biases and different numerical methods for determining the kinematics and morphology can affect the resulting profiles and hence their interpretation. With these considerations in mind, the current aims of the community have been to develop methods for overcoming both the biases of individual users and single viewpoint observations in order to robustly determine the kinematics and morphology of CMEs with the greatest possible precision.

1.4 Outline of Thesis

The work presented in this thesis improves the understanding of the kinematics and morphology of CMEs as they propagate through the solar corona. To date the quantification of these properties has been subject to various sources of error, most notably as a result of the innate difficulties in tracking CMEs with traditional image processing techniques. Moreover the projected two-dimensional nature of coronagraph observations has been a persistent source of error that scientists have striven to overcome when studying CMEs. This thesis outlines

new methods of multiscale image processing and ellipse characterisation of the CME front in coronagraph observations so as to reduce the error on the derived kinematic and morphological parameters. This is extended for applying to stereoscopic image data whereby an elliptical tie-pointing methodology for reconstructing the CME front in three-dimensions leads to a study of its true kinematics and morphology as it propagates into the heliosphere.

Chapter 2 details the space-borne instrumentation used in this work for observing CMEs through the solar corona and heliosphere. Chapter 3 discusses the current CME catalogues in use and their limitations, leading to the implementation of new multiscale techniques and an ellipse characterisation for studying CME propagation. The efforts to automate this process for the development of a new cataloguing database are also discussed. These methods are the basis of the studies undertaken in Chapter 4, where a selection of CME events are presented and their kinematics and morphologies discussed in light of theory. Chapter 5 presents an important extension of these methods for performing a three-dimensional reconstruction of a CME front, which overcomes the issues of projection effects and so provides insight into the true kinematics and morphology of the eruption. A detailed discussion of the results and conclusions drawn from this work are also presented. Finally Chapter 6 presents the main conclusions of the thesis and details possible future work that could follow on from these new developments.

Chapter 2

Instrumentation

CMEs were initially observed in the early 1970s with the launch of the first space-borne coronagraph onboard the seventh Orbiting Solar Observatory (OSO 7) providing daily white-light coronal images with a field-of-view $2.8-10~\rm R_{\odot}$, observing about a dozen CMEs from 1971-1974 (Koomen et al., 1975; Tousey & Koomen, 1972). Skylab, a U.S. space station launched in 1973, housed a coronagraph with field-of-view $1.5-6~\rm R_{\odot}$ that imaged the corona every 6-8 hours and conclusively established these transient ejections as a common occurrence, observing $\sim 100~\rm in~1973/74$ (MacQueen et al., 1974). Following Skylab, several more coronagraphs were flown: SOLWIND on satellite P78-1 observed over 1,500 CMEs from 1979-1985 (Sheeley et al., 1980); the High Altitude Observatory Coronagraph/Polarimeter onboard Solar Maximum Mission (SMM) observed $\sim 1,350$ CMEs from 1980-1989 (MacQueen et al., 1980); the Large Angle Spectrometric Corongraph suite (LASCO; Brueckner et al., 1995) onboard the Solar and Heliospheric Observatory (SOHO; Domingo et al., 1995) has observed thousands of CMEs from 1995 to present; and the COR1/2 coronagraphs of the Sun-Earth

Connection Coronal and Heliospheric Imaging suite (SECCHI; Howard et al., 2008a) onboard the Solar Terrestrial Relations Observatory (STEREO; Kaiser et al., 2008) have been observing CMEs from 2006 to present. Data from the instruments of SOHO/LASCO and STEREO/SECCHI are used in the research presented throughout this thesis.

2.1 SOHO/LASCO

SOHO is a joint European Space Agency (ESA) and National Aeronautics and Space Administration (NASA) mission, launched on 2 December 1995 to undertake scientific investigations of (1) helioseismology, the study of the interior solar structure, and (2) the physical processes that account for the heating and acceleration of the solar wind, or more broadly the nature of evolutionary change in the Sun's outer atmosphere. It is situated in orbit about the Lagrangian L1 point approximately 1.5×10^6 km sunward of the Earth for an uninterrupted view of the Sun. Onboard are twelve complementary science instruments: three helioseismology experiments to probe the Sun's inner structure through measurements of solar oscillations; three solar wind experiments to measure the in-situ properties of the ambient wind (densities, speeds, charge states, etc.); and six telescopes and spectrometers to study the solar disk and atmosphere. A schematic of SOHO and its instrument suites is shown in Figure 2.1.

The LASCO instrument suite is a set of three coronagraphs C1, C2 and C3 that image the solar corona from 1.1-3, 1.5-6 and 3.7-30 R $_{\odot}$ respectively (however the C1 coronagraph has not been in operation since 1998 when contact with the SOHO spacecraft was lost for several weeks). The coronagraph was invented




Figure 2.1: A schematic of the SOHO spacecraft and onboard instrument suites, reproduced from Domingo *et al.* (1995).

by French astronomer Bernard Lyot in 1939 to artificially eclipse the Sun for observing the solar corona. It essentially blocks light rays from the centre of the telescope field-of-view by occulting the solar disk, in order to increase the relative intensity of the surrounding coronal light which is on the order of one million times fainter. The externally occulted Lyot coronagraph design of LASCO/C2 and C3 is illustrated in Figure 2.2. The top diagram demonstrates how the optical assembly images the coronal light, while the bottom diagram demonstrates how stray light is suppressed. Light is incident through aperture A0 where the external occulter D1 eclipses the solar disk. The light then enters aperture A1 and is focused by the objective lens O1, through the field stop, onto the inner occulter D2 which apodizes the bright fringe of the external occulter. Field lens O2 then collimates the light onto the Lyot stop A3 that intercepts the light rays diffracted off the entrance aperture A1. A relay lens O3 is placed behind A3 to




Figure 2.2: The externally occulted Lyot coronagraph, reproduced from Brueckner et al. (1995), showing: front aperture A0 and external occulter D1; entrance aperture A1 and objective lens O1; the field stop; inner occulter D2 and field lens O2; Lyot stop A3 and relay lens O3 with Lyot spot; filter and polariser wheels F/P; and the focal plane F.

focus the coronal image on to the plane F. O3 contains the Lyot spot for intercepting residual diffracted light from D1 and ghost images created by O1. In front of the focal plane F are the color filters and linear polarising filters F/P. The color filters distinguish specific bandpasses of the coronal light, in the ranges 400-850 nm for C2 and 400-1050 nm for C3. The polariser wheel is used to obtain total brightness B or polarised brightness pB images through combinations of polariser positions $I_a = -60^{\circ}$, $I_b = 0^{\circ}$, and $I_c = 60^{\circ}$, according to the




Figure 2.3: A LASCO/C3 image of the solar corona out to $\sim 30~R_{\odot}$.

equations (Billings, 1966):

$$B = 2(I_a + I_b + I_c) (2.1)$$

$$pB = \frac{4}{3}[(I_a + I_b + I_c)^2 - 3(I_aI_b + I_aI_c + I_bI_c)]^{1/2}$$
 (2.2)

A CCD is placed at the focal plane F and the final images are 1024×1024 pixels, subtending an angle of 11.4 arcseconds per pixel in C2, and 56 arcseconds per pixel in C3 (see Section 2.3 for CCD details).

2.2 STEREO/SECCHI

STEREO is the third mission in NASA's Solar Terrestrial Probes program (Kaiser et al., 2008). It was launched on 25 October 2006, and employs two nearly identical space-based observatories; one ahead of Earth in its orbit, and the other behind, separating at $\pm 22^{\circ}$ each year. This arrangement provides the first ever stereoscopic observations of the Sun and inner heliosphere. The main objectives of STEREO are to:

- Understand the causes and mechanisms of CME initiation.
- Characterize the propagation of CMEs through the heliosphere.
- Discover the mechanisms and sites of energetic particle acceleration in the low corona and the interplanetary medium.
- Improve the determination of the structure of the ambient solar wind.

STEREO hosts four instrument suites to achieve this, as illustrated in Figure 2.4. The SECCHI suite comprises five scientific telescopes: firstly the Sun




Figure 2.4: Payload diagram of one of the STEREO spacecraft, indicating the positions of the four instrument suites onboard: Sun-Earth Connection Coronal and Heliospheric Imagers (SECCHI); In-situ Measurements of Particles and CME Transients (IMPACT); Plasma and SupraThermal Ion Composition (PLASTIC); STEREO/WAVES radio burst tracker (SWAVES). Image credit: stereo.gsfc.nasa.gov.

Centred Imaging Package (SCIP) consisting of an Extreme Ultraviolet Imager (EUVI) of the solar disk out to 1.7 R_{\odot} and two coronagraphs (COR1/2) with fields-of-view 1.4–4 and 2–15 R_{\odot} ; and secondly the Heliospheric Imagers (HI) consisting of two wide-angle visible light imagers positioned on the sides of the STEREO spacecraft for fields-of-view extending out to Earth at 1 A.U. (astronomical unit, based on the distance from the Earth to the Sun which is approximately 1.49×10^8 km).




Figure 2.5: Diagram of the EUVI in the STEREO/SECCHI suite, reproduced from Howard *et al.* (2000).

2.2.1 EUVI

The Extreme Ultraviolet Imager is a normal-incidence Ritchey-Chrétien telescope that images the solar disk out to 1.7 R_{\odot} at four wavelengths of emission that span a temperature range of 0.1 to 20 MK (Wuelser *et al.*, 2004). Radiation from the Sun enters through a thin aluminium filter of 150 nm thickness that suppresses most of the ultraviolet, visible and infrared wavelengths of light. The radiation passes one of four quadrants that are each optimised for one of the EUV wavelength lines (listed in Table 2.1). The primary and secondary mirrors direct the light through a filter wheel that has a redundant thin-film aluminium filter to remove the remainder of the visible and IR radiation. A shutter in the path controls the exposure time, and 2048×2048 pixel images are produced by the CCD subtending an angle of 1.6 arcseconds per pixel (for details on the CCD see Section 2.3).




Figure 2.6: Diagram of the COR1 coronagraph in the STEREO/SECCHI suite, reproduced from Howard et al. (2000). The measurements are specified in millimetres.

Table 2.1: Summary of EUVI wavelengths.

Principal emission lines	Wavelength
Fe IX	172 Å
Fe XII	$194~\mathrm{\AA}$
Fe XV	284 Å
He II	304 Å

2.2.2 COR1

COR1 is a classic Lyot internally occulting refractive coronagraph (Thompson et al., 2003). Light enters through the front aperture of the telescope and is focused by the objective lens onto the occulter, with a series of baffles in place to minimise scattering of light within the telescope (Figure 2.6). The occulter is cone shaped to reject light from the centre of the field-of-view into a surrounding light trap. The field lens focuses the rest of the light down the telescope to the Lyot stop which removes light diffracted by the edge of the front aperture. A Lyot spot is also glued to the doublet lens immediately behind the Lyot stop in order to remove any ghosting of the objective lens. Two doublet lenses focus the light onto the CCD detector, with a bandpass filter 10 nm wide (centred on the $H\alpha$ line at 656 nm) and a linear polariser in between them. To extract both total brightness B and polarised brightness pB images, three sequential images are taken with polarisations of $I_a = -60^{\circ}$, $I_b = 0^{\circ}$, and $I_c = 60^{\circ}$, and combined using equations 2.1 and 2.2. The cut-on frequency (at 350 nm) of light through COR1 is set by the transmission of the BK7-G18 glass in the objective lens, and the cut-off frequency (at 1100 nm) is set by the band gap of the silicon CCD detector. The final images are 2×2 binned onboard to 1024×1024 pixels, subtending an angle of 7.5 arcseconds per pixel. The field is unvignetted except for a small area around the edge of the occulter and near the field stop in the corner of the images. The average radial profile for both instruments is well below 10^{-6} B/B $_{\odot}$ though some discrete ring-shaped areas of increased brightness in the Behind instrument are caused by features on the front surface of the field lens.

2.2.3 COR2

COR2 is an externally occulted Lyot coronagraph, similar to the LASCO/C2 and C3 telescopes (Figure 2.7). An array of internal baffles sits behind the external occulter to reduce stray light entering the telescope, and an internal occulter and Lyot stop minimise diffraction effects. The final images are produced at three different polarisations as in COR1 for creating total brightness and polarised brightness images. The final images are 2048×2048 pixels, subtending an angle of 14.7 arcseconds per pixel. The image is vignetted throughout the field-of-view, at a level of 40-50% around the occulter pylon, and reaching a minimum of 20% at about $10~R_{\odot}$ before increasing again towards the image edge.

2.2.4 Heliospheric Imagers

The Heliospheric Imagers (HI1/2; Eyles et al., 2009) are two small, wide-angle, visible-light camera systems mounted to the side of each STEREO spacecraft to image along the Sun-Earth line from elongations of 4–88.7° (Figure 2.8). This has provided several new opportunities for CME research, notably the ability to track their evolution as they propagate through the inner heliosphere and potentially impact at Earth or one of either STEREO spacecraft which allows a comparison of in-situ data and white-light imagery of CMEs. The basic design




Figure 2.7: Diagram of the COR2 coronagraph in the STEREO/SECCHI suite, reproduced from Howard et al. (2000). The measurements are specified in millimetres.




Figure 2.8: Diagram of the heliospheric imagers HI-1/2 in the STEREO/SECCHI suite, reproduced from Howard *et al.* (2000).

of the HI comprises a number of occulting baffles that achieve the required level of light rejection for imaging CMEs of intensities about two orders of magnitude weaker than the F-corona (Figure 2.9). The final images are 2×2 binned onboard to 1024×1024 pixels, subtending an angle of 70 arcseconds per pixel for HI1 and 4 arcminutes per pixel for HI2. In order to sufficiently image the low intensity CME signal over the stellar background and zodiacal light, a series of short exposures are summed together since long exposures are dominated by cosmic ray hits. The combination of summing (30 images for HI1, and 99 images for HI2) and 2×2 binning increases the signal-to-noise ratio by about $14 \times$.




Figure 2.9: The intensity profile of a CME compared to the K & F coronae observed at elongations up to 90°, and the corresponding fields-of-view of the Heliospheric Imagers (HI1/2), reproduced from Howard *et al.* (2000).

2.3 CCD Detectors

A charge-coupled device (CCD) is used in the LASCO and SECCHI instruments for detecting the incident photons and converting them to a digital output to generate images. Essentially a CCD converts light into electrons which are read and converted into numeric values used to display image intensities. The CCD is a small silicon chip divided into a grid of cells, or pixels. The electrons in the silicon atoms lie in discrete energy bands. In the ground state the outermost electrons lie in the valence band and can be excited to the conduction band by the




Figure 2.10: Illustration of a thick front-side illuminated CCD. *Image credit:* www.inq.iac.es.

absorption of a photon, via the photoelectric effect, leaving behind a 'hole'. In a CCD an electric field is introduced to prevent recombination of the electron-hole pair. Thus an electric charge is accumulated proportional to the light intensity at that location. The charge is read out pixel by pixel to a charge amplifier which converts it to a voltage, then this voltage is digitised and stored in memory.

A thick front-side illuminated CCD (Figure 2.10) is cheap to produce, but because photons are incident at the surface electrodes they can be reflected or absorbed which gives low quantum efficiency (a measure of the percentage of photons detected: $QE = N_e/N_{\nu}$). The LASCO/C2 and C3 detectors are front-side illuminated Textronix CCDs that have a quantum efficiency of about 0.3-0.5 in the 500 to 700 nm spectral range. They are 1024×1024 pixels in size, each pixel being a square measuring 21 μ m on a side.

To increase the quantum efficiency back-side illumination is used so the electrodes do not obstruct the photons. But the silicon in a back-side illuminated CCD must be chemically etched down (thinned) to a thickness of about 15 μ m, which is an expensive process (Figure 2.11). Silicon also has a high refractive




Figure 2.11: Illustration of a thinned back-side illuminated CCD. *Image credit:* www.ing.iac.es.

index leading to strong photon reflection. It must therefore be coated with an anti-reflective material with a refractive index less than that of silicon (3.6) and preferably with an optical thickness of 1/4 at a chosen wavelength of 550 nm (close to the middle of the optical spectrum). Hafnium dioxide is regularly used to significantly reduce the reflectivity of the CCD. Due to their high quantum efficiency, almost all current astronomical CCDs are thinned and back-side illuminated. Each of the SECCHI instruments uses a back-side illuminated E2V 42–40 CCD detector, that has a quantum efficiency of roughly 0.8 at 500 nm, 0.88 at 650 nm, 0.64 at 800 nm, and 0.34 at 900 nm. They are 2048×2048 pixels in size, each measuring $13.5~\mu m$ on a side. This CCD has an operational temperature range of 153-323 K ($-120-50^{\circ}$ C).

Sources of noise in CCD imaging must be noted when performing image analysis. Thermal noise, or dark current, is due to thermal excitations of electrons in the CCD. A dark frame may be generated to correct for thermal noise by taking a closed shutter exposure of some known duration to study the effects on the resultant image, though this form of noise is minimal for space-borne instruments operating at temperatures of $\lesssim 200$ K.

While such temperatures are ideal for detecting faint coronal light they also mean the CCD is sensitive to energetic particles. Hot pixels can result from these energetic particles or cosmic rays causing ionisation in the silicon, since the resulting free electrons are indistinguishable from photo-generated ones. CCD read-out noise can occur when charge is converted to voltage since electronic amplifiers are not perfect. A high charge transfer efficiency is also important during shift operations in the read-out process to minimise count errors.

Calibrations of CCD images must be performed to remove imperfections. CCDs are not always linear (measuring one count for one photon incident). A flat-field calibration removes variations in sensitivity across the surface of the CCD, due to silicon or manufacturing defects and vignetting effects. Flat-field images are normally generated in the lab by taking an exposure when the CCD is evenly illuminated by a light source, and dividing this into future images for linearity.

Similar to dark frames, bias frames may also be generated. A bias frame is a zero duration exposure taken with no light incident on the CCD (the shutter remains closed). Thus structures which appear in bias frames are as a result of defects in the CCD electronics and must be removed from future images.

The charge capacity of a CCD pixel is limited and when full it can overflow, leading to blooming. While this is somewhat unavoidable when taking long exposures, especially if a bright star or comet comes into view for example, most CCD design ensures blooming only occurs in one direction; normally vertically with respect to the orientation of the CCD imager.

2.4 Coordinate Systems

The pixel coordinates from the CCD images must be transformed to the relevant coordinate system for studying and interpreting observations, especially when comparing images from multiple viewpoints (such as STEREO and SOHO as discussed in Section 5.3.1). First the pixel coordinates $(p_1, p_2, p_3, ...)$ must be transformed to intermediate world coordinates $(x_1, x_2, x_3, ...)$, meaning they are converted into the relevant units (e.g. metres or arcseconds) but are not necessarily corrected for the reference point of the observations nor geometric or projection effects:

$$x_i = s_i \sum_{j=1}^{N} m_{ij} (p_j - r_j)$$
 (2.3)

where s_i is the scale function, N is the number of axes, m_{ij} is the transformation matrix, and r_j is the reference pixel (Calabretta & Greisen, 2002). Then these can then be transformed into one of the Sun-centred coordinate systems described below.

2.4.1 Heliographic Coordinates

Features on the Sun are located by the coordinates of latitude Θ and longitude Φ with respect to the solar equator and rotational axis. In the Stonyhurst approach, the zero point of longitudinal measurements is set at the intersection of the solar equator and central meridian as seen from Earth (Figure 2.12a). In the Carrington approach, the central meridian is fixed to its observation on 9 November 1853, and the rotations since then are counted and labelled as the Carrington rotation number (Carrington, 1863).




Figure 2.12: Schematics of two Sun-centred coordinate systems, reproduced from Thompson (2006). (a) Stonyhurst heliographic coordinates commonly used to locate features on-disk. (b) Heliocentric-cartesian coordinates commonly used for spatially localising features in the vicinity of the Sun.

2.4.2 Heliocentric Coordinates

Heliocentric coordinates specify the location of a feature in space with respect to the centre of the Sun. The Heliocentric-Cartesian coordinate system (x, y, z) has the z-axis from Sun-centre along the Sun-observer line, the y-axis is perpindicular to this and lies in the plane containing solar north, and the x-axis is perpindicular to both y and z and increases towards solar west (Figure 2.12b). The Heliocentric-Radial coordinate system shares the same z-axis but measures features in cylindrical coordinates with radial distance ρ from the z-axis, and position angle ψ counterclockwise from solar north.

With observations from the STEREO/SECCHI instrument suite centred on the Sun and extending out along the full Sun-Earth line, the optimal coordinate systems are the heliocentric coordinates, which can be defined in three possible manners.

2.4.2.1 Heliocentric Earth Equatorial (HEEQ)

The HEEQ system is closely related to the Stonyhurst heliographic coordinates with the z-axis parallel to the solar rotational axis, and the x-axis towards the intersection of the solar equator and central meridian as seen from Earth, obtained by the following transformations:

$$X_{HEEQ} = r \cos \Theta \cos \Phi$$

 $Y_{HEEQ} = r \cos \Theta \sin \Phi$ (2.4)
 $Z_{HEEQ} = r \sin \Theta$

2.4.2.2 Heliocentric Earth Ecliptic (HEE)

The HEE system has the x-axis towards the Earth from Sun centre, and the z-axis is perpindicular to the plane of Earth's orbit around the Sun called the ecliptic, and the y-axis perpendicular to both x and z.

2.4.2.3 Heliocentric Aries Ecliptic (HAE)

The HAE system has the x-axis towards the First Point of Aries (the direction to the point of intersection between Earth's equatorial plane and the plane of the ecliptic), the z-axis perpendicular to the ecliptic, and the y-axis perpendicular to both x and z.

2.4.3 Helioprojective Coordinates

When considered on a large scale it is more intuitive to take the projection of the Heliocentric coordinates onto the celestial sphere (an imaginary sphere of




Figure 2.13: Schematic of the celestial sphere and ecliptic plane. The celestial equator is a projection of the Earth's equator, and the ecliptic is a projection of the Earth's orbit about the Sun.

arbitrarily large radius, centred on the Earth such that all observations may be considered as projections upon it). This results in the Helioprojective-Cartesian coordinates $(\theta_x, \theta_y, \zeta)$:

$$\theta_x \approx \left(\frac{180^{\circ}}{\pi}\right) \frac{x}{d} , \quad \theta_y \approx \left(\frac{180^{\circ}}{\pi}\right) \frac{y}{d} , \quad \zeta = D_{\odot} - d$$
 (2.5)

where d is the distance between the observer and the feature, and D_{\odot} is the distance between the observer and Sun centre. This similarly results in the Helioprojective-Radial coordinates $(\delta_{\rho}, \psi, \zeta)$:

$$\delta_{\rho} \equiv \theta_{\rho} - 90^{\circ} \text{ where } \theta_{\rho} \approx \left(\frac{180^{\circ}}{\pi}\right) \frac{\rho}{d}$$
 (2.6)

Chapter 3

Detecting and Tracking CMEs

In coronagraph images CMEs are observed as outwardly moving regions of stronger brightness intensities than the background corona. Different approaches to threshold the intensity of CMEs in these images have been employed in order to detect their appearance and track their motion through the field-of-view, leading to a cataloguing of their kinematics and morphology. However these techniques suffer several drawbacks and as such different catalogues can vary significantly in their description of events. We introduce a method of multiscale analysis to overcome certain drawbacks of previous detection and tracking methods. In multiscale decompositions of images noise and small-scale features are removed to leave only larger-scale features of interest such as CMEs. This allows them to be tracked through the image sequences in order to determine their changing kinematics and morphology (see Chapter 4). Unfortunately coronal streamers tend to appear on similar size scales to CMEs, making their automatic detection difficult. Streamers do, however, tend to remain static on timescales comparable to CME propagation through the field-of-view and contain much less angular information

than the typically curved structure of CMEs, so they may be removed through spatio-temporal filtering of multiscale CME images. This chapter discusses the previous CME detection catalogues, and outlines our use of new methods of multiscale filtering to detect the CME edges in single images. We discuss our efforts to extend this to an automated CME detection algorithm. We also outline an ellipse characterisation of the CME front for study.

3.1 CME Detection Catalogues

Current methods of CME detection have their limitations, mostly since these diffuse objects have been difficult to identify using traditional image processing techniques. These difficulties arise from the varying nature of the CME morphology, the scattering effects and non-linear intensity profile of the surrounding corona, the presence of coronal streamers, and the addition of noise due to cosmic rays and solar energetic particles (SEPs) that impact the coronagraph detectors. The images are also prone to numerous instrumental effects and possible data dropouts. The following standard preprocessing methods are usually applied to optimise the images for CME studies. The coronagraph images are normalized with regard to exposure time in order to correct for temporal variations in the image statistics. A filter may be applied to remove pixel noise, for example to replace hot pixels with a median value of the surrounding pixel intensities, or to reduce the effects of background stars in the image. A correction for vignetting effects and/or lens distortion may be applied to the images. A background subtraction may also be applied, obtained from the minimum of the daily median pixels across a time span of a month. The occulting disc is normally masked, along




Figure 3.1: Raw (left) and pre-processed image (right) of a CME observed by LASCO/C2 on 1 April 2004. The pre-processing includes normalising the image statistics, taking a background subtraction, and masking the occulter disk. The white circle (right) indicates the relative size and position of the Sun behind the occulter.

with any data drop-outs in the images. These steps lead to a clear improvement in the image quality for CME study (Figure 3.1).

3.1.1 CDAW

The CME catalogue hosted at the Coordinated Data Analysis Workshop (CDAW¹) Data Center grew out of a necessity to record a simple but effective description and analysis of each event observed by SOHO/LASCO (Gopalswamy et al., 2009b). The catalogue is wholly manual in its operation, with a user tracking the CME through C2 and C3 running-difference images and producing a height-time plot of each event. A linear fit to the height-time profiles provides a 1st-order estimate for the plane-of-sky velocity, and a quadratic fit then provides a 2nd-


¹http://cdaw.gsfc.nasa.gov/CME_list

order velocity fit and an acceleration for the event. The central position angle and angular width of the CME are also deduced from the images, and the event flagged as a halo if it spans 360° , partial halo if it spans $\geq 120^{\circ}$, and wide if it spans $\geq 60^{\circ}$. The catalogue itself lists each CME's first appearance in C2, central position angle, angular width, linear speed, 2nd-order speed at final height, 2nd-order speed at 20 R_{\odot} , acceleration, mass, kinetic energy, and measurement position angle (the angle along which the heights of the CME are determined). While the human eye is supremely effective at distinguishing CMEs in coronagraph images, errors may be introduced to the manual cataloguing procedure through the biases of different operators; for example, in deciding how the images are scaled, where along the CME the heights are measured, or whether a CME is even worth including in, or discarding from, the catalogue. In an effort to overcome such biases, different automated catalogues have been developed to perform robust CME detections over large data-sets. This is also of great benefit for future missions where the data rate is expected to be too high for manual cataloguing to remain feasible.


3.1.2 CACTus

The Computer Aided CME Tracking catalogue (CACTus¹; Robbrecht & Berghmans, 2004) was the first automated CME detection algorithm, in operation since 2004. It is based upon the detection of CMEs as bright ridges in time-height slices (t, r) at each angle θ around a coronagraph image. The images are preprocessed as standard, then a running-difference technique is applied and each image unwrapped into Sun-centred polar coordinates (r, θ) , rebinned, and the C2 and C3

¹http://sidc.oma.be/cactus/



(a) The (t, r) slice for a given angle θ , with a mirrored illustration of the resulting CME intensity ridge detections.



(b) Left: the (t, r) slice for a given angle θ , with an example ridge drawn from onset time t_0 with duration Δt across the field-of-view from r_{min} to r_{max} . Right: the corresponding accumulator space $(t_0, \Delta t)$ where the ridge will appear as a point with a magnitude corresponding to the ridge intensity. This modified Hough transform is used to threshold the most significant ridges in the slice, automatically detecting the CME in the coronagraph image.

Figure 3.2: The top image (a) shows the detection of ridges in the (t, r) stacks of the CACTus catalogue, through the use of the Hough transform detailed in the bottom image (b), reproduced from Robbrecht & Berghmans (2004).

fields-of-view combined. These are then stacked in time, and for each angle the corresponding (t, r) slice undergoes a modified Hough transform for detecting intensity ridges across it. This works by parameterising the (t, r) slice by the variables t_0 and Δt , corresponding to the coordinate intersection point with the time axis, and the distance along the time axis respectively (together called the accumulator space; see Figure 3.2). So the equation of a line corresponding to an intensity ridge in the slice is given by:

$$r = \frac{r_{max} - r_{min}}{\Delta t} (t - t_0) + r_{min} \tag{3.1}$$

Thresholding the most significant ridges in the resultant accumulator space filters out the progression of CMEs, with the variables for each ridge characterised by onset time t_R , the velocity v_R ($\sim \frac{1}{\Delta t}$), for angle θ_R , to give a characteristic variable $I_R = (v_R, \theta_R, t_R)$. A 3D scatter plot (v, θ, t) of all detected ridges I_R is then integrated along the v-direction to identify clusters in the resulting (θ, t) map which illustrates the angular span and duration time of the detected CMEs in the coronagraph data. A median velocity across the angular span is quoted as the CME speed.

The running-difference cadence, the ridge intensity threshold, and the imposed limit on how many frames a CME may exist (and indeed the definition of a CME) all affect how successful the detection can be. However, Robbrecht & Berghmans (2004) show the algorithm to be robust in reproducing well the detections of a human user by direct comparison with the CDAW catalogue. The main drawback of the CACTus catalogue for studying CMEs is the imposed zero acceleration of the detection algorithm, since the Hough transform thresholds the ridges as

straight lines whose slopes provide a constant velocity. The velocity itself may also be underestimated since it is a median across the span of the CME. The angular spans are possibly over-estimated since side outflows in the images are enhanced by the running-difference and may also include streamer deflections. It is also difficult to distinguish when one CME has fully progressed from the field-of-view and another CME has entered it, so in some cases trailing portions of a CME are detected as separate events.

3.1.3 **SEEDS**

The Solar Eruptive Event Detection System (SEEDS¹; Olmedo *et al.*, 2008) is an automated CME detection algorithm for tracking an intensity thresholded CME front in running-difference images from LASCO/C2. The images are preprocessed as standard, unwrapped into Sun-centred polar coordinates (r, θ) , and a normalised running-difference technique is applied using the following equation:

$$u_i = \left[n_i - n_{i-1} \left(\frac{\bar{n}_i}{\bar{n}_{i-1}} \right) \right] \frac{\alpha}{\Delta t}$$
 (3.2)

where u_i is the running-difference image, \bar{n} is the mean of the pixels in the entire field-of-view of the image n, Δt is the time difference between images (in minutes), α is a constant set to approximately the smallest time difference (Δt) between any image pair, where i is the current image and i-1 the preceding image. This normalised difference ensures that the mean of the new image (u_i) will effectively be zero.

The pixel intensities (positive values only) are then summed along angles and

¹http://spaceweather.gmu.edu/seeds/




Figure 3.3: Example of the SEEDS CME detection and height determination, reproduced from Olmedo et al. (2008). (a) shows the running-difference image unwrapped into Sun-centre polar coordinates, showing a CME observed by LASCO/C2 on 12 September 2002. The black line distribution across the image represents the positive value intensity count along each angle, and the two vertical black lines mark the angular span at one standard deviation above the mean intensity. (b) shows the new angular span following the region growing technique. (c) shows the intensity within the angular span averaged across heights, and the 'Half-Max-Lead' is taken as the CME height in the image.

thresholded at a certain number of standard deviations above the mean intensity: $\mu + N\sigma$ (c.f. equation 3.12) as in Figure 3.3a. This determines the 'core angles' of the CME, and a region growing technique based on a secondary threshold of intensities in the rest of the image is applied to open the angular span to include the full CME (Figure 3.3b). Issues arise when streamer deflections occur that will offset the region growing technique and overestimate the CME angular width. An intensity average across the angles within the span of the CME is then determined, and where the forward portion of this intensity profile equals half its maximum value is taken as the CME height (Figure 3.3c). The velocity and acceleration are determined from the heights through consecutive images and these results are output with the CME position angle and angular width in the SEEDS catalogue.

Along with the issues of streamer deflections and the tracking being limited to C2 images, the choice of the 'Half-Max-Lead' as the CME height is dependant on the overall CME brightness, and thus any brightness changes as the CME propagates will affect this measurement. This would add to the error on the height-time profile which, along with the error in time as a result of the running-difference technique, makes it difficult to accurately determine the velocity and acceleration.

3.1.4 ARTEMIS

The Automatic Recognition of Transient Events and Marseille Inventory from Synoptic maps (ARTEMIS¹; Boursier *et al.*, 2009b) is an automated CME detection algorithm that works by identifying signatures of transients in synoptic maps.

¹http://www.oamp.fr/lasco/

These maps are generated as (t, θ) slices for specific heights r in the coronagraph images of LASCO/C2. The images are prepared through the standard preprocessing steps. Then at a specific height (e.g. $r = 3 \text{ R}_{\odot}$) the intensity is plotted across all angles θ for each image through time t with transient events appearing as vertical streaks through the more persistent streamer intensities (Figure 3.4). A method of image filtering and intensity thresholding is applied to distinguish the streaks in the synoptic map, and image segmentation then discards small features and closes off regions-of-interest (ROIs) to produce a binary map of the streaks. Specific parameters of these streaks are also computed, such as their total radiances, areas and centres of gravity. Merging with high-level knowledge helps to associate ROIs of the same CME if three criteria are met: the difference between the x-coordinates of the centres of two ROIs differs by less than two pixels; the difference between the y-coordinates of their centres differs by less than 60 pixels (corresponding to a 60° angular span); and the ratio of their radiances calculated at their centres (on the original synoptic map) ranges from 0.25 to 4. The result is a binary CME detection map in (t, θ) space for different heights in the corona: 3, 3.5, 4, 4.5, 5 and 5.5 R_{\odot} .

With the CME detections in place, estimates of the velocity may be made. A first estimate is taken by testing a range of constant velocities $50-2,000~\rm km~s^{-1}$ to determine which best matches the shifting of the CME detection in synoptic maps at subsequent heights through the corona. The binary maps are shifted by an amount corresponding to velocity steps of 10 km s⁻¹, such that the one which provides the maximum pixel value (with a minimum limit of 3) indicates the best velocity estimate of the event. A second estimate is taken by cross-correlating the detected CME ROIs on the original synoptic maps at 3 R_{\odot} and 5.5 R_{\odot}




Figure 3.4: An example of how the synoptic maps are generated for the ARTEMIS catalogue, reproduced from Boursier et al. (2009b). At a chosen height in the coronagraph image an annulus is unwrapped (indicated with the dashed line and blue square, circle and triangle) and these are then stacked together to illustrate how the intensity at that height changes through time. Vertical streaks represent transient events occurring on smaller time-scales than the more persistant streamers in the images.

and inspecting the intensity shift in time (pixel shift in x-direction) to obtain the velocity estimate. A third estimate is taken by similar cross-correlation but specifically on each individual line of the ROIs to obtain a distribution of velocities across the angular span of the CME, the median of which is taken as the actual velocity. Boursier $et\ al.\ (2009b)$ compare histograms of the three different velocity estimates for the ARTEMIS CME detections over a twelve year interval and find that, globally, the three estimates are highly consistent with each other.

ARTEMIS is limited to the C2 field-of-view and it provides kinematics only in the $3-5.5~\rm R_{\odot}$ range. The velocity determinations themselves are not specific to either the CME front nor other identifiable feature, and carry all the inaccuracies resulting from the image rebinning, intensity averaging, filtering and segmentation techniques in generating the final detection masks.

Due to the drawbacks of each of the catalogues above, the motivation exists to study the kinematics and morphology of CMEs with as great an accuracy as possible in order to better compare with theory. To this end we outline below the application of multiscale analysis to remove small scale noise/features and enhance the larger scale CME in single coronagraph frames, allowing the CME front edges to be detected and a geometrical characterisation applied to study its propagation with increased accuracy for deriving the kinematics and morphology.

3.2 Multiscale Filtering

In this section a new multiscale method of analysing CMEs is described. The use of multiscale methods in astrophysics have proven effective at denoising spectra and images (Fligge & Solanki, 1997; Murtagh et al., 1995), analysing solar active region evolution (Hewett et al., 2008), and enhancing solar coronal images (Stenborg & Cobelli, 2003; Stenborg et al., 2008). A particular application of multiscale decompositions uses high and low pass filters convolved with the image data to exploit the multiscale nature of the CME (Young & Gallagher, 2008). This highlights its intensity against the background corona as it propagates through the field-of-view, while neglecting small scale features (essentially denoising the data). It also leads to the use of non-maxima suppression to trace the edges in the CME images, and Young & Gallagher (2008) show the power of multiscale methods over previous edge detectors such as Roberts and Sobel. With these methods for defining the front of the CME we can characterise its kinematics (position, velocity, acceleration) and morphology (width, orientation) in coronagraph images. Multiscale analysis also has the benefit of working on independent images without any need for differencing, so the temporal errors involved are on the order of the exposure time of the instrument (\sim a few seconds).

The fundamental idea behind wavelet analysis is to highlight details apparent on different scales within the data. An example of this is the removal of noise from images, which tends to occur only on the smallest scales. Wavelets have benefits over previous methods (e.g. Fourier transforms) because they are localised in space and are easily dilated and translated in order to operate on multiple scales, the basic equation being:

$$\psi_{a,b}(t) = \frac{1}{\sqrt{b}} \psi(\frac{t-a}{b}) \tag{3.3}$$

where a and b represent the shifting (translation) and scaling (dilation) of the

mother wavelet ψ which can take several forms depending on the required use.

We explore a method of multiscale decomposition in 2D through the use of low and high pass filters; using a discrete approximation of a Gaussian θ and its derivative ψ respectively (Stenborg & Cobelli, 2003). Since $\theta(x,y)$ is separable, i.e. $\theta(x,y) = \theta(x)\theta(y)$, we can write the wavelets as the first derivative of the smoothing function:

$$\psi_x^s(x,y) = s^{-2} \frac{\partial \theta(s^{-1}x)}{\partial x} \theta(s^{-1}y)$$
(3.4)

$$\psi_y^s(x,y) = s^{-2}\theta(s^{-1}x)\frac{\partial\theta(s^{-1}y)}{\partial y}$$
(3.5)

where s is the dyadic scale factor such that $s = 2^j$ where j = 1, 2, 3, ..., J. Successive convolutions of an image with the filters produces the scales of decomposition, with the high-pass filtering providing the wavelet transform of image I(x, y) in each direction:

$$W_x^s I \equiv W_x^s I(x, y) = \psi_x^s(x, y) * I(x, y)$$
 (3.6)

$$W_y^s I \equiv W_y^s I(x, y) = \psi_y^s(x, y) * I(x, y)$$
 (3.7)

Akin to a Canny edge detector (Young & Gallagher, 2008), these horizontal and vertical wavelet coefficients are combined to form the gradient space $\Gamma^s(x,y)$ for each scale:

$$\Gamma^s(x,y) = \left[W_x^s I, \ W_y^s I \right] \tag{3.8}$$

The gradient information has an angular component α and a magnitude (edge




Figure 3.5: Top left, the horizontal detail, and top right, the vertical detail from the high-pass filtering at one scale of the multiscale decomposition (called the rows and columns respectively). Bottom left, the corresponding magnitude (edge strength) and bottom right, the angle information $(0-360^{\circ})$ taken from the gradient space, for a CME observed in LASCO/C2 on 1 April 2004 (Byrne *et al.*, 2009).




Figure 3.6: The vectors plotted represent the magnitude and angle determined from the gradient space of the high-pass filtering at scale 3. The CME of 2004 April 1 shown here is highlighted very effectively by this method (Byrne *et al.*, 2009).

strength) M:

$$\alpha^{s}(x,y) = \tan^{-1}\left(W_{y}^{s}I / W_{x}^{s}I\right) \tag{3.9}$$

$$M^{s}(x,y) = \sqrt{(W_{x}^{s}I)^{2} + (W_{y}^{s}I)^{2}}$$
 (3.10)

The resultant horizontal and vertical detail coefficients, and the magnitude and angular information are illustrated in Figure 3.5.

At a particular scale the signal-to-noise ratio of the CME is highest and this is the optimum scale for determining the edges in the image. The angular component α of the gradient specifies a direction which points across the greatest intensity change in the data (an edge). A threshold is specified with regard to this gradient direction in order to chain pixels along maxima to highlight the edges. The changes in magnitude and angular information may then be imple-

mented in a spatiotemporal filter for distinguishing those edges corresponding to the CME only. Overlaying a mesh of vector arrows on the data shows how the combined magnitude and angular information illustrate the progression of the CME. Each vector is rooted on a pixel in the gradient space, and has a length corresponding to the magnitude M with an angle from the normal α (Figure 3.6). Using this information, it becomes possible to create a specific detection mask which is used to pull out the edges along the CME front to study its propagation (as done for a sample of events in Chapter 4). However, for cases of faint CMEs or strong streamer deflections the filter is presently limited by exploiting the information from only one scale and ignoring all other scales, meaning it currently often requires the user to remove/include certain edges that the algorithm has mistakenly retained/discarded. Extending the algorithm to work on more than one scale may help alleviate this issue in order to develop a fully automated CME detection and characterisation routine, as outlined in the following section.

3.3 Automated Multiscale CME Detection

For the most part, CMEs exist on size scales larger than noise and any small scale features in coronagraph images that are redundant for studying CME propagation. This fact has led to the development and implementation of multiscale decompositions that highlight the CME in images from SOHO/LASCO and STEREO/SECCHI (Byrne *et al.*, 2009; Stenborg & Cobelli, 2003; Young & Gallagher, 2008). However, coronal streamers (plasma outflows from open magnetic field regions on the Sun) can persist through coronagraph images with significant brightness intensities and tend to appear on similar scales as the CME in

multiscale image analysis. If a CME propagates through an image with a strong streamer present, it becomes difficult to distinguish the two features by intensity thresholding alone, and this is one reason why differencing techniques have been widely used in CME analysis. In an effort to move away from differencing and the large errors involved in the subtraction of images from each other, since the goal is to obtain kinematics with the greatest precision, we endeavour to separate CME and streamer features from one another using multiscale methods alone. These efforts involve exploiting the angular distribution that exists across a curved CME front compared to the more linear streamers in single independent images. To do this, the coronagraph images must first be normalised for their radial gradient in intensity, since the drop-off across the field-of-view is too steep to effectively segment a single entire streamer from the inner to outer edge of an image. This can serve to enhance the noise at the edge of the images, but this is again suppressed by the multiscale analysis. Occasions when the CME propagates directly in front of or behind a streamer remain problematic, as do strong streamer deflections that can occur when a CME propagates into or expands alongside a streamer.

3.3.1 Normalising Radial Graded Filter

Since the intensity drop-off of the corona is large, falling from approximately $10^{-6}-10^{-9}$ B_{\odot} across heights of 1-6 R_{\odot} (Kimura & Mann, 1998), a method for radially normalising coronagraph images to enhance features across this steep intensity gradient was developed by Morgan *et al.* (2006). It works by normalising




Figure 3.7: A normalised, background subtracted, LASCO/C2 image (left) of a CME on 1 April 2004, and the resulting NRGF image (right). The image radial intensity is scaled such that structure along streamers and the CME becomes visible across the field-of-view

the intensity in radial coordinates of the image according to the equation

$$I'(r,\phi) = \frac{I(r,\phi) - I(r)_{\langle\phi\rangle}}{\sigma(r)_{\langle\phi\rangle}}$$
(3.11)

where $I'(r,\phi)$ is the processed and $I(r,\phi)$ is the original intensity at height r and position angle ϕ , and $I(r)_{<\phi>}$ and $\sigma(r)_{<\phi>}$ are the mean and standard deviation of intensities calculated over all position angles at height r. Figure 3.7 shows the result when the NRGF is applied to a LASCO/C2 image of the 1 April 2004 CME.

The multiscale decomposition introduced in Young & Gallagher (2008) provides magnitude and angular information of the edges in the image. This information is combined to chain the strongest edges within the image on a scale which provides an optimal signal-to-noise ratio for studying the CME. Byrne *et al.*




Figure 3.8: A chosen scale of the decomposed NRGF image provides a magnitude image of the edge strengths displayed on the left, which is thresholded at one standard deviation of the mean intensity to obtain contoured regions of interest that could contain a CME (sample contours indicated in red). As shown, the streamers have edges which appear on the same scale as the CME edges in this image. The angular information from the decomposition is displayed on the right, and the contoured regions of interest overlaid for comparison. The grey scale indicates angles from $0-360^{\circ}$ and it is clear that streamers tend to have a linear colour scale while the CME has a gradient of colours across the scale.

(2009) obtain the CME front edges in this manner, which are then used to fit an ellipse to characterise the CME propagation in the image sequence. In order to automate the algorithm, thresholds on the magnitude information (e.g. CME edges appear on larger scales than noisy features) and angular information (e.g. CME edges appear more curved than streamer edges) were investigated. The thresholding is strengthened by the inclusion of more than one scale in localising the CME and distinguishing it from the streamers, detailed below.

3.3.2 Thresholding

The magnitude information corresponds to the strength of the edges in the image, and so can be thresholded at several scales to discard the small scale noise. For the NRGF image (Figure 3.7), a hard threshold T is set at one standard deviation σ of the mean μ of the image intensity I to contour out regions of interest that may be a CME according to the equations:

$$T = \mu_I + \sigma_I = \underbrace{\frac{1}{n} \sum_{i=1}^n x_i}_{\text{mean}} + \underbrace{\sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}}_{\text{standard deviation}}$$
(3.12)

where x_i are the pixel intensity values of the image, with a mean of \bar{x} . The left image of Figure 3.8 illustrates this thresholding with a sample of contoured regions (outlined in red) on the multiscale decomposition of a CME observed in LASCO/C2 on 1 April 2004.

It is apparent from the right of Figure 3.8 that the CME will contain edges whose normals are widely distributed across $0-360^{\circ}$ compared to the more linear, radially directed, streamer edges. The angular distribution of each region is determined and then normalised and folded into $0-180^{\circ}$ range, centred on 90° , due to the symmetry of the edge normals. This is illustrated for four selected contour regions in Figure 3.9. The resulting angular distributions are then thresholded with regard to their median value, since the distribution of angles across the CME will be wider and have a higher median value than for a distribution of angles along a streamer.

This thresholding is repeated across four scales of the multiscale decomposition, neglecting smaller scales dominated by noise and larger scales that are




Figure 3.9: Left: four contoured regions (at one standard deviation of the mean image intensity) highlighted on the magnitude information from the multiscale decomposition of the 1 April 2004 CME. Right: the corresponding angular distribution of each region, normalised and folded into the $0-180^{\circ}$ range (centred on 90°). The angular distribution may be thresholded with respect to its median value to distinguish regions corresponding to CMEs from those along streamers.

overly smoothed. Assigning a score to the regions that may contain a CME at each scale, it becomes possible to build a detection mask as in Figure 3.10.

The scoring system is chosen arbitrarily to best work with the chosen thresholds, and these may be changed and refined as an analysis of more CMEs is done. For example, the current thresholds from working on a sample of ~ 10 CMEs are as follows:

- 1. The magnitude information is thresholded at one standard deviation (1σ) of the mean intensity across the image.
- 2. The 15 largest contoured regions across the image are investigated (there are rarely more than ~ 5 streamers of similar intensity to a CME, and we allow for disjointed contours along structures).




Figure 3.10: The resulting CME detection mask from combining the thresholded regions of strongest magnitude and angular distribution at four scales of the decomposition.

- 3. If the median angular value is > 20% of the angular distribution peak then the region is deemed a CME and assigned a score of 3 (the pixels in that region of the mask are given the value 3). If it is > 10% the score is 2 (potential CME structure), or > 5% the score is 1 (weak CME structure or portion thereof).
- 4. The final CME mask through the combination of scores at each scale results in a dominant region that localises the CME front in the image and can be used to characterise the front, or input into a spatio-temporal filter if subsequent CME images are available in order to refine the masked region whenever streamers are still present.

The resultant set of detection masks for three of the frames of the CME observed on 1 April 2004 are shown in Figure 3.11.




Figure 3.11: The NRGF (left) and resulting detection masks (middle and right) for different frames of a CME observed by LASCO/C2 at times of 00:00 UT (top), 00:40 UT (middle) and 01:20 UT (bottom) on 2 April 2004. The location of the CME front is highlighted very efficiently by this method, although the detection masks may contain artefacts of the chosen thresholds which must be discarded when characterising the CME front.

3.3.3 Faint CMEs and Streamer Interactions/Deflections

Due to the nature of the hard thresholds in place on the magnitude and angular information, there are problems arising when the algorithm mistakenly disregards a CME or includes a streamer, or portions thereof. Firstly, if a CME is faint enough that the intensity falls below the 1σ magnitude threshold, it will not be detected as a region of interest in the image. Secondly, if the CME interacts with a streamer, the two features may be contoured together and this will skew the angular distribution and affect the detection mask. And thirdly, if the CME causes a significant streamer deflection, it will lead to a wider distribution of angles along the streamer and the algorithm may thus detect it as part of the CME. This is why the above scoring system was introduced in an effort to minimise these effects, which are highlighted in Figure 3.12 for a CME observed on 23 April 2001. The event is too faint compared with the streamers across it for it to be easily distinguished in the image, and parts of the streamers are then mistakenly included in the final detection mask. This is where the current algorithm requires a user to specify which parts of the edges correspond to the CME for characterisation. Such limitations in current wavelet analysis of CMEs may be overcome by extending these algorithms to work with ridgelets or curvelets that better suit the curved form of a typical CME front as discussed in Gallagher et al. (2010).

3.4 Characterising the CME Front

Using a model such as an ellipse to characterise the CME front across a sequence of images, has the benefit of providing the kinematics and morphology of a mov-




Figure 3.12: Example of the difficulty in detecting the faint CME observed by LASCO on 23 April 2001. If the CME is far-sided and/or of low intensity it becomes difficult to threshold its edges in the image compared to the coronal streamers.

ing and/or expanding structure. The ellipse's multiple parameters, namely its changeable axes lengths and tilt angle, is adequate for approximating the varying curved structures of CMEs. Chen et al. (1997) suggest an ellipse to be the two-dimensional projection of a flux rope, and Krall & St. Cyr (2006) use ellipses to parameterise CMEs and explore their geometrical properties. It also serves as the observed projection of the base of the cone model applied to CME images (Xie et al., 2004; Xue et al., 2005; Zhao et al., 2002). We fit ellipses to the points determined to be along the CME front by means of a Levenberg-Marquardt least squares algorithm. A kinematic analysis then provides height, velocity and acceleration profiles; while the ellipse's changing morphology provides the inclination angle and angular width (see the example in Figure 3.14). Measuring these properties in the observed data is vitally important for accurate comparison with theoretical models.

Following Schrank (1961), we may determine the polar equation of an inclined ellipse by starting with the standard equation for an ellipse with centre point (x_0, y_0) , semimajor axis a, and semiminor axis b:

$$\frac{(x-x_0)^2}{a^2} + \frac{(y-y_0)^2}{b^2} = 1 (3.13)$$

This is written in polar coordinates by $x = \rho \cos \omega$, $y = \rho \sin \omega$ and centred on the origin $(x_0 = 0, y_0 = 0)$ to give:

$$\frac{\rho^2 \cos^2 \omega}{a^2} + \frac{\rho^2 \sin^2 \omega}{b^2} = 1 \tag{3.14}$$

where ρ is a radial line from the centre to any point on the ellipse, at an angle ω





Figure 3.13: Ellipse inclined at angle γ , with semimajor axis a, semiminor axis b, and radial line ρ inclined at angle ω to the semimajor axis.

to the semimajor axis a (Figure 3.13). Allowing for a tilt angle γ on the ellipse, we may define $\omega' = \omega + \gamma$ to obtain:

$$\rho^2 = \frac{a^2b^2}{\left(\frac{a^2+b^2}{2}\right) - \left(\frac{a^2-b^2}{2}\right)\cos(2\omega' - 2\gamma)}$$
(3.15)

This gives a first approximation which can then be used to iteratively float the ellipse parameters until a best fit to the points along the CME front is obtained, as in Figure 3.14.



on 24 January 2007. The white dots along the CME front indicate the multiscale edge detection to which the ellipse is fit. The black cross on the ellipse indicates the furthest point measured from Sun centre which provides the height-time Figure 3.14: A depiction of three frames sampled from the ellipse characterisation of a CME observed by LASCO/C2 profile for determining the kinematics. The opening angle from Sun centre measures the CME width.

Chapter 4

The Kinematics and Morphology of CMEs using Multiscale Methods

The diffuse morphology and transient nature of CMEs make them difficult to identify and track using traditional image processing techniques. We apply multiscale methods to enhance the visibility of the faint CME front. This enables an ellipse characterisation to objectively study the changing morphology and kinematics of a sample of events imaged by SOHO/LASCO and STEREO/SECCHI. The accuracy of these methods allows us to test the CMEs for non-constant acceleration and expansion. This chapter is founded on work published in Byrne et al. (2009).

4.1 Introduction

To date, most CME kinematics are derived from difference images; a technique based either on the subtraction of a single pre-event image (fixed-difference) or the subtraction of each image from the next in an event sequence (running-difference). These techniques are applied in order to highlight regions of changing intensity, increasing the relative brightness of the CME against the background coronal features. However, drawbacks do exist. Numerical differencing can enhance noise to a level comparable to the signal. The noise can be suppressed to a certain degree by using a standard box-car or median filter, but this will also smooth out CME features such as structure along the CME front and its environs. An additional issue resulting from differencing is the introduction of spatio-temporal cross-talk in difference frames. Since it is used to highlight non-stationary features in both space and time, then the differencing of subsequent images of a moving feature will show a signature at the position where the feature was initially observed and a signature at the position that the feature has moved to when next observed. Since the signature of motion in the difference images is heavily dependent on the time between frames and how many pixels the feature has moved, it may be considered to blend spatial and temporal information in a non-trivial manner: an effect referred to as spatio-temporal cross-talk. This can serve to blur out CME features and introduce ambiguity in estimating positions and times, critical to accurately deriving the kinematics of the event. Furthermore, user bias is introduced by the choice of intensity scaling and thresholding when determining the location of CME features by point-and-click methods or automated detection algorithms, as discussed in Section 3.1.

In this work we apply multiscale methods for analysing CMEs as described in Chapter 3, which has the benefit of working on independent images without any need for differencing. Once the edges of the CME front are resolved, an ellipse characterisation is applied to determine the CME kinematics (position, velocity, acceleration) and morphology (width, orientation) in coronagraph images. CME height measurements are taken as the height of the furthest point on the ellipse from Sun centre. The angular width is taken as the opening angle of the ellipse from Sun centre, and the tilt of the ellipse is given by the calculated angle γ . (Note that in cases where the code produces an extremely large and oblate ellipse with one apex approximating the CME front, the width and tilt information is deemed redundant. Hence the resulting analysis of some events can have less data points included in the width and tilt plots than in the height-time plots.) Following previous concerns on the errors in CME heights (e.g. Wen et al., 2007), multiscale methods allow us to determine the kinematics to a high degree of accuracy in order to improve confidence in their interpretation and comparison to theory. These methods also show potential for future automation. In this chapter a sample of CMEs observed by SOHO/LASCO and STEREO/SECCHI are studied, namely the gradual events of 2 January 2000, 18 April 2000, 23 April 2001, 1 April 2004, 8 October 2007 and 16 November 2007, and the impulsive events of 23 April 2000 and 21 April 2002. We compare our results with the catalogues of CDAW, CACTus and SEEDS (note ARTEMIS was not included due to difficulties interpreting the entries in its database that correspond to the chosen CMEs).

4.2 Error Analysis

The front of the CME is determined through the multiscale decomposition and consequent rendering of a gradient magnitude space. At scale 3 of the decomposition the smoothing filter is 2³ pixels wide, which we use as our error estimate in edge position. This error is input to the ellipse fitting algorithm for weighting the ellipse parameters, and a final error output is produced for each ellipse fit. In the case of a fading leading edge the reduced amount of points along the front will increase the error on our analysis. The final errors are displayed in the height-time plots of the CMEs, and are used in the velocity and acceleration calculations. The derivative is a 3-point Lagrangian interpolation, so there is an enhancement of error at the edges of the data sets as explained below.

4.2.1 3-Point Lagrangian Interpolation

3-point Lagrangian interpolation is used on the discrete set of given data points in order to determine the first and second derivatives corresponding to the velocity and acceleration of the CME height-time measurements in a more robust manner than simple forward- or reverse-derivatives. Considering three data points, $(x_0, y_0), (x_1, y_1), (x_2, y_2)$, the Langrangian interpolation polynomial is given by:

$$L(x) = \sum_{j=0}^{k} y_{j} l_{j}(x)$$

$$\dots l_{j}(x) = \prod_{i=0, i \neq j}^{k} \frac{x - x_{i}}{x_{j} - x_{i}}$$
(4.1)

$$\Rightarrow L(x) = y_0 l_0(x) + y_1 l_1(x) + y_2 l_2(x)$$

$$= y_0 \left(\frac{x - x_1}{x_0 - x_1} \frac{x - x_2}{x_0 - x_2} \right) + y_1 \left(\frac{x - x_0}{x_1 - x_0} \frac{x - x_2}{x_1 - x_2} \right) + y_2 \left(\frac{x - x_0}{x_2 - x_0} \frac{x - x_1}{x_2 - x_1} \right)$$

So the derivative is determined to be:

$$L' \equiv \frac{\partial L(x)}{\partial x}$$

$$= y_0 \frac{2x - x_1 - x_2}{(x_0 - x_1)(x_0 - x_2)} + y_1 \frac{2x - x_0 - x_2}{(x_1 - x_0)(x_1 - x_2)} + y_2 \frac{2x - x_0 - x_1}{(x_2 - x_0)(x_2 - x_1)}$$
(4.2)

And the edge point $x = x_0$ (and similarly for $x = x_n$) is weighted as follows:

$$d_0 = y_0 \frac{2x_0 - x_1 - x_2}{(x_0 - x_1)(x_0 - x_2)} + y_1 \frac{x_0 - x_2}{(x_1 - x_0)(x_1 - x_2)} + y_2 \frac{x_0 - x_1}{(x_0 - x_2)(x_1 - x_2)}$$
(4.3)

In the case where the points are equally spaced this is simply:

$$d_0 = \frac{1}{2} \left[-3y_0 + 4y_1 - y_2 \right] \tag{4.4}$$

The error propagation equation is used to determine the errors on the resulting derivative points in $L' \equiv f(L(x), x)$:

$$\sigma_{L'}^2 = \sigma_L^2 \left(\frac{\partial L'}{\partial L}\right)^2 + \sigma_x^2 \left(\frac{\partial L'}{\partial x}\right)^2 + \dots \tag{4.5}$$

$$= \frac{\sigma_L^2}{\partial x^2} + \frac{\sigma_x^2}{\partial x^2} \left(\frac{\partial L}{\partial x}\right)^2 \tag{4.6}$$

Or more appropriately written in this context as:

$$\sigma_d^2 = \frac{\sigma_{y_{n+1}}^2 + \sigma_{y_{n-1}}^2}{dx^2} + \frac{\sigma_{x_{n+1}}^2 + \sigma_{x_{n-1}}^2}{dx^2} \left(\frac{dy}{dx}\right)^2 \tag{4.7}$$

So the errors on the end points become:

$$\sigma_{d_0}^2 = \frac{9\sigma_{y_0}^2 + 16\sigma_{y_1}^2 + \sigma_{y_2}^2}{\left(x_2 - x_0\right)^2} + \frac{\sigma_{x_2}^2 + \sigma_{x_0}^2}{\left(x_2 - x_0\right)^2} \left(\frac{3y_0 - 4y_1 + y_2}{x_2 - x_0}\right)^2 \tag{4.8}$$

$$\sigma_{d_n}^2 = \frac{9\sigma_{y_n}^2 + 16\sigma_{y_{n-1}}^2 + \sigma_{y_{n-2}}^2}{\left(x_n - x_{n-2}\right)^2} + \frac{\sigma_{x_{n-2}}^2 + \sigma_{x_n}^2}{\left(x_{n-2} - x_n\right)^2} \left(\frac{3y_n - 4y_{n-1} + y_{n-2}}{x_{n-2} - x_n}\right)^2 \tag{4.9}$$

This effect is reflected in the larger errorbars on the end points of the derived kinematics of Section 4.3.

The errors on the heights are used to constrain the best fit to a constant acceleration model of the form:

$$h(t) = at^2 + v_0 t + h_0 (4.10)$$

where t is time and a, v_0 and h_0 are the acceleration, initial velocity and initial height respectively. This provides a linear fit to the derived velocity points and a constant fit to the acceleration. An important point to note is the small time error (taken to be the image exposure time of the coronagraph data) since the analysis is performed upon the observed data frames individually. Previous methods of temporal-differencing would increase this time error. With these more accurate measurements we are better able to determine the velocity and acceleration errors, leading to improved constraints upon the data and providing greater confidence in comparing to theoretical models.




Figure 4.1: A sample of ellipse fits to the multiscale edge detection of the events studied. For each event the upper and lower image show LASCO/C2 and C3, except for the 2007 events which show SECCHI/COR1 and COR2.

4.3 Results

This section outlines events which have been analysed using our multiscale methods. We use data from the LASCO/C2 and C3, and SECCHI/COR1 and COR2 instruments, and preprocess the images as discussed in Section 3.1. The ellipse fitting algorithm applied to each event gives consistent heights of the CME front measured from Sun centre to the maximum height on the ellipse, and these lead to velocity and acceleration profiles of our events. The ellipse fitting also provides the angular widths and orientations, as shown below. The velocity, acceleration and angular width results of each method are highlighted in Tables 4.1, 4.2 and 4.3. In each instance we include the values from CACTus, CDAW and SEEDS. Note that CACTus lists a median speed of the CME; CDAW provide the speed at the final height and from the velocity profile we infer the speed at the initial height; and the SEEDS detection applies only to the LASCO/C2 field-of-view but doesn't currently provide a velocity range or profile. Note also that the CMEs of 8 October 2007 and 16 November 2007 are analysed in SECCHI images by CACTus and our multiscale methods (marked by asterisks in the Tables), while CDAW and SEEDS currently only provide LASCO analysis. It is clear that many of the CACTus, CDAW and SEEDS results lie outside the results and error ranges of our analysis.

4.3.1 Arcade Eruption: 2 January 2000

This CME was first observed in the south-west at 06:06 UT on 2 January 2000 and appears to be a far-side event associated with an arcade eruption consisting of one or more bright loops.

Table 4.1: Summary of CME velocities as measured by CACTus, CDAW, SEEDS and multiscale methods. Asterisks indicate analysis of SECCHI data rather than LASCO.

Date	CACTus	CDAW	SEEDS	Multiscale
		${\rm km~s^{-1}}$		
02 Jan 2000	512	370 - 794	396	396 - 725
18 Apr 2000	463	410 - 923	339	324 - 1049
23 Apr 2000	1041	1490 - 898	595	1131 - 1083
23 Apr 2001	459	540 - 519	501	581 - 466
21 Apr 2002	1103	2400 - 2388	702	2195 - 2412
01 Apr 2004	487	300 - 613	319	415 - 570
08 Oct 2007	235*	85 - 331	103	71 - 330*
16 Nov 2007	337*	210 - 437	154	131 - 483*

Table 4.2: Summary of CME accelerations as measured by CACTus, CDAW, SEEDS and multiscale methods. Asterisks indicate analysis of SECCHI data rather than LASCO.

Date	CACTus	CDAW	SEEDS	Multiscale
		$\rm m\ s^{-2}$		
02 Jan 2000	0	21.3	-5.8	14.7 ± 3.6
18 Apr 2000	0	23.1	17.5	32.3 ± 3.5
23 Apr 2000	0	-48.5	-8.9	-4.8 ± 20.6
23 Apr 2001	0	-0.7	-1.4	-4.8 ± 4.1
21 Apr 2002	0	-1.4	33.5	32.5 ± 26.6
01 Apr 2004	0	7.1	12.9	4.4 ± 2.0
08 Oct 2007	0*	3.4	2.4	$5.7 \pm 0.9*$
16 Nov 2007	0*	4.9	11.0	$13.7 \pm 1.7^*$

The height-time plot has a trend not unlike that of CDAW (overplotted in top Figure 4.2 with a dashed line). However the offset of the CDAW heights - which puts them outside our error bounds - may be due to how the difference images are scaled for display. This is a problem multiscale methods avoid. From Figure 4.2, the velocity-fit was found to be increasing from 396 to 725 km s⁻¹, giving an acceleration of 14.7 ± 3.6 m s⁻². The ellipse fit spans approximately

Table 4.3: Summary of CME angular widths as measured by CACTus, CDAW, SEEDS and multiscale methods. Asterisks indicate analysis of SECCHI data rather than LASCO.

Date	CACTus	CDAW	SEEDS	Multiscale
		degrees		
02 Jan 2000	160	107	96	50 - 95
18 Apr 2000	106	105	108	68 - 110
23 Apr 2000	352	360	130	96 - 130
23 Apr 2001	124	91	74	55 - 60
21 Apr 2002	352	360	186	53 - 65
01 Apr 2004	66	79	58	44 - 38
$08 \ {\rm Oct} \ 2007$	52*	82	59	23 - 60*
16 Nov 2007	68*	78	54	40 - 55*

50–70° of the field-of-view in the inner portion of C2, and expands to over 95° in C3. This expansion may simply be attributed to the inclusion of one or more loops in the ellipse fit as the arcade traverses the LASCO/C2 and C3 fields-of-view. The orientation of the ellipse as a function of time is shown in bottom Figure 4.2. It can be seen that the orientation angle of the CME increases to approximately 100° before decreasing toward 60°.

The constant acceleration model is not a sufficient fit to the data in this event. The kinematics produced from the multiscale edge detection would be better fit with a non-linear velocity and a non-constant acceleration. This would show the CME to have a period of decreasing acceleration in the C2 field-of-view, leveling off to zero in C3 (if not decelerating further).

4.3.2 Gradual/Expanding CME: 18 April 2000

This CME was first observed off the south limb at 16:06 UT on 18 April 2000 and exhibits a flux rope type structure.




Figure 4.2: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 2 January 2000 CME observed by LASCO/C2 and C3. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The CDAW heights are over-plotted with a dashed line. The height and velocity fits are based upon the constant acceleration model.

The height-time plot for this event has a trend similar to that of CDAW (overplotted in top Figure 4.3 with a dashed line). The velocity-fit was found to be linearly increasing from 324 to over 1,000 km s⁻¹, giving an acceleration of 32.3 ± 3.5 m s⁻². The ellipse fit spans from 68° of the field-of-view in the inner portion of C2, to approximately 110° in C3. The orientation of the ellipse as a function of time is shown to increase from just above 0° to over 60° in Figure 4.3.

This CME is fitted well with the constant acceleration model but shows an increasing angular width implying expansion across the field-of-view.

4.3.3 Impulsive CME: 23 April 2000

This impulsive CME was first observed in the west at 12:54 UT on 23 April 2000 and exhibits strong streamer deflection.

The height-time plot derived using our methods has a trend which diverges from that of CDAW (overplotted in top Figure 4.4 with a dashed line). The velocity-fit was found to be linearly decreasing from 1,131 to 1,083 km s⁻¹, giving a constant deceleration of -4.8 ± 20.6 m s⁻². The CME is present for one frame in C2 with an ellipse fit spanning 96°, increasing to approximately $120-130^{\circ}$ in the C3 field-of-view, and the orientation of the ellipse as a function of time is shown to rise from 71° to 95° then fall to 64° (see bottom Figure 4.4).

This event is modeled satisfactorily with a constant deceleration. However, due to the impulsive nature of the CME there are only a few frames available for analysis, making it difficult to constrain the kinematics.




Figure 4.3: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 18 April 2000 CME observed by LASCO/C2 and C3. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The CDAW heights are over-plotted with a dashed line. The height and velocity fits are based upon the constant acceleration model.




Figure 4.4: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 23 April 2000 CME observed by LASCO/C2 and C3. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The CDAW heights are over-plotted with a dashed line. The height and velocity fits are based upon the constant acceleration model.

4.3.4 Faint CME: 23 April 2001

This CME was first observed in the south-west at 12:39 UT on 23 April 2001 and exhibits some degree of streamer deflection.

The height-time plot has a similar trend to CDAW (overplotted in top Figure 4.5 with a dashed line). The velocity-fit was found to be linearly decreasing from 581 to 466 km s⁻¹, giving a deceleration of -4.8 ± 4.1 m s⁻². The ellipse fit spans approximately $55-60^{\circ}$ of the field-of-view throughout the event, and the orientation of the ellipse as a function of time is shown to decrease from approximately 50° to almost 0° (see Figure 4.5).

This CME is fitted well with the constant acceleration model.

4.3.5 Fast CME: 21 April 2002

This CME was first observed in the west from 01:27 UT on 21 April 2002.

The height-time plot follows a similar trend to that of CDAW (overplotted in top Figure 4.6 with a dashed line). The velocity-fit was found to be linearly increasing from 2,195 to 2,412 km s⁻¹, giving a constant acceleration fit of 32.5 ± 26.6 m s⁻². The ellipse fit spans 53° in C2, and shows an increasing trend to 65° in C3. The orientation of the ellipse as a function of time is shown to scatter about 115° though it drops to approximately 81° in the final C3 image.

The kinematics of this event are not modeled satisfactorily by the constant acceleration model, since the fits do not lie within all error bars. The argument for a non-linear velocity profile, with a possible early decreasing acceleration, is justified for this event, although the instrument cadence limits the data set available for interpretation. The previous analysis of Gallagher *et al.* (2003)




Figure 4.5: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 23 April 2001 CME observed by LASCO/C2 and C3. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The CDAW heights are over-plotted with a dashed line. The height and velocity fits are based upon the constant acceleration model.

resulted in a velocity of $\sim 2{,}500~\rm km~s^{-1}$ past $\sim 3.4~R_{\odot}$ which is consistent with our results past $\sim 6~R_{\odot}$ in Figure 4.6.

4.3.6 Flux-Rope/Slow CME: 1 April 2004

This CME was first observed in the north-east from approximately 23:00 UT on 1 April 2004, is in the field-of-view for over 9 hours, and exhibits a bright loop front, cavity and twisted core.

The height-time plot follows a similar trend to that of CDAW (overplotted in top Figure 4.7 with a dashed line). The velocity-fit was found to be linearly increasing from 415 to 570 km s⁻¹, giving an acceleration of 4.4 ± 2.0 m s⁻². Note also that the kinematics of this event exhibit non-linear structure clearly seen in the velocity and acceleration profiles. The ellipse fit spans approximately 44° in C2, stepping down to approximately 38° in C3. The orientation of the ellipse as a function of time is shown to jump down from approximately 130° in C2 to approximately $70-80^{\circ}$ in C3.

This event shows unexpected structure in the velocity and acceleration profiles which indicates a complex eruption not satisfactorily modeled with constant acceleration.

4.3.7 STEREO-B Event: 8 October 2007

This CME was first observed in the west from approximately 12:00 UT on 8 October 2007, and is best viewed from the STEREO-B spacecraft. It is noted that the kinematics as measured by SOHO and STEREO will be different due to projection effects (Howard *et al.*, 2008b; Vršnak *et al.*, 2007). On this date




Figure 4.6: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 21 April 2002 CME observed by LASCO/C2 and C3. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The CDAW heights are over-plotted with a dashed line. The height and velocity fits are based upon the constant acceleration model.




Figure 4.7: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 1 April 2004 CME observed by LASCO/C2 and C3. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The CDAW heights are over-plotted with a dashed line. The height and velocity fits are based upon the constant acceleration model.

STEREO-B was at an angular separation of 16.5° from Earth.

The height-time plot for this event is shown in Figure 4.8. The velocity-fit was found to be linearly increasing from 71 to 330 km s⁻¹, giving an acceleration of 5.7 ± 0.9 m s⁻². The ellipse fit in COR1 spans approximately 23° stepping up to a scatter about $40-50^{\circ}$ which rises slightly to $50-60^{\circ}$ in COR2. The orientation of the ellipse as a function of time is shown to increase from $55-110^{\circ}$ then jumps to an approximately steady scatter about $180-190^{\circ}$.

This CME is fitted well with the constant acceleration model.

4.3.8 STEREO-A Event: 16 November 2007

This CME was first observed in the west from approximately 08:26 UT on 16 November 2007, and is best viewed from the STEREO-A spacecraft. On this date STEREO-A was at an angular separation of 20.3° from Earth.

The height-time plot for this event is shown in Figure 4.9. The velocity-fit was found to be linearly increasing from 131 to 483 km s⁻¹, giving an acceleration of 13.7 ± 1.7 m s⁻². The ellipse fit in COR1 spans approximately $40-50^{\circ}$ stepping up slightly to a scatter about $45-55^{\circ}$ in COR2. The orientation of the ellipse as a function of time is shown to start at 153° and end at 120° with the mid points scattered about 170° .

This CME is fitted well with the constant acceleration model.

4.4 Discussion & Conclusions

A variety of theoretical models have been proposed to describe CMEs, especially their early propagation phase. Observational studies, such as those outlined




Figure 4.8: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 8 October 2007 CME observed by SECCHI/COR1 and COR2. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The height and velocity fits are based upon the constant acceleration model.




Figure 4.9: Kinematic and morphological profiles for the ellipse fit to the multiscale edge detection of the 16 November 2007 CME observed by SECCHI/COR1 and COR2. The plots from top to bottom are height, velocity, angular width, and ellipse tilt. The height and velocity fits are based upon the constant acceleration model.

above, are necessary to determine CME characteristics. We argue that the results of previous methods are limited in this regard due mainly to large kinematic errors which fail to constrain a model, an artefact of CME detection based upon either running- (or fixed-) difference techniques or other operations. Current methods fit either a linear model to the height-time curve, implying constant velocity and zero acceleration (e.g. CACTus) or a second order polynomial, producing a linear velocity and constant acceleration (e.g. CDAW, SEEDS). The implementation of a multiscale decomposition provides a time error on the scale of seconds (the exposure time of the instrument) and a resulting height error on the order of a few pixels. The height-time error is used to determine the errors of the velocity and acceleration profiles of the CMEs. It was shown that for certain events the results of CACTus, CDAW and SEEDS can differ significantly from our methods, as illustrated in the Tables of Section 4.3.

Our results clearly confirm that the constant acceleration model may not always be appropriate. The 2 January 2000 and 21 April 2002 CMEs are good examples of the possible non-linear velocity profile and consequent non-constant acceleration profile (see Figure 4.2 and Figure 4.6). Indeed these events are shown to have a decreasing acceleration, possibly to zero or below, as the CMEs traverse the field-of-view. Simulations of the breakout model outlined in Lynch et al. (2004) resulted in constant acceleration fits which do not agree with these observations. It may be further noted that the events of 23 April 2001 and 1 April 2004 show a possible decreasing acceleration phase early on, though within errors this cannot be certain (see Figure 4.5 and Figure 4.7). Furthermore, the structure seen in some events would indicate that the CME does not progress smoothly. The velocities of the 1 April 2004 CME in Figure 4.7 and the 16 November 2007

CME in Figure 4.9 show non-smooth profiles and may imply a form of bursty reconnection or other staggered energy release driving the CME. Other profiles such as Figure 4.2 and to a lesser extent Figures 4.5 and 4.6 may show a stepwise pattern, indicative of separate regimes of CME progression. None of the current CME models indicate a form of non-smooth progression, although the flux-rope model does describe an early acceleration regime giving a non-linear velocity to the eruption (see Figure 11.5 in Priest & Forbes, 2000).

It may be concluded that the angular widths of the events are indicative of whether the CME expands radially or otherwise in the plane-of-sky. For the CMEs studied above, the observations of 18 April 2000, 23 April 2000, and 21 April 2002 show an angular width expansion (see Figure 4.3, Figure 4.4, and Figure 4.6). These events also show high velocities, obtaining top speeds of up to 1,000 km s⁻¹, over 1,100 km s⁻¹ and 2,500 km s⁻¹ respectively, and may therefore indicate a link between the CME expansion and speed. Furthermore, it is suggested by Krall & St. Cyr (2006) that the flux-rope model can account for different observed expansion rates due to the axial versus broadside view of the erupting flux system.

The observed morphology of the ellipse fits may be further interpreted through the tilt angles plotted in Section 4.3. In knowing the ellipse tilt and the direction of propagation of the CME it is possible to describe the curvature of the front. For the events above, the changing tilt and hence curvature is possibly significant for the 18 April 2000, 1 April 2004, and 8 October 2007 events (see bottom Figure 4.3, Figure 4.7, and Figure 4.8). The elliptical flux rope model of Krall *et al.* (2006) was shown to have a changing orientation of the magnetic axis which results in a dynamic radius of curvature of the CME, possibly accounting for these observed

ellipse tilts.

The work outlined here is an initial indication that the zero and constant acceleration models in CME analysis are not an accurate representation of all events, and the over-estimated angular widths are not indicative of the true CME expansion. The ellipse characterisation has provided additional information on the system through its changing width and orientation. This work will be further explored and developed with STEREO data whereby the combined view-points can give additional kinematic constraints and lead to a correction for projection effects through 3D reconstructions (discussed in Chapter 5).

Chapter 5

Propagation of an Earth-Directed CME in Three-Dimensions

Although CMEs are long known to be significant drivers of adverse space weather at Earth, the physics governing their propagation is not fully understood, due predominantly to the plane-of-sky projection effects of low-cadence imagers with restricted fields-of-view. The launch of the STEREO mission in 2006 has provided new insight into their 3D propagation in the heliosphere, although the mechanisms governing their evolution remain unclear due to difficulties in reconstructing their true 3D structure. Here we use a new elliptical tie-pointing technique to reconstruct a full CME front in 3D, enabling us to quantify an early acceleration profile, deflected motion along the ecliptic, increasing angular width and 'pancaking' of the CME front as it propagates from $2-46~{\rm R}_{\odot}~(\sim 0.2~{\rm AU})$. Beyond $7~{\rm R}_{\odot}$, we show that its motion is determined by aerodynamic drag in the solar wind and, using our reconstruction as input for a 3D MHD simulation, we determine an accurate arrival time at the L1 point near Earth. This chapter is

founded on research published in Byrne et al. (2010).

5.1 Introduction

It is predominantly believed that magnetic reconnection is responsible for the destabilisation of magnetic flux-ropes on the Sun, which then erupt through the corona into the solar wind to form CMEs (Moore & Sterling, 2006). There is much debate as to the specific processes which trigger the eruption of CMEs, and different models exist to explain these (Antiochos et al., 1999; Chen, 1996; Forbes & Priest, 1995; Kliem & Török, 2006; van der Holst et al., 2007). In the low solar atmosphere, it is postulated that high latitude CMEs undergo deflection since they are often observed at different position angles with respect to their associated source region locations (Xie et al., 2009). It has been suggested that field lines from polar coronal holes may guide high-latitude CMEs towards the equator (Kilpua et al., 2009), or that the initial magnetic polarity of a flux-rope relative to the background magnetic field influences its trajectory (Chané et al., 2005; Filippov et al., 2001). During this early phase, CMEs are observed to expand outwards from their launch site, though plane-of-sky measurements of their increasing sizes and angular widths are ambiguous in this regard (Gopalswamy et al., 2009a). This expansion has been modelled as a pressure gradient between the flux-rope and the background solar wind (Odstrčil & Pizzo, 1999; Riley & Crooker, 2004). At larger distances in their propagation, CMEs are predicted to interact with the solar wind and the interplanetary magnetic field. Studies that compare in-situ CME velocity measurements with initial eruption speeds through the corona show that slow CMEs must be accelerated toward the speed of the solar wind, and fast CMEs decelerated (González-Esparza et al., 2003; Maloney et al., 2009). It has been suggested that this is due to the effects of drag acting on the CME in the solar wind (Cargill, 2004; Tappin, 2006). However, the quantification of drag, along with that of both CME expansion and non-radial motion, is currently lacking, due primarily to the limits of observations from single fixed viewpoints with restricted fields-of-view.

Efforts to reconcile 2D plane-of-sky images with the true 3D morphology of CMEs have been underway since they were first observed in the 1970s. The inherent difficulties in this are predominantly due to the single, fixed-position imagers with restricted fields-of-view, as well as the difficulties in observing the optically thin coronal plasma of these dynamic events. Before the launch of STEREO, there was limited ability to infer the 3D CME morphology from the available observations such as SOHO/LASCO. Coronagraphs mainly measure the Thomson scattered light of the free electrons in the coronal plasma, providing white-light images of CMEs against the plane-of-sky that are not trivial to deconvolve, and the projected 2D nature of these images introduces uncertainties in kinematical and morphological analyses (Vršnak et al., 2007). Some efforts were based upon a pre-assumed geometry of the CME, such as the cylindrical model (Cremades & Bothmer, 2004) or the cone model (Xue et al., 2005; Zhao et al., 2002), whose shapes were simply oriented to best match the 2D observations. Others used either a comparison of multiple events to infer a statistical relationship between plane-of-sky measurements and true CME motion (Howard & Tappin, 2005; Schwenn et al., 2005), or a comparison of observations with in-situ data and/or signatures on-disk (Démoulin et al., 2008; Howard et al., 2008b). One prominent method was the use of 3D polarisation analysis of LASCO images (Moran & Davila, 2004), whereby the line-of-sight averaged distance from the plane-of-sky is determined from the brightness ratio of polarised to unpolarised electron scattered emissivity (K-corona). However, this lacks in details such as whether the feature is truly unique along the line-of-sight, and if so is it towards or away from the observer with respect to the plane-of-sky. Polarisation analysis itself is only acceptable up to heights of $\sim 5~\rm R_{\odot}$, since beyond this distance the dust-scattered F-corona may no longer be considered unpolarised (Billings, 1966). These issues motivated the launch of the STEREO mission to further our understanding of CMEs.

5.2 The STEREO Era

The two near-identical spacecraft of the STEREO mission provide simultaneous observations of CMEs from independent viewpoints to better observe their true morphology. Unfortunately there are limitations on how much 3D information can be extracted from the combined plane-of-sky observations, especially when the object is optically thin and its boundaries ill-defined. In order to determine the morphology of an object in 3D from only two viewpoints, techniques must be applied within the context of an epipolar geometry, as illustrated in Figure 5.1 (Inhester, 2006). This epipolar coordinate system for considering the 3D space observed from two independent viewpoints is built up as follows. A line is drawn to connect the two observers, called the stereo base line. The two observer locations and any third object point or location in the observing space then define a plane. Numerous object points will define numerous planes who share an intersection with the stereo base line. These are the epipolar planes of Figure 5.1.




Figure 5.1: Schematic of the epipolar geometry used to relate the observations from the two STEREO spacecraft, reproduced from Inhester (2006). This geometry enables us to localise features in 3D space by the triangulating sight-lines across epipolar planes.

The plane-of-sky from each observer then intersects the epipolar planes such that they appear as epipolar lines across the image, and will converge on a point along the stereo base line referred to as the epipole of that image. So if a line-of-sight from observer 1 is drawn across an epipolar plane, it will appear as a single point on image 1, but as a complete line across the corresponding epipolar plane in image 2 as seen by observer 2, who is then able to triangulate upon an object in 3D space by the intersecting lines-of-sight.

The technique of tie-pointing lines-of-sight across epipolar planes is best for




Figure 5.2: Schematic of how tie-pointing a curved surface within an epipolar geometry is limited in its ability to resolve the true feature, since lines-of-sight will be tangent to different edges of the surface and not necessarily intersect upon it. Reproduced from Inhester (2006).

Under the assumption that the same feature may be tracked in coronagraph images many CME studies have also employed tie-pointing techniques (Liewer et al., 2009; Mierla et al., 2008; Srivastava et al., 2009; Temmer et al., 2009; Wood et al., 2009). However, when measuring the kinematics of the CME front this technique alone doesn't hold true, since it is inevitable that the same part of the curved front cannot be confidently resolved from both viewpoints once the CME has traversed a certain distance in space, nor similarly once the spacecraft have moved beyond a certain angular separation during the mission (Figure 5.2). Furthermore, triangulating CME observations using only the COR images confines the kinematic and morpohological analyses to within the 20 R_{\odot} field-of-view. The additional use of the heliospheric imagers allows a study of CMEs out to distances of ~ 1 AU, however (instrumental effects aside) a 3D analysis can only be carried out if the




Figure 5.3: Schematic of the Graduated Cylindrical Shell (GCS) model of a flux rope CME, reproduced from Thernisien *et al.* (2009). Indicated are the model parameters of front height \mathbf{h}_{front} , leg height \mathbf{h} , angle between the legs 2α , cross-sectional radius \mathbf{a} , and distance from Sun centre \mathbf{O} to a point on the edge of the shell \mathbf{r} . Two views of the GCS model are shown; (\mathbf{a}) 'face-on', and (\mathbf{b}) 'end-on'. The positional parameters of longitude ϕ and latitude θ are illustrated in (\mathbf{c}).

CME propagates along a trajectory between the two spacecraft so that it is observed by both HI instruments. Otherwise, assumptions of its trajectory have to be inferred from either its association with a source region on-disk (Howard & Tappin, 2008) or its trajectory through the COR data (Maloney et al., 2009), or derived by assuming a constant velocity through the HI fields-of-view (Davis et al., 2009). Triangulation of CME features using time-stacked intensity slices at fixed latitude, named 'J-maps' due to the characteristic propagation signature of a CME, has also been developed (Davis et al., 2010; Liu et al., 2010). This technique is hindered by the same limitation of standard tie-pointing techniques; namely that the curvature of the feature is not considered, and the intersection of sight-lines may not occur upon the surface of the observed feature.

An alternative to tie-pointing is a method called forward modeling which

presumes a given shape of the CME and seeks to match it with observational data. Thernisien et al. (2006) employ a graduated cylindrical shell which is warped to form a flux rope model overlaid on CME images (Figure 5.3). The parameters governing the model's shape and orientation may be changed by the user to fit the model to STEREO-Ahead and Behind data simultaneously and obtain a 3D flux rope characterisation of the CME as it propagates, though this may not always be appropriate (Jacobs et al., 2009). Boursier et al. (2009a) outline a similar forward model which assumes one of three pre-assigned shapes: a hemispherical cap, a flux rope, or a cloud-like model. However, in each of these methods the predetermined shape of the CME model has a spherical cross-section and must adhere to some quasi-similarity (self-invariance) over the sequence of images. So while forward modelling better accounts for the curved nature of the CME being observed, the inherent restrictions of the imposed model still limits the analysis of the true 3D structure and dynamics of the CME as it propagates.

5.3 Elliptical Tie-Pointing

In the epipolar geometry outlined above, 3D information may be gleaned from two independent viewpoints of a feature using tie-pointing techniques to triangulate lines-of-sight in space. However, when the object is known to be a curved surface, sightlines will be tangent to it and not necessarily intersect upon it (Figure 5.2). Consequently CMEs cannot be reconstructed by tie-pointing alone, but rather their localisation may be constrained by intersecting sightlines tangent to the leading edges of a CME (de Koning et al., 2009; Pizzo & Biesecker, 2004). Following the multiscale edge detection and ellipse characterisation outlined in

Chapter 3, it is possible to extract the intersection of a given epipolar plane through the ellipse fits of both the STEREO-Ahead and Behind images. This defines a quadrilateral in 3D space which localises the ellipse characterisation of the CME front in that plane.

Inscribing an ellipse within the quadrilateral such that it is tangent to all four sides (detailed below) provides a slice through the CME that matches the observations from each spacecraft (Figure 5.4a). A full reconstruction is achieved by stacking ellipses from numerous epipolar slices (Figure 5.4b). Since the positions and curvatures of these inscribed ellipses are constrained by the characterised curvature of the CME front in the stereoscopic image pair, the modeled CME front is considered an optimum reconstruction of the true CME front. This is repeated for every frame of the eruption to build the reconstruction as a function of time and view the changes to the CME front as it propagates in 3D (Figure 5.4c).

Following Horwitz (2002, 2005), we inscribe an ellipse within a quadrilateral using the following steps (see Figure 5.5):

- 1. Apply an isometry to the plane such that the quadrilateral has vertices (0,0), (A,B), (0,C), (s,t), where in the case of an affine transformation we set A=1, B=0 and C=1, with s and t variable.
- 2. Set the ellipse centre point (h, k) by fixing h somewhere along the open line segment connecting the midpoints of the diagonals of the quadrilateral and hence determine k from the equation of a line, for example:

$$h = \frac{1}{2} \left(\frac{s}{2} + \frac{A}{2} \right), \quad k = \left(h - \frac{s}{2} \right) \left(\frac{t - B - C}{s - A} \right) + \frac{t}{2}$$
 (5.1)




Figure 5.4: The elliptical tie-pointing technique developed to reconstruct the 3D CME front, shown here for the 12 December 2008 event. One of any number of epipolar planes will intersect the ellipse characterisation of the CME at two points in each image from STEREO-A and B. (a) illustrates how the resulting four sight-lines intersect in 3D space to define a quadrilateral that constrains the CME front in that plane. Inscribing an ellipse within the quadrilateral such that it is tangent to each sight-line provides a slice through the CME that matches the observations from each spacecraft. (b) illustrates how a full reconstruction is achieved by stacking multiple ellipses from the epipolar slices to create a model CME front that is an optimum reconstruction of the true CME front. (c) illustrates how this is repeated for every frame of the eruption to build the reconstruction as a function of time and view the changes to the CME front as it propagates in 3D. While the ellipse characterisation applies to both the leading edges and, when observable, the flanks of the CME, only the outermost part of the reconstructed front is shown here for clarity.




Figure 5.5: An ellipse inscribed within a convex quadrilateral. An isometry of the plane is applied such that the quadrilateral has vertices (0,0), (A,B), (0,C), (s,t). The ellipse has center (h,k), semimajor axis a, semiminor axis b, tilt angle δ , and is tangent to each side of the quadrilateral.

3. To solve for the ellipse tangent to the four sides of the quadrilateral, we can solve for the ellipse tangent to the three sides of a triangle whose vertices are the complex points

$$z_1 = 0, \quad z_2 = A + Bi, \quad z_3 = -\frac{At - Bs}{s - A}i$$
 (5.2)

and the two ellipse foci are then the zeroes of the equation

$$p_h(z) = (s-A)z^2 - 2(s-A)(h-ik)z - (B-iA)(s-2h)C$$
 (5.3)

whose discriminant can be denoted by $r(h) = r_1(h) + ir_2(h)$ where

$$r_{1} = 4 \left((s - A)^{2} - (t - B - C)^{2} \right) \left(\frac{h - A}{2} \right)^{2}$$

$$+ 4 (s - A) (A (s - A) + B (B - t) + C (C - t)) \left(\frac{h - A}{2} \right)$$

$$+ (s - A)^{2} (A^{2} - (C - B)^{2})$$

$$+ 2 \left((s - A) (At + Cs + Bs - 2AB) \left(\frac{h - A}{2} \right) \right)$$

$$+ 2 A (s - A)^{2} (B - C)$$
(5.5)

Thus we need to determine the quartic polynomial $u(h) = |r(h)|^2 = r_1(h)^2 + r_2(h)^2$ and we can then solve for the ellipse semimajor axis, a, and semiminor axis, b, from the equations

$$a^{2} - b^{2} = \sqrt{\frac{1}{\left(16\left(s - A\right)^{4}\right)}u(h)}$$
 (5.6)

$$a^{2}b^{2} = \frac{1}{4} \left(\frac{C}{(s-A)^{2}} \right) \left(2 \left(Bs - A \left(t - C \right) \right) h - ACs \right) \left(2h - A \right) \left(2h - s \right) (5.7)$$

by parameterising $R = a^2 - b^2$ and $W = a^2b^2$ to obtain

$$a = \sqrt{\frac{1}{2} \left(\sqrt{R^2 + 4W} + R \right)}, \quad b = \sqrt{\frac{1}{2} \left(\sqrt{R^2 + 4W} - R \right)}$$
 (5.8)

4. Knowing the axes we can generate the ellipse and float its tilt angle δ until it sits tangent to each side of the quadrilateral, using the inclined ellipse

equation (3.15) introduced in Section 3.4.

5.3.1 SOHO as a Third Perspective

The elliptical tie-pointing technique was used to reconstruct the front of a CME observed by STEREO on 26 April 2008 in order to test its efficacy by comparing it with observations from SOHO - a third perspective on the event. The CME appears as a halo from STEREO-B, so a running-difference technique is used to highlight the faint CME front in the images. The front is defined in the images by a point-and-click methodology and characterised with an ellipse fit (outlined in Section 3.4). From STEREO-A the event appears off the east limb and shows a strong streamer deflection to the south-east (in fact the CME would probably be considered only as the northern portion of the erupting material in STEREO-A if it were not shown by STEREO-B to expand further south).

With the ellipse characterisations determined for the CME front in the simultaneous images from COR1 and COR2 onboard STEREO-A and B, the elliptical tie-pointing technique is performed and the CME front reconstructed in 3D. This reconstruction is then back-projected onto the LASCO plane-of-sky in order to compare it with observations of the CME from SOHO's vantage point at L1 (Figure 5.6). This back-projection is performed by standard geometry of lines-of-sight from the observer position $O(x_0, y_0, z_0)$ through the 3D reconstruction points $P(x_i, y_i, z_i)$ and determining where they intersect the plane-of-sky




Figure 5.6: The back-projection of a STEREO 3D CME front reconstruction onto the SOHO/LASCO plane-of-sky, from the observations by STEREO-A (red) and STEREO-B (green) at 16:22 UT on 26 April 2008.

 $Q(x_j = 0, y_j, z_j)$ as follows:

$$\tan \alpha = \frac{y_i - y_0}{x_i - x_0} = \frac{y_j - y_0}{x_j - x_0} = \frac{y_j - y_i}{x_j - x_i} \quad \Rightarrow \quad y_j = x_i \left(\frac{y_i - y_0}{x_i - x_0}\right) + y_i \quad (5.9)$$

$$\tan \beta = \frac{z_i - z_0}{x_i - x_0} = \frac{z_j - z_0}{x_j - x_0} = \frac{z_j - z_i}{x_j - x_i} \quad \Rightarrow \quad z_j = x_i \left(\frac{z_i - z_0}{x_i - x_0}\right) + z_i \quad (5.10)$$

$$\tan \beta = \frac{z_i - z_0}{x_i - x_0} = \frac{z_j - z_0}{x_j - x_0} = \frac{z_j - z_i}{x_j - x_i} \Rightarrow z_j = x_i \left(\frac{z_i - z_0}{x_i - x_0}\right) + z_i \quad (5.10)$$

Due to the different instrument cadences of the SECCHI and LASCO coronagraphs, frames which lie closest in time were chosen for comparison. Figure 5.7 shows the COR2 frames from STEREO-A and B at 16:22 UT on 26 April 2008 with the ellipse characterisations of the CME front (left and right panels), and the back-projected 3D front reconstruction as compared with the LASCO/C2 frame from SOHO at 16:30 UT (middle panel). The reconstruction from the STEREO observations adequately fits with the SOHO observations given the time offset, and so gives credence to the elliptical tie-pointing technique.

5.4 Earth-Directed CME

On 12 December 2008 an erupting prominence was observed by STEREO while the spacecraft were in near quadrature at 86.7° separation (Figure 5.8a). The eruption is visible at 50-55° north from 03:00 UT in SECCHI/EUVI images, obtained in the 304 Å passband, in the northeast from the perspective of STEREO-A and off the northwest limb from STEREO-B. The prominence is considered to be the inner material of the CME which was first observed in COR1-B at 05:35 UT (Figure 5.8b). For our analysis, we use the two coronagraphs (COR1/2) and the inner Heliospheric Imagers (HI1) (Figure 5.8c). In each image the front of the CME is fitted with an ellipse that characterises its propagation across the planeof-sky (Byrne et al., 2009). This ellipse fitting is sensitive predominantly to the leading edges of the CME but equal weight is given to the CME flank edges as they enter the field-of-view of each instrument. The 3D reconstruction is then performed using a method of curvature constrained tie-pointing within epipolar planes containing the two STEREO spacecraft (detailed in Section 5.3). An example of the ellipse characterisation to the CME front and the corresponding back-projected 3D reconstruction is shown in Figure 5.9 for COR1, Figure 5.10 for COR2, and Figure 5.11 for HI1. Corresponding frames from a 3D visuali-




Figure 5.7: The back-projection of a STEREO 3D CME front reconstruction onto the SOHO/LASCO plane-of-sky, at 16:22 UT on 26 April 2008. The left panel shows the multiscale edge detection and ellipse characterisation of the CME front observed off the east limb from STEREO-A. The right panel shows the running-difference point-and-click ellipse characterisation of the CME front observed as a halo event from STEREO-B. . The middle panel shows the LASCO/C2 observation of the CME at 16:30 UT (8 minutes later than the STEREO observations) with the back-projected 3D CME front reconstruction overplotted for comparison.




Figure 5.8: Composite of STEREO-Ahead and Behind images from EUVI, COR1, COR2, and HI1 (Byrne *et al.*, 2010). (a) indicates the STEREO spacecraft locations, separated by an angle of 86.7° at the time of the event. (b) shows the prominence eruption observed in EUVI-B off the NW limb from approximately 03:00 UT which is considered to be the inner material of the CME. The multiscale edge detection and corresponding ellipse characterisation are overplotted in COR1. (c) shows that the CME is Earth-directed, being observed off the east limb in STEREO-A and the west limb in STEREO-B.




Figure 5.9: STEREO-Ahead and Behind COR1 images at 07:35 UT on the 12 December 2008. Overplotted are: the multiscale edge detections of the CME front (red); the ellipse characterisations (blue); and the resulting 3D reconstructions back-projected onto the plane-of-sky (white).




Figure 5.10: STEREO-Ahead and Behind COR2 images at 14:52 UT on the 12 December 2008. Overplotted are: the multiscale edge detections of the CME front (red); the ellipse characterisations (blue); and the resulting 3D reconstructions back-projected onto the plane-of-sky (white).




Figure 5.11: STEREO-Ahead and Behind HI1 images at 01:29 UT on the 13 December 2008. Overplotted are: the running difference edge detections of the CME front (red); the ellipse characterisations (blue); and the resulting 3D reconstructions back-projected onto the plane-of-sky (white).




Figure 5.12: The 3D CME front reconstruction as it propagates along the Sun-Earth line into interplanetary space. This particular frame of the reconstruction is from the observations of COR1 at 07:35 UT on 12 December 2008.




Figure 5.13: The 3D CME front reconstruction as it propagates along the Sun-Earth line into interplanetary space. This particular frame of the reconstruction is from the observations of COR2 at 14:52 UT on 12 December 2008.




Figure 5.14: The 3D CME front reconstruction as it propagates along the Sun-Earth line into interplanetary space. This particular frame of the reconstruction is from the observations of HI1 at 01:29 UT on 13 December 2008.

sation of the event are shown in Figure 5.12 for COR1, Figure 5.13 for COR2, and Figure 5.14 for HI1, showing the relative locations of the Sun, Earth and STEREO spacecrafts in the inner heliosphere.

5.5 Results

The resulting kinematics and morphology of the CME are measured along an angular span through the reconstructed CME front in the out-of-ecliptic plane along the Sun-Earth line (Figure 5.15). These were taken by first closing tangents to the CME front ('Northern/Southern Flanks'), and then measuring the height along an angle midway between these ('Midpoint of Front'), and then similarly along the two angles midway between the midpoint and the flanks ('Midtop/Midbottom of Front'). Although these measurements are fixed along the Sun-Earth line, investigating how the CME height profile would change if taken along a trajectory slightly off the Sun-Earth line shows no significant deviation within the associated errors and thus has negligible effect on the kinematics. The same is true if the overall maximum height (of varying location) on each individual CME front is instead taken and the kinematics reanalysed.

5.5.1 3D Error Propagation

When considering the errors that propagate from the 2D plane-of-sky of each image onto the 3D quadrilateral localising the CME, we may assume that the lines-of-sight within the error range are essentially parallel. This means the error interval on the coordinate being tie-pointed in 3D is given by a trapezoid surrounding the intersection of the lines-of-sight, illustrated in Figure 5.16. This is




Figure 5.15: The 3D CME front reconstruction from the COR2 Ahead and Behind frames at 14:52 UT on 12 December 2008. The lines drawn from Sun-centre indicate the 'Midpoint of Front' (solid blue), the 'Northern/Southern Flanks' (solid red/brown), and the 'Midtop/Midbottom of Front' at angles in between (dashed red/brown). By taking these measurements across all frames we may determine the kinematics and morphology of the CME as plotted in Figures 5.17 and 5.18.




Figure 5.16: The error trapezoid on the tie-pointing of two lines-of-sight in 3D space, reproduced from Inhester (2006). The error w in localising a point on each plane-of-sky results in a trapezoid with a diagonal measuring $w/\sin(\alpha/2)$ as shown.

done for each corner of the quadrilateral within a given epipolar plane. For a spacecraft separation of angle α and errorbar of magnitude w on the 2D image we can define the error trapezoid as having diagonals of length $w/\cos(\alpha/2)$ and $w/\sin(\alpha/2)$.

So in the case of COR1/2, the optimum filter size in the multiscale decomposition was 2^3 pixels wide, giving an error of ± 8 pixels, so w = 16. Over the course of the 12 December 2008 CME the average STEREO spacecraft separation was 86.75° , so we calculate the error trapezoid as having diagonals of size:

$$\left[\frac{w}{\cos\left(\frac{\alpha}{2}\right)}, \frac{w}{\sin\left(\frac{\alpha}{2}\right)}\right] = [11.0, 11.6] \tag{5.11}$$

This provides a 3σ height error of 11.6 pixels, so the corresponding 1σ height error is given by 68(11.6)/99.7 = 7.9 pixels. The time error for the multiscale edge detections is given by the exposure time of the individual frames: 1.69984 seconds for COR1 and 2.00090 seconds for COR2.

In the case of HI1 a 1σ plane-of-sky error of 3 pixels was determined (Maloney et al., 2009), so w = 6 and the error trapezoid is deduced to be [4.1, 4.4]. So the height error for HI1 is taken as 4.4 pixels and the time error is given by the thirty summed 60 second images to result in 1800 seconds (Eyles et al., 2009).

These errors are transformed first into arcseconds by multiplying by the plate scale of the instruments (7.5043001 arcsec/pixel for COR1, 14.7 arcsec/pixel for COR2, and 71.927554 arcsec/pixel for HI1), and then into metres, knowing the respective size of the Sun in arcseconds on the plane-of-sky observed by each instrument (given 1 $R_{\odot} = 6.95508 \times 10^8$ m). The resultant errors are then propagated with the 3-point Lagrangian interpolation (detailed in Section 4.2.1) from the height-time curves into the velocity and acceleration profiles of Figure 5.18. Due to the potentially large deviation of the end points from the general trend in 3-point Lagrangian interpolation, the endpoints of the EUVI prominence data, the COR1/2 coronagraph data, and the HI data are each removed as outliers from the velocity and acceleration plots.

5.5.2 Prominence & CME Acceleration

In determining the CME kinematics for the 'Midpoint of Front' we find a steep increase in the velocity corresponding to an early impulsive acceleration phase of 94 ± 58 m s⁻² (bottom of Figure 5.17). The prominence rises with a velocity




Figure 5.17: The kinematics of the prominence and 3D reconstructed CME front of 12 December 2008. The prominence is observed as the inner material of the CME, with both undergoing acceleration from $\sim 06:00-07:00$ UT, peaking at ~ 40 m s⁻² and ~ 94 m s⁻² respectively, before reducing to scatter about zero. Measurement uncertainties are indicated by one standard deviation error bars.

of $\sim 50~\rm km~s^{-1}$ before the system fully erupts and the prominence undergoes an acceleration of $40\pm 5~\rm m~s^{-2}$ behind the CME front. This is indicative of the onset of explosive reconnection or other loss-of-equilibrium in the system whereby the internal magnetic pressure increases sufficiently and/or the external magnetic pressure decreases sufficiently to allow the eruption to proceed in the height range $\sim 1.5-3~\rm R_{\odot}$ (top panel of Figure 5.18). The velocity profile is synonymous with those produced by the 2D flux-rope model as in Figure 11.5 of Priest & Forbes (2000). The acceleration then reduces to scatter about zero as the explosive nature of the eruption due to the Lorentz force diminishes and the drag force due to the ambient solar wind pressure begins to dominate (see equation 1.26).

5.5.3 Non-radial Prominence & CME Motion.

It is immediately evident from the reconstruction (illustrated in Figure 5.4c) that the CME propagates non-radially away from the Sun. The CME flanks change from an initial latitude span of $16-46^{\circ}$ to finally span approximately $\pm 30^{\circ}$ of the ecliptic (middle panel of Figure 5.18). The mean declination, θ , of the CME is well fitted by a power-law of the form $\theta(r) = \theta_0 r^{-0.92}$ (2 $R_{\odot} < r < 46 R_{\odot}$) as a result of this non-radial propagation. Tie-pointing the prominence apex and fitting a power-law to its declination angle results in $\theta^{prom}(r) = \theta_0^{prom} r^{-0.82}$ (1 $R_{\odot} < r < 3 R_{\odot}$), implying a source latitude of $\theta_0^{prom}(1 R_{\odot}) \approx 54^{\circ}$ N in agreement with EUVI observations. Previous statistics on CME position angles have shown that, during solar minimum, they tend to be offset closer to the equator as compared to those of the associated prominence eruption (Gopalswamy *et al.*, 2003). The non-radial motion we quantify here may be evidence of the drawn-out magnetic




Figure 5.18: Kinematic and morphological properties of the 3D reconstruction of the 12 December 2008 CME front. The top panel shows the velocity of the middle of the CME front with corresponding drag model and, inset, the early acceleration peak. Measurement uncertainties are indicated by one standard deviation errorbars. The middle panel shows the declinations from the ecliptic (0°) of an angular spread across the front between the CME flanks with a power-law fit indicative of non-radial propagation. The bottom panel shows the angular width of the CME with a power-law expansion.

dipole field of the Sun, an effect predicted at solar minimum due to the influence of the solar wind pressure (e.g. Figure 8 in Pneuman & Kopp (1971) and Figure 2 in Banaszkiewicz et al. (1998)). Other possible influences include changes to the internal current of the magnetic flux rope (Filippov et al., 2001), or the orientation of the magnetic flux rope with respect to the background field (Chané et al., 2005), whereby magnetic pressure can act asymmetrically to deflect the flux rope pole-ward or equator-ward depending on the field configurations.

5.5.4 CME Angular Width Expansion

Over the height range $2-46~\rm R_{\odot}$ the CME angular width ($\Delta\theta=\theta_{max}-\theta_{min}$) increases from $\sim 30^{\circ}$ to $\sim 60^{\circ}$ with a power-law of the form $\Delta\theta(r)=\Delta\theta_{0}r^{0.22}$ (2 $\rm R_{\odot}< r<46~\rm R_{\odot}$) (bottom panel of Figure 5.18). This angular expansion is evidence for an initial overpressure of the CME relative to the surrounding corona (coincident with its early acceleration inset in top panel of Figure 5.18). The expansion then tends to a constant during the later drag phase of CME propagation, as it expands to maintain pressure balance with heliocentric distance. It is theorised that the expansion may be attributed to two types of kinematic evolution, namely spherical expansion due to simple convection with the ambient solar wind in a diverging geometry, and expansion due to a pressure gradient between the flux rope and solar wind (Tappin, 2006). It is also noted that the southern portions of the CME manifest the bulk of this expansion below the ecliptic (best observed by comparing the relatively constant 'Midtop of Front' measurements with the more consistently decreasing 'Midbottom of Front' measurements in middle panel of Figure 5.18). Inspection of a Wang-Sheeley-Arge (WSA) solar wind model run

(Arge & Pizzo, 2000) reveals higher speed solar wind flows ($\sim 650 \; \mathrm{km \; s^{-1}}$) emanating from open-field regions at high/low latitudes (approximately 30° north/south of the solar equator). Once the initial prominence/CME eruption occurs and is deflected into a non-radial trajectory, it undergoes asymmetric expansion in the solar wind. It is prevented from expanding upwards into the open-field high-speed stream at higher latitudes, and the high internal pressure of the CME relative to the slower solar wind near the ecliptic accounts for its expansion predominantly to the south. In addition, the northern portions of the CME attain greater distances from the Sun than the southern portions as a result of this propagation in varying solar wind speeds, an effect predicted to occur in previous hydrodynamic models (Odstrčil & Pizzo, 1999).

5.5.5 CME Drag in the Inner Heliosphere

Investigating the midpoint kinematics of the CME front, we find the velocity profile increases from approximately $100-300~{\rm km~s^{-1}}$ over the first $2-5~{\rm R}_{\odot}$, before rising more gradually to a scatter between $400-550~{\rm km~s^{-1}}$ as it propagates outward (top panel of Figure 5.18). The acceleration peaks at approximately $100~{\rm m~s^{-2}}$ at a height of $\sim 3~{\rm R}_{\odot}$, then decreases to scatter about zero. This early phase is generally attributed to the Lorentz force whereby the dominant outward magnetic pressure overcomes the internal and/or external magnetic field tension, while the subsequent increase in velocity, at heights above $\sim 7~{\rm R}_{\odot}$, is predicted by theory to result from the effects of drag (Tappin, 2006). At large distances from the Sun, during this postulated drag-dominated epoch of CME propagation, the

equation of motion can be cast in the form:

$$M_{cme} \frac{dv_{cme}}{dt} = -\frac{1}{2} \rho_{sw} (v_{cme} - v_{sw}) |v_{cme} - v_{sw}| A_{cme} C_D$$
 (5.12)

This describes a CME of velocity v_{cme} , mass M_{cme} , and cross-sectional area A_{cme} propagating through a solar wind flow of velocity v_{sw} and density ρ_{sw} . The drag coefficient, C_D , is found to be of the order of unity for typical CME geometries (Cargill, 2004), while the density and area are expected to vary as power-law functions of distance R. Thus, we parameterise the density and geometric variation of the CME and solar wind using a power-law (Vršnak & Gopalswamy, 2002) to obtain:

$$\frac{dv_{cme}}{dR} = -\alpha R^{-\beta} \frac{1}{v_{cme}} \left(v_{sw} - v_{cme} \right)^{\gamma} \tag{5.13}$$

where γ describes the drag regime, which can be either viscous ($\gamma = 1$) or aerodynamic ($\gamma = 2$), and α and β are constants primarily related to the cross-sectional area of the CME and the density ratio of the solar wind flow to the CME (ρ_{sw}/ρ_{cme}). We determine a theoretical estimate of the CME velocity as a function of distance by numerically integrating this equation using a 4th order Runge-Kutta scheme and fitting the result to the observed velocities from $\sim 7-46~\rm R_{\odot}$. The initial CME height, CME velocity, asymptotic solar wind speed, and α , β , and γ are obtained from a bootstrapping procedure which provides a final best-fit to the observations and confidence intervals for the parameters. Best-fit values for α and β were found to be $(4.55^{+2.30}_{-3.27})\times 10^{-5}$ and $-2.02^{+1.21}_{-0.95}$ which agree with values found in previous modelling work (Vršnak, 2001). The best-fit value for the exponent of the velocity difference between the CME and the solar wind, γ , was found to be $2.27^{+0.23}_{-0.30}$, which is clear evidence that aerodynamic drag

 $(\gamma=2)$, and not viscous drag $(\gamma=1)$ acts during the propagation of the CME in interplanetary space.

5.5.6 CME Arrival Time

The drag model provides an asymptotic CME velocity of 555^{+114}_{-42} km s⁻¹ when extrapolated to 1 AU, which predicts the CME to arrive one day before the Advanced Composition Explorer (ACE) or WIND spacecrafts detect it at the L1 point. We investigate this discrepancy by using our 3D reconstruction to simulate the continued propagation of the CME from the Alfvén radius ($\sim 21.5~\rm R_{\odot}$) to Earth using the ENLIL with Cone Model (Xie et al., 2004) at NASA's Community Coordinated Modeling Center. ENLIL is a time-dependent 3D magnetohydrodynamic (MHD) code that models CME propagation through interplanetary space. An ideal fluid approximation is used to describe the solar wind plasma, under time-dependent MHD processes (neglecting microscopic processes). The plasma is treated as a fully ionised hydrogen gas with equal electron and proton densities ($n = n_e = n_p$) and temperatures ($T = T_e = T_p$), and the basic equations of MHD theory applied (such as outlined in Section 1.3.1).

We use the height, velocity, and width from our 3D reconstruction as initial conditions for the simulation, and find that the CME is actually slowed to $\sim 342~\rm km~s^{-1}$ at 1 AU. This is as a result of its interaction with an upstream, slow-speed, solar wind flow at distances beyond 50 R_{\odot}, as seen by inspection of the solar wind profile along the trajectory of the CME in the ENLIL simulation (Figure 5.19). This CME velocity is consistent with in-situ measurements of solar wind speed ($\sim 330~\rm km~s^{-1}$) from the ACE and WIND spacecraft at L1. Track-




Figure 5.19: The 3D CME front parameters are used as initial conditions for an ENLIL with Cone Model MHD simulation (Xie et al., 2004) and the output density (top) and velocity (bottom) profiles of the inner heliosphere are illustrated here for the time-stamp of 06:00 UT on 14 December 2008. Beyond distances of $\sim 50~{\rm R}_{\odot}$ the CME is slowed by its interaction with the upstream, slow-speed, solar wind flow along its trajectory towards Earth, and this accounts for its arrival time as detected in-situ by the ACE and WIND spacecraft at the L1 point near Earth.




Figure 5.20: The in-situ solar wind plasma and magnetic field data observed by the WIND spacecraft. From top to bottom the panels show proton density, bulk flow speed, proton temperature, and magnetic field strength and components. The red dashed lines indicate the arrival time of the density enhancement predicted from our ENLIL with Cone Model run providing 08:09 UT on 16 December 2008, with a potential offset error between our reconstruction and the derived model height profiles accounting for an arrival time up to 13:20 UT. We observe a magnetic cloud signature behind the front, as highlighted with blue dash-dotted lines.

ing the peak density of the CME front from the simulation gives an arrival time at L1 of $\sim 08:09$ UT on 16 December 2008. Accounting for the offset in CME front heights between our 3D reconstruction and ENLIL simulation at distances of 21.5 R_{\odot} < r < 46 R_{\odot} gives an arrival time in the range 08:09-13:20 UT on 16 December 2008. This prediction interval agrees well with the earliest derived arrival times of the CME front plasma pileup ahead of the magnetic cloud flux rope from the in-situ data of both ACE and WIND (Figure 5.20) before its subsequent impact at Earth (Davis *et al.*, 2009; Liu *et al.*, 2010).

5.5.7 CME 'Pancaking'

From the ENLIL simulation it is apparent that the CME undergoes an effect known as 'pancaking' whereby the middle portion of the CME may be slowed while the flanks of the CME maintain or increase speed such that the front distorts to become concave outwards in shape. This is illustrated in the density plot for the cross-section of the CME along the Sun-Earth line in top of Figure 5.19, and the effect increases with distance from the Sun. We investigate the curvature of the 3D CME front reconstruction along the Sun-Earth line in the distance $2-46~{\rm R}_{\odot}$ by fitting the front in this plane with an ellipse and inspecting the changing morphology with distance. A plot of this characterisation against height is shown in Figure 5.21 where it can be seen that the curvature of the front decreases as the CME propagates, with the front initially optimised by high-curvature horizontal ellipses, then becoming better optimised by more spherical ellipses, before finally being optimised by smoother vertical ellipses. The observations through the latter half of the HI1 into the HI2 fields-of-view also show this pancaking effect

on the CME (though geometrical and instrumental effects must be considered when interpreting these images).

5.6 Discussion & Conclusions

Since its launch, the dynamic twin-viewpoints of STEREO have enabled studies of the true propagation of CMEs in 3D space. Our new elliptical tie-pointing technique uses the curvature of the CME front as a necessary third constraint on the two viewpoints to build an optimum 3D reconstruction of the front. Here the technique is applied to an Earth-directed CME, to reveal numerous forces at play throughout its propagation.

The early acceleration phase results from the rapid release of energy when the CME dynamics are dominated by outward magnetic and gas pressure forces. Different models can reproduce the early acceleration profiles of CME observations though it is difficult to distinguish between them with absolute certainty (Lin et al., 2010b; Schrijver et al., 2008). For this event the acceleration phase coincides with a strong angular expansion of the CME in the low corona, which tends toward a constant in the later observed propagation in the solar wind. While, statistically, expansion of CMEs is a common occurrence (Bothmer & Schwenn, 1994), it is difficult to accurately determine the magnitude and rate of expansion across the 2D plane-of-sky images for individual events. Some studies of these single-viewpoint images of CMEs use characterisations such as the cone model (Xie et al., 2004; Xue et al., 2005; Zhao et al., 2002) but assume the angular width to be constant (rigid cone) which is not always true early in the events (Byrne et al., 2009; Gopalswamy et al., 2009a). Our 3D front reconstruction




Figure 5.21: Top: Ellipses characterising the 3D CME front reconstruction in the out-of-ecliptic plane along the Sun-Earth line. Bottom: The ellipse tilt angle is indicative of the initial effects of 'pancaking', as the curvature of the CME front decreases with increasing height due to its changing morphology in the solar wind.

overcomes the difficulties in distinguishing expansion from image projection effects, and we show that early in this event there is a non-constant, power-law, angular expansion of the CME. Theoretical models of CME expansion generally reproduce constant radial expansion, based on the suspected magnetic and gas pressure gradients between the erupting flux rope and the ambient corona and solar wind (Berdichevsky et al., 2003; Cargill et al., 2000; Odstrčil & Pizzo, 1999). To account for the angular expansion of the CME, a combination of internal overpressure relative to external gas and magnetic pressure drop-offs, along with convective evolution of the CME in the diverging solar wind geometry, must be considered (Riley & Crooker, 2004).

During this early phase evolution the CME is deflected from a high-latitude source region into a non-radial trajectory as indicated by the changing inclination angle (middle panel of Figure 5.18). While projection effects again hinder interpretations of CME position angles in single images, statistical studies show that, relative to their source region locations, CMEs have a tendency to deflect toward lower latitudes during solar minimum (Gopalswamy et al., 2003; Yashiro et al., 2004). It has been suggested that this results from the guiding of CMEs towards the equator by either the magnetic fields emanating from polar coronal holes (Kilpua et al., 2009; Xie et al., 2009) or the flow pattern of the background coronal magnetic field and solar wind/streamer influences (Cremades & Bothmer, 2004; MacQueen et al., 1986; Xie et al., 2009). Other models show that the internal configuration of the erupting flux rope can have an important effect on its propagation through the corona. The orientation of the flux rope, either normal or inverse polarity, will determine where magnetic reconnection is more likely to occur, and therefore change the magnetic configuration of the system to guide

the CME either equator- or pole-ward (Chané et al., 2005). Alternatively, modelling the filament as a toroidal flux rope located above a mid-latitude polarity inversion line results in non-radial motion and acceleration of the filament, due to the guiding action of the coronal magnetic field on the current motion (Filippov et al., 2001). Both of these models have a dependence on the chosen background magnetic field configuration, and so the suspected drawn-out magnetic dipole field of the Sun by the solar wind (Banaszkiewicz et al., 1998; Pneuman & Kopp, 1971) may be the dominant factor in deflecting the prominence/CME eruption into this observed non-radial trajectory.

At larger distances from the Sun (>7 R_{\odot}) the effects of drag become important as the CME velocity approaches that of the solar wind. The interaction between the moving magnetic flux rope and the ambient solar wind has been suggested to play a key role in CME propagation at large distances where the Lorentz driving force and the effects of gravity become negligible (Chen, 1996). Comparisons of initial CME speeds and in-situ detections of arrival times have shown that velocities converge on the solar wind speed (González-Esparza et al., 2003; Maloney et al., 2009). For this event we find that the drag force is indeed sufficient to accelerate the CME to the solar wind speed, and quantify that the kinematics are consistent with the quadratic regime of aerodynamic drag (turbulent, as opposed to viscous, effects dominate). The importance of drag becomes further apparent through the CME interaction with a slow-speed solar wind stream ahead of it, slowing it to a speed that accounts for the observed arrival time at L1 near Earth. This agrees with the conjecture that Sun-Earth transit time is more closely related to the solar wind speed than the initial CME speed (Vršnak et al., 2009). Other kinematic studies of this CME through the HI fields-of-view quote velocities of 411 ± 23 km s⁻¹ (Ahead) and 417 ± 15 km s⁻¹ (Behind) when assumed to have zero acceleration during this late phase of propagation (Davis *et al.*, 2009), or an average of 363 ± 43 km s⁻¹ when triangulated in time-elongation J-maps (Liu *et al.*, 2010). These speeds through the HI fields-of-view, lower than those quantified through the COR1/2 fields-of-view, agree somewhat with the deceleration of the CME to match the slow-speed solar wind ahead of it in our MHD simulation. Ultimately we are able to predict a more accurate arrival time of the CME front at L1.

A cohesive physical picture for how the CME erupts, propagates, and expands in the solar atmosphere remains to be fully developed and understood from a theoretical perspective. Realistic MHD models of the Sun's global magnetic field and solar wind are required to explain all processes at play, along with a need for adequate models of the complex flux rope geometries within CMEs. Additionally, ambitious space exploration missions, such as Solar Orbiter (ESA; Hassler *et al.*, 2009) and Solar Probe+ (NASA; McComas *et al.*, 2008), will be required to give us a better understanding of the fundamental plasma processes responsible for driving CMEs and determining their adverse effects at Earth.

Chapter 6

Conclusions & Future Work

This thesis has sought to increase our understanding of solar activity and its effects on Earth through a study of the phenomenon known as CMEs. This was undertaken by studying CME propagation with new data and techniques. This chapter presents the main results and conclusions, and outlines possible future directions for this work.

6.1 Principal Results

The primary objective of this study was to further our understanding of the kinematic and morphological evolution of CMEs as they propagate from the Sun into the heliosphere. This was done by applying new methods of multiscale image processing to CME observations in order to identify and track the CME front and characterise its propagation through coronagraph data. Following this, the development of a new elliptical tie-pointing technique applied to STEREO observations allowed a reconstruction of a CME front in 3D, overcoming projection

effects and revealing its true 3D motion. The principal results arising from these studies may be summarised as follows:

- 1. The multiscale nature of CMEs was revealed through the application of a high and low pass image filtering technique (Young & Gallagher, 2008). The multiscale filtering was shown to effectively remove noise and small-scale features in order to reveal CME structure on a scale that best identifies the CME front. The specific implementation of multiscale filtering introduced by Young & Gallagher (2008) also allowed the chaining of pixels along edges in the decomposed images to reveal the CME front for tracking through time. Such an algorithm provides a robustness in CME front detection that alleviates issues of subjective user biases and unreproducibility of results. The technique is also more accurate than running-difference which is widely used for determining CME heights, both spatially since it requires no arbitrary scaling and/or thresholding to find the edges, and temporally as it operates on individual images without a need for subtracting antecedent frames. The technique was also extended for use as a potentially automated CME front detection algorithm, discussed further in Section 6.2.1 below.
- 2. An ellipse characterisation of the CME front in coronagraph data was implemented and shown to be an effective method for retrieving information on the changing CME morphology through sequences of observations. An ellipse was chosen for the innate freedom in its parameters, having the ability to best fit the varying curvature of CME fronts across different image sequences, while still being constrained to close upon itself. The characterisation provided a robust method for obtaining CME heights since it is not

affected by deviations along the often ill-defined and/or kinked CME front. The opening cone angle to the ellipse also provided a measure of angular width for testing CME expansion, and the eccentricity of the ellipse provided information on the changing CME front curvature as it was observed to propagate across the plane-of-sky.

- 3. The degree of accuracy provided by the multiscale methods and characterisation of the CME front allowed a test of the constant acceleration model upon a variety of CMEs, and it was found not to be true of all cases. This has implications for how CMEs may be modelled theoretically since the forces acting must have different regimes of dominance to cause non-constant acceleration in certain events. This warrants further investigation, especially if projection effects can be corrected for using STEREO data, and the kinematics determined with the best affordable accuracy (see the discussion of Sections 6.2.3 and 6.2.4 below).
- 4. The accuracy of these methods further revealed an early acceleration phase for some CMEs, and those with high speeds tended to reach greater angular widths, indicative of an initially dominant outward driving force ramping up the CME speed and expansion within the first few solar radii. Testing for this outward force that drives the CME acceleration and causes it to expand, is vitally important for comparing with theory in an effort to understand the interplay of forces acting on the CME as it propagates through the corona and heliosphere.
- 5. The newly developed elliptical tie-pointing technique was shown to overcome the plane-of-sky projection effects of previous single vantage-point

observations in a manner better suited to studying the CME of 12 December 2008 than previous stereoscopic efforts. By using the characterised curvature of the CME front in the observations from the twin-viewpoints of STEREO it was possible to constrain a reconstruction of the true CME front between the two planes-of-sky. This revealed the CME's true 3D motion, thus greatly increasing the accuracy with which we can interpret its kinematics and morphology.

- 6. The acceleration phase of the prominence eruption and CME was determined in 3D to occur within the first 3 R_☉, with the speed of the outer CME front being higher than the speed of the prominence that forms the inner core material of the CME. Their different speeds are indicative of the CME expansion, and indeed the early phase acceleration corresponds to the event's strongest characterised (out-of-ecliptic) angular width increase. This is again indicative of an initially dominant outward driving force that causes the CME speed and width to increase dramatically (generally considered to be the Lorentz magnetic force of CME initiation and propagation).
- 7. The deflected motion of the prominence eruption and CME was quantified in 3D; originating at high solar latitudes of $\sim 55^{\circ}$ but determined to move on a trajectory along the ecliptic at $\sim 0^{\circ}$. This is indicative of the drawn-out background magnetic field of the quiet Sun. The pressure of the solar wind acts to drag out field lines in the $\beta > 1$ coronal regime, influencing the overall magnetic field configuration right down to the surface of the quiet Sun. At these low heights the magnetic pressure of the drawn-out field guides the motion of the gradual eruption into a non-radial trajectory,

placing it on a course towards Earth. This highlights the importance of the overall solar magnetic field with respect to the CME and its source region, and reveals how necessary an understanding of their interdependence is, both in the context of CME physics and space weather monitoring.

8. The effects of drag on the 3D CME motion were investigated and it was found that, subsequent to the early acceleration phase of the CME, its speed continued to increase towards the speed of the ambient solar wind. A comparison of the CME velocity with the in-situ detection of its arrival at L1 implied the CME had to be further slowed along its trajectory, revealed to be true when investigated with the 3D MHD 'ENLIL with Cone' Model. This highlights the importance of drag on the CME throughout its propagation as it was found to undergo further deceleration by the slow-speed solar wind stream ahead of it as it propagated through interplanetary space. Understanding these drag effects is thus of great importance for predicting CME arrival times at Earth and improving space weather forecasts.

6.2 Future Work

The methods developed and implemented in this thesis are a first use of multiscale analysis and characterisation in obtaining CME kinematics and morphology with better constrained errors than previous efforts, of benefit in testing theoretical CME models and forecasting space weather at Earth or elsewhere in the heliosphere. The possibility for automation has been demonstrated and would provide a new catalogue of CMEs working in realtime with SolarMonitor.org for example. Extending these multiscale techniques to curvelets or ridgelets may better

suit the detection of the typically curved nature of CMEs in coronagraph data. Furthermore, the data from the twin viewpoints of the STEREO mission have allowed us to overcome plane-of-sky projection effects in studying CMEs with our newly developed elliptical tie-pointing technique, resulting in a very cohesive description of how the CME of 12 December 2008 propagates from the Sun to the Earth. There are numerous candidate events which should also be studied in this manner to test the conclusions drawn from the results of the CME studied here and gain further insight into CME dynamics. Other methods for deriving the velocity and acceleration profiles of CMEs also warrant investigation, since the 3-point Lagrangian interpolation is sensitive to scatter in the data (though less so than the standard forward/reverse difference) and bootstrapping or inversion techniques for example may help overcome this.

6.2.1 Automation

As discussed in Chapter 3 there is great benefit in implementing an automated CME detection and tracking algorithm for cataloguing their kinematics and similar properties of interest in large data sets. The algorithms (specifically CACTus, SEEDS and ARTEMIS) remove the need for a manual inspection of the images (as performed in the CDAW catalogue), which can be both laborious and suffer from user-specific biases. The automated techniques also produce a robust output of parameters for large statistical analyses of CME properties over long periods of solar activity, since the thresholds for detection and height/width measurements are hard-coded into the algorithm. However, the ultimate aim is to determine the kinematics and morphology of CMEs with as great an accuracy as possible




Figure 6.1:

with the available data, thus operations such as image rebinning, smoothing, and differencing are not ideal. Also, accurately measuring the CME height (and subsequently deriving the velocity and acceleration) can be difficult since the CME front is often diffuse and ill-defined. These issues motivated the application of multiscale filtering to enhance the CME front in coronagraph images without reducing the quality of the image since noise and small-scale features can be removed in the multiscale decomposition for optimum CME detection. The multiscale technique outlined in Chapter 3 also results in an edge detection on the image akin to the Canny edge detector as a result of the horizontal and vertical directions in which the multiscale filter is applied. CME detection is shown to be




Figure 6.2: An example interface which could be developed for the potentially automated multiscale filtering and ellipse characterisation for tracking CMEs observed by STEREO-A and B. Images from the respective instruments appear on the left, and the resulting parameters from the characterisation appear on the right.

effective through the thresholding of angular spreads across the edge detections of the image, since CMEs will have a large distribution of edge normals compared with the generally radial nature of the corona and structures within (specifically streamers). However, the algorithm currently lacks a satisfactory image segmentation technique for discarding the regions of an image that do not correspond to the detected CME and thus still requires a user to perform an inspection of the edges to be maintained/discarded in the images for characterisation. Including multiple scales of the decomposition, rather than just the one with the best signal-to-noise ratio for the CME, allows a scoring system to refine the CME detection mask, although cases of streamer interaction or deflection are prone to skew the resulting detection. The algorithm needs to be improved with a form of image segmentation included to make the CME detection masks more robust, and minimise the effects of streamers. Then the ellipse characterisation can be automatically applied for studying the CME front kinematics and morphology with greater accuracy, and produce a multiscale-methods based catalogue of events (e.g. Figure 6.2).

6.2.2 Ridgelets/Curvelets

The implementation of multiscale analysis has been demonstrated for its efficacy in highlighting CMEs and coronal structure against noise and small-scale features in coronagraph images (Byrne *et al.*, 2009; Stenborg & Cobelli, 2003; Stenborg *et al.*, 2008; Young & Gallagher, 2008). However, wavelets are better suited to identifying point-like features in images, but their extension to ridgelets and curvelets has been shown to better resolve the visibility of the curved form of a




Figure 6.3: An example ridgelet, reproduced from Gallagher *et al.* (2010). The first graph shows a typical ridgelet, and the second to fourth graphs are obtained from simple geometrical manipulations, namely rotation, rescaling and shifting.

typical CME front (Gallagher *et al.*, 2010). Thus their development may provide a more reliable CME detection algorithm than the multiscale technique investigated in this work.

The continuous wavelet transform of an image may be defined as:

$$w(s, a, b) = \int \int I(x, y) \psi_{s,a,b}(x, y) dx dy$$
(6.1)

where w(s, a, b) are the wavelet coefficients of the image I(x, y), $\psi_s(x, y)$ is the mother wavelet, and s is the term describing scale at a position (a, b). The mother wavelet can take many forms, e.g., the Morlet wavelet or the Mexican hat




Figure 6.4: The curvelet filtered image of a CME observed by LASCO/C2 on 18 April 2000, reproduced from Gallagher *et al.* (2010). The detail along the curved CME front is enhanced as a result of the curvelet technique following the removal of certain coefficients probably due to noise.

wavelet. The ridgelet transform takes a similar mathematical form to the wavelet transform, i.e., it is a convolution of an image with a predefined basis function, but they are anisotropic and thus more directionally sensitive (Figure 6.3). The ridgelet uses a radon transform that transforms lines into points, upon which a wavelet transform may be applied to provide a sparse representation of the points. The basis function of the ridgelet takes the form:

$$\psi_{s,b,\theta}(x,y) = s^{-1/2}\psi\left(\frac{x\cos\theta + y\sin\theta - b}{s}\right)$$
 (6.2)

The ridgelet is constant along lines $x \cos \theta + y \sin \theta = const.$ The ridgelet coefficients are given by the convolution:

$$R_{I}(s,b,\theta) = \int \int I(x,y) \psi_{s,b,\theta}(x,y) dx dy$$
(6.3)




Figure 6.5:

Curvelets generalise the idea of ridgelets to multiscale curves in images. The detailed maths on the application of curvelets to CME images is discussed in Gallagher *et al.* (2010) based on the developments by Candés & Donoho (1999). Figure 6.4 shows how well the curvelet filter performs on an image of a CME.

6.2.3 3D CME Reconstructions

The development of the elliptical tie-pointing technique has proven effective at reconstructing the CME front of the 12 December 2008 event, discussed in detail in Chapter 5. This event was an ideal case for study since the STEREO space-craft were at an angular separation of almost 90°, so the lines-of-sight intersected to form optimum quadrilaterals localising the CME front. Very small, or very large, angles of separation would not be as effective since the quadrilaterals will be skewed and so too will the corresponding inscribed ellipses. So while a recon-

struction would match the observations of both spacecraft it may not perfectly represent the true curvature of the CME front (although it will still offer a better approximation than tie-pointing alone). The fact that the 12 December 2008 CME propagated along the Sun-Earth line midway between the two spacecraft meant the reconstruction could be performed out to heights of almost 50 R_{\odot} before the 'pancaking' of the CME (and concerns on the scattering geometry and instrumental effects) meant the ellipse fit was no longer appropriate. If these concerns could be corrected for, or simply included in the uncertainty, then it is possible to use more than one ellipse fit to characterise the different portions of the CME front and perform elliptical tie-pointing on these to gain an insight into the 3D propagation through the rest of the HI1 field-of-view. It would also be interesting to test how the method fares with the observations of HI2. A number of other CMEs will warrant studying with the elliptical tie-pointing technique throughout the lifetime of the STEREO mission. Many events have been observed through the COR1 and COR2 fields of view and been studied by several authors through a variety of stereoscopic methods (Boursier et al., 2009a; de Koning et al., 2009; Liewer et al., 2009; Mierla et al., 2008; Srivastava et al., 2009; Temmer et al., 2009; Wood et al., 2009), and so direct comparisons may be made with the results of these studies and future stereoscopic CME analyses.

To potentially reveal some of the inner structure of CMEs, the polarisation technique of Moran & Davila (2004) could be used in conjunction with the elliptical tie-pointing reconstruction. The technique relies upon the geometric dependence of the polarisation of Thomson-scattered light, whereby the polarisation fraction in CME emission provides a line-of-sight averaged 'mean distance to the plane of the sky' for selected or all CME pixels. The validity of the method was




Figure 6.6: A theoretical model for a CME with constant acceleration 2 m s^{-2} and initial velocity 300 km s^{-1} , and two simulations of how the resulting profiles for a noisy sample of datapoints behave using 3-point Lagrangian interpolation.

tested and proven using STEREO observations of a CME on 21 August 2007 and shown to be in good agreement with other triangulation methods (Moran et al., 2010). A combined use of the technique with the geometric localisation on COR2 beacon data has been explored by De Koning et al. (2009), although the polarisation techniques becomes untrustworthy at heights $\gtrsim 5~\rm R_{\odot}$ (Billings, 1966).

6.2.4 Deriving CME Kinematics

The standard method for determining the kinematics of a CME is to obtain a sequence of height-time measurements and perform a 3-point Lagrangian interpolation to derive the velocity and acceleration of the event (detailed in Section 4.2.1). This method alone is somewhat dated since the advent of strong numerical computing power and development of bootstrapping and spline-fitting techniques for example. As a first approximation the 3-point Lagrangian provides a good estimate of how the velocity and acceleration cooresponding to a height-time curve are likely to behave, by revealing the trends in the profiles that indicate increasing/decreasing velocity and/or acceleration. However, this is only true if the scatter and errors in the height measurements are not unreasonably large, since a large scatter would be enhanced by the derivatives and a large error will increase the uncertainty on the derivative points such that trends may become untestable. A quick simulation of how the 3-point Lagrangian fares for a theoretical model CME undergoing a constant acceleration of 2 m s^{-2} and initial velocity of 300 km s⁻² reveals the potential unreliability of the resulting kinematic profiles. A scatter of height-time datapoints is chosen with varying levels of noise upto $\sim 20\%$. An errorbar on each datapoint is determined by its distance from the theoretical height-time profile. Various instances of datapoint scatters result in erroneous trends in the velocity and acceleration profiles - even with the proper error treatment. Figure 6.6 shows two examples of how different the derived kinematics can be, in comparison with the theoretical model. They show how different scatters of the datapoints can result in what appear to be completely opposing acceleration trends, meaning the nature of the scatter is




Figure 6.7: The resulting velocity profile for the 3D reconstructed CME front of the 12 December 2008 using the inversion technique of Kontar & MacKinnon (2005).

not satisfactorily reflected in the derived errorbars of the resulting profiles. This warrants further investigation for future CME, and other, kinematic studies.

It should also be noted that the 3-point Lagrangian counter-intuitively increases the errorbars on the resulting velocity and acceleration profiles when the number of height-time measurements are increased (i.e. for a smaller cadence) due to the algorithm's inverse dependance on the spacing between points. It is therefore worth investigating how other techniques might be applied to more confidently derive the kinematics of CMEs. Bootstrapping of a presumed model fit to the CME height-time profile would result in the best match of parameters to the data, but questions would remain on the appropriateness of the chosen model itself. Alternatively a simulation of data could be bootstrapped regarding the resulting derived kinematics and an estimate of the errors involved could be deduced to apply to true observations, though this again may be model dependant. An inversion technique could also be investigated for obtaining derivatives

(Kontar & MacKinnon, 2005), an example of which is shown in deriving the velocity of the 12 December 2008 CME (Figure 6.7). It works by essentially solving for the smoothest spline fit that minimises the distance between the end-points while still being bound by any constraints on the data. If the quantification of the height-time errors are sound, then inversion techniques may provide a more robust determination of the kinematic profiles.

References

- Antiochos, S.K., Devore, C.R. & Klimchuk, J.A. (1999). A Model for Solar Coronal Mass Ejections. *Astrophysical Journal*, **510**, 485–493.
- Arge, C.N. & Pizzo, V.J. (2000). Improvement in the prediction of solar wind conditions using near-real time solar magnetic field updates. *Journal of Geophysical Research*, **105**, 10465–10480.
- Aschwanden, M.J. (2005). Physics of the Solar Corona. An Introduction with Problems and Solutions (2nd edition).
- ASCHWANDEN, M.J., WÜLSER, J.P., NITTA, N.V. & LEMEN, J.R. (2008). First Three-Dimensional Reconstructions of Coronal Loops with the STEREO A and B Spacecraft. I. Geometry. *Astrophysical Journal*, **679**, 827–842.
- BABCOCK, H.W. (1961). The Topology of the Sun's Magnetic Field and the 22-YEAR Cycle. Astrophysical Journal, 133, 572-+.
- Bahcall, J.N. (1989). Neutrino astrophysics.
- Banaszkiewicz, M., Axford, W.I. & McKenzie, J.F. (1998). An analytic solar magnetic field model. *Astronomy & Astrophysics*, **337**, 940–944.

- Berdichevsky, D.B., Lepping, R.P. & Farrugia, C.J. (2003). Geometric considerations of the evolution of magnetic flux ropes. *Physical Review E*, **67**, 036405–+.
- BIERMANN, L. (1948). Über die Ursache der chromosphärischen Turbulenz und des UV-Exzesses der Sonnenstrahlung. Zeitschrift fr Astrophysik, 25, 161-+.
- BILLINGS, D.E. (1966). A guide to the solar corona.
- BISKAMP, D. (1986). Magnetic Reconnection Via Current Sheets (Invited paper).
 In M. A. Dubois, D. Grésellon, & M. N. Bussac, ed., Magnetic Reconnection and Turbulence, 19-+.
- BOTHMER, V. & SCHWENN, R. (1994). Eruptive prominences as sources of magnetic clouds in the solar wind. *Space Science Reviews*, **70**, 215–220.
- Boursier, Y., Lamy, P. & Llebaria, A. (2009a). Three-Dimensional Kinematics of Coronal Mass Ejections from STEREO/SECCHI-COR2 Observations in 2007-2008. *Solar Physics*, **256**, 131–147.
- Boursier, Y., Lamy, P., Llebaria, A., Goudail, F. & Robelus, S. (2009b). The ARTEMIS Catalog of LASCO Coronal Mass Ejections. Automatic Recognition of Transient Events and Marseille Inventory from Synoptic maps. *Solar Physics*, **257**, 125–147.
- Brueckner, G.E., Howard, R.A., Koomen, M.J., Korendyke, C.M., Michels, D.J., Moses, J.D., Socker, D.G., Dere, K.P., Lamy, P.L., Llebaria, A., Bout, M.V., Schwenn, R., Simnett, G.M., Bedford,

- D.K. & EYLES, C.J. (1995). The Large Angle Spectroscopic Coronagraph (LASCO). Solar Physics, **162**, 357–402.
- Byrne, J.P., Gallagher, P.T., McAteer, R.T.J. & Young, C.A. (2009). The kinematics of coronal mass ejections using multiscale methods. *Astronomy & Astrophysics*, **495**, 325–334.
- Byrne, J.P., Maloney, S.A., McAteer, R.T.J., Refojo, J.M. & Gal-Lagher, P.T. (2010). Propagation of an Earth-directed coronal mass ejection in three-dimensions. *Nat. Commun.*, ?
- Calabretta, M.R. & Greisen, E.W. (2002). Representations of celestial coordinates in FITS. Astronomy & Astrophysics, **395**, 1077–1122.
- Candés, E.J. & Donoho, D.L. (1999). Ridgelets: a key to higher-dimensional intermittency? *Philosophical Transactions of The Royal Society*, **357**, 2495–2509.
- CARGILL, P.J. (2004). On the Aerodynamic Drag Force Acting on Interplanetary Coronal Mass Ejections. *Solar Physics*, **221**, 135–149.
- Cargill, P.J., Chen, J., Spicer, D.S. & Zalesak, S.T. (1996). Magnetohydrodynamic simulations of the motion of magnetic flux tubes through a magnetized plasma. *Journal of Geophysical Research*, **101**, 4855–4870.
- Cargill, P.J., Schmidt, J., Spicer, D.S. & Zalesak, S.T. (2000). Magnetic structure of overexpanding coronal mass ejections: Numerical models. *Journal of Geophysical Research*, **105**, 7509–7520.

- Carrington, R.C. (1863). Observations of the Spots on the Sun. London: Williams & Norgate.
- CHANÉ, E., JACOBS, C., VAN DER HOLST, B., POEDTS, S. & KIMPE, D. (2005). On the effect of the initial magnetic polarity and of the background wind on the evolution of CME shocks. *Astronomy & Astrophysics*, **432**, 331–339.
- CHAPMAN, S. & ZIRIN, H. (1957). Notes on the Solar Corona and the Terrestrial Ionosphere. *Smithsonian Contributions to Astrophysics*, **2**, 1–+.
- Chen, J. (1996). Theory of prominence eruption and propagation: Interplanetary consequences. *Journal of Geophysical Research*, **101**, 27499–27520.
- Chen, J. & Krall, J. (2003). Acceleration of coronal mass ejections. *Journal* of Geophysical Research (Space Physics), 108, 1410–+.
- Chen, J., Howard, R.A., Brueckner, G.E., Santoro, R., Krall, J., Paswaters, S.E., St. Cyr, O.C., Schwenn, R., Lamy, P. & Simnett, G.M. (1997). Evidence of an Erupting Magnetic Flux Rope: LASCO Coronal Mass Ejection of 1997 April 13. Astrophysical Journal Letters, 490, L191+.
- CREMADES, H. & BOTHMER, V. (2004). On the three-dimensional configuration of coronal mass ejections. *Astronomy & Astrophysics*, **422**, 307–322.
- DAL LAGO, A., SCHWENN, R. & GONZALEZ, W.D. (2003). Relation between the radial speed and the expansion speed of coronal mass ejections. *Advances in Space Research*, **32**, 2637–2640.

- Davis, C.J., Davies, J.A., Lockwood, M., Rouillard, A.P., Eyles, C.J. & Harrison, R.A. (2009). Stereoscopic imaging of an Earth-impacting solar coronal mass ejection: A major milestone for the STEREO mission. *Geophysical Research Letters*, **36**, 8102–+.
- DAVIS, C.J., KENNEDY, J. & DAVIES, J.A. (2010). Assessing the Accuracy of CME Speed and Trajectory Estimates from STEREO Observations Through a Comparison of Independent Methods. *Solar Physics*, **263**.
- DE KONING, C.A., PIZZO, V.J. & BIESECKER, D.A. (2009). Calculating CME Velocity in Near-Real-Time Using Geometric and Polarimetric Techniques. In AAS/Solar Physics Division Meeting, vol. 40 of AAS/Solar Physics Division Meeting, 1609.
- DE KONING, C.A., PIZZO, V.J. & BIESECKER, D.A. (2009). Geometric Localization of CMEs in 3D Space Using STEREO Beacon Data: First Results. Solar Physics, 256, 167–181.
- DÉMOULIN, P., NAKWACKI, M.S., DASSO, S. & MANDRINI, C.H. (2008). Expected in Situ Velocities from a Hierarchical Model for Expanding Interplanetary Coronal Mass Ejections. *Solar Physics*, **250**, 347–374.
- DEVORE, C.R. & Antiochos, S.K. (2008). Homologous Confined Filament Eruptions via Magnetic Breakout. *Astrophysical Journal*, **680**, 740–756.
- Domingo, V., Fleck, B. & Poland, A.I. (1995). The SOHO Mission: an Overview. *Solar Physics*, **162**, 1–2.

- Eyles, C.J., Harrison, R.A., Davis, C.J., Waltham, N.R., Shaughnessy, B.M., Mapson-Menard, H.C.A., Bewsher, D., Crothers, S.R., Davies, J.A., Simnett, G.M., Howard, R.A., Moses, J.D., Newmark, J.S., Socker, D.G., Halain, J.P., Defise, J.M., Mazy, E. & Rochus, P. (2009). The Heliospheric Imagers Onboard the STEREO Mission. *Solar Physics*, **254**, 387–445.
- FILIPPOV, B.P., GOPALSWAMY, N. & LOZHECHKIN, A.V. (2001). Non-radial motion of eruptive filaments. *Solar Physics*, **203**, 119–130.
- FLIGGE, M. & SOLANKI, S.K. (1997). Noise reduction in astronomical spectra using wavelet packets. Astronomy & Astrophysics Supplemental, 124, 579–587.
- Forbes, T.G. & Isenberg, P.A. (1991). A catastrophe mechanism for coronal mass ejections. *Astrophysical Journal*, **373**, 294–307.
- FORBES, T.G. & PRIEST, E.R. (1995). Photospheric Magnetic Field Evolution and Eruptive Flares. *Astrophysical Journal*, **446**, 377–+.
- GABRIEL, A.H. & MASON, H.E. (1982). Solar physics. In H. S. W. Massey & D. R. Bates, ed., Applied Atomic Collision Physics, Volume 1: Atmospheric Physics and Chemistry, vol. 1, 345–397.
- Gallagher, P.T., Lawrence, G.R. & Dennis, B.R. (2003). Rapid Acceleration of a Coronal Mass Ejection in the Low Corona and Implications for Propagation. *Astrophysical Journal Letters*, **588**, L53–L56.

- Gallagher, P.T., Young, C.A., Byrne, J.P. & McAter, R.T.J. (2010). Coronal Mass Ejection Detection using Wavelets, Curvelets and Ridgelets: Applications for Space Weather Monitoring. *Advances in Space Research*.
- GARY, G.A. (2001). Plasma Beta above a Solar Active Region: Rethinking the Paradigm. *Solar Physics*, **203**, 71–86.
- Gold, T. (1962). Magnetic Storms. Space Science Reviews, 1, 100–114.
- González-Esparza, J.A., Lara, A., Pérez-Tijerina, E., Santillán, A. & Gopalswamy, N. (2003). A numerical study on the acceleration and transit time of coronal mass ejections in the interplanetary medium. *Journal of Geophysical Research (Space Physics)*, **108**, 1039–+.
- Goossens, M., ed. (2003). An introduction to plasma astrophysics and magnetohydrodynamics, vol. 294 of Astrophysics and Space Science Library.
- Gopalswamy, N., Lara, A., Lepping, R.P., Kaiser, M.L., Berdichevsky, D. & St. Cyr, O.C. (2000). Interplanetary acceleration of coronal mass ejections. *Geophysical Research Letters*, **27**, 145–148.
- Gopalswamy, N., Shimojo, M., Lu, W., Yashiro, S., Shibasaki, K. & Howard, R.A. (2003). Prominence Eruptions and Coronal Mass Ejection: A Statistical Study Using Microwave Observations. *Astrophysical Journal*, **586**, 562–578.
- Gopalswamy, N., Dal Lago, A., Yashiro, S. & Akiyama, S. (2009a). The Expansion and Radial Speeds of Coronal Mass Ejections. *Central European Astrophysical Bulletin*, **33**, 115–124.

- Gopalswamy, N., Yashiro, S., Michalek, G., Stenborg, G., Vourlidas, A., Freeland, S. & Howard, R. (2009b). The SOHO/LASCO CME Catalog. *Earth Moon and Planets*, **104**, 295–313.
- HASSLER, D., HORBURY, T., LEWIS, W., MASON, G., OWEN, C., SOLANKI, S., VELLI, M. & WIMMER-SCHWEINGRUBER, R. (2009). Solar Orbiter: Exploring the Sun-Heliosphere Connection. *ESA/SRE(2009)5*.
- HEWETT, R.J., GALLAGHER, P.T., MCATEER, R.T.J., YOUNG, C.A., IRE-LAND, J., CONLON, P.A. & MAGUIRE, K. (2008). Multiscale Analysis of Active Region Evolution. *Solar Physics*, **248**, 311–322.
- HORWITZ, A. (2002). Finding ellipses and hyperbolas tangent to two, three, or four given lines. Southwest Journal of Pure and Applied Mathematics, 1, 6–32.
- HORWITZ, A. (2005). Ellipses of maximal area and minimal eccentricity inscribed in a convex quadrilateral. The Australian Journal of Mathematical Analysis and Applications, 2, 1–12.
- HOWARD, R.A., MOSES, J.D. & SOCKER, D.G. (2000). Sun-Earth connection coronal and heliospheric investigation (SECCHI). In S. Fineschi, C. M. Korendyke, O. H. Siegmund, & B. E. Woodgate, ed., Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol. 4139 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 259–283.
- HOWARD, R.A., MOSES, J.D., VOURLIDAS, A., NEWMARK, J.S., SOCKER, D.G., PLUNKETT, S.P., KORENDYKE, C.M., COOK, J.W., HURLEY, A., DAVILA, J.M., THOMPSON, W.T., ST CYR, O.C., MENTZELL, E.,

Mehalick, K., Lemen, J.R., Wuelser, J.P., Duncan, D.W., Tarbell, T.D., Wolfson, C.J., Moore, A., Harrison, R.A., Waltham, N.R., Lang, J., Davis, C.J., Eyles, C.J., Mapson-Menard, H., Simnett, G.M., Halain, J.P., Defise, J.M., Mazy, E., Rochus, P., Mercier, R., Ravet, M.F., Delmotte, F., Auchere, F., Delaboudiniere, J.P., Bothmer, V., Deutsch, W., Wang, D., Rich, N., Cooper, S., Stephens, V., Maahs, G., Baugh, R., McMullin, D. & Carter, T. (2008a). Sun Earth Connection Coronal and Heliospheric Investigation (SECCHI). Space Science Reviews, 136, 67–115.

- HOWARD, T.A. & TAPPIN, S.J. (2005). Statistical survey of earthbound interplanetary shocks, associated coronal mass ejections and their space weather consequences. *Astronomy & Astrophysics*, **440**, 373–383.
- HOWARD, T.A. & TAPPIN, S.J. (2008). Three-Dimensional Reconstruction of Two Solar Coronal Mass Ejections Using the STEREO Spacecraft. *Solar Physics*, **252**, 373–383.
- HOWARD, T.A. & TAPPIN, S.J. (2009). Interplanetary Coronal Mass Ejections Observed in the Heliosphere: 1. Review of Theory. *Space Science Reviews*, **147**, 31–54.
- Howard, T.A., Nandy, D. & Koepke, A.C. (2008b). Kinematic properties of solar coronal mass ejections: Correction for projection effects in spacecraft coronagraph measurements. *Journal of Geophysical Research (Space Physics)*, **113**, 1104–+.

- Hundhausen, A.J., Sawyer, C.B., House, L., Illing, R.M.E. & Wagner, W.J. (1984). Coronal mass ejections observed during the solar maximum mission Latitude distribution and rate of occurrence. *Journal of Geophysical Research*, 89, 2639–2646.
- ILLING, R.M.E. & HUNDHAUSEN, A.J. (1985). Observation of a coronal transient from 1.2 to 6 solar radii. *Journal of Geophysical Research*, **90**, 275–282.
- INHESTER, B. (2006). Stereoscopy basics for the STEREO mission. ArXiv Astrophysics e-prints.
- Isenberg, P.A., Forbes, T.G. & Demoulin, P. (1993). Catastrophic Evolution of a Force-free Flux Rope: A Model for Eruptive Flares. *Astrophysical Journal*, 417, 368–+.
- Jackson, J.D. (1975). Classical electrodynamics.
- Jacobs, C., Roussev, I.I., Lugaz, N. & Poedts, S. (2009). The Internal Structure of Coronal Mass Ejections: Are all Regular Magnetic Clouds Flux Ropes? *Astrophysical Journal Letters*, **695**, L171–L175.
- Kaiser, M.L., Kucera, T.A., Davila, J.M., St. Cyr, O.C., Guhathakurta, M. & Christian, E. (2008). The Stereo Mission: An Introduction. *Space Science Reviews*, **136**, 5–16.
- KIEPENHEUER, K.O. (1953). Solar Activity, 322-+.
- KILPUA, E.K.J., POMOELL, J., VOURLIDAS, A., VAINIO, R., LUHMANN, J., LI, Y., SCHROEDER, P., GALVIN, A.B. & SIMUNAC, K. (2009). Stereo

- observations of interplanetary coronal mass ejections and prominence deflection during solar minimum period. *Annales Geophysicae*, **27**, 4491–4503.
- Kimura, H. & Mann, I. (1998). Brightness of the solar F-corona. Earth, Planets, and Space, **50**, 493–499.
- KITCHIN, C.R. (1987). Stars, nebulae and the interstellar medium. Observational physics and astrophysics.
- KLIEM, B. & TÖRÖK, T. (2006). Torus Instability. *Physical Review Letters*, **96**, 255002-+.
- KLIMCHUK, J.A. (2001). Theory of Coronal Mass Ejections. Space Weather (Geophysical Monograph 125), ed. P. Song, H. Singer, G. Siscoe (Washington: Am. Geophys. Un.), 143 (2001), 125, 143-+.
- Kontar, E.P. & Mackinnon, A.L. (2005). Regularized Energy-Dependent Solar Flare Hard X-Ray Spectral Index. *Solar Physics*, **227**, 299–310.
- Koomen, M.J., Detwiler, C.R., Brueckner, G.E., Cooper, H.W. & Tousey, R. (1975). White Light Coronagraph in OSO-7. *Applied Optics*, **14**, 743–751.
- KOUTCHMY, S., ZIRKER, J.B., STEINOLFSON, R.S. & ZHUGZHDA, I.D. (1991). Coronal activity, 1044–1086.
- Krall, J. & St. Cyr, O.C. (2006). Flux-Rope Coronal Mass Ejection Geometry and Its Relation to Observed Morphology. *Astrophysical Journal*, **652**, 1740–1746.

- Krall, J., Chen, J., Duffin, R.T., Howard, R.A. & Thompson, B.J. (2001). Erupting Solar Magnetic Flux Ropes: Theory and Observation. *Astrophysical Journal*, **562**, 1045–1057.
- Krall, J., Yurchyshyn, V.B., Slinker, S., Skoug, R.M. & Chen, J. (2006). Flux Rope Model of the 2003 October 28-30 Coronal Mass Ejection and Interplanetary Coronal Mass Ejection. Astrophysical Journal, 642, 541–553.
- KULSRUD, R.M. (2001). Magnetic reconnection: Sweet-Parker versus Petschek.
 Earth, Planets, and Space, 53, 417–422.
- LIEWER, P.C., DE JONG, E.M., HALL, J.R., HOWARD, R.A., THOMPSON, W.T., CULHANE, J.L., BONE, L. & VAN DRIEL-GESZTELYI, L. (2009). Stereoscopic Analysis of the 19 May 2007 Erupting Filament. *Solar Physics*, **256**, 57–72.
- LIN, C., GALLAGHER, P.T. & RAFTERY, C.L. (2010a). Investigating the driving mechanisms of coronal mass ejections. *Astronomy & Astrophysics*, **516**, A44+.
- LIN, C., GALLAGHER, P.T. & RAFTERY, C.L. (2010b). Investigating the driving mechanisms of coronal mass ejections. *ArXiv e-prints*.
- Lin, J. & Forbes, T.G. (2000). Effects of reconnection on the coronal mass ejection process. *Journal of Geophysical Research*, **105**, 2375–2392.

- LIN, J., RAYMOND, J.C. & VAN BALLEGOOIJEN, A.A. (2004). The Role of Magnetic Reconnection in the Observable Features of Solar Eruptions. *Astrophysical Journal*, **602**, 422–435.
- LINDEMANN, F.A. (1919). Note on the theory of magnetic storms. *Philosophical Magazine Series* 6, **38**, 669–684.
- Liu, Y., Davies, J.A., Luhmann, J.G., Vourlidas, A., Bale, S.D. & Lin, R.P. (2010). Geometric Triangulation of Imaging Observations to Track Coronal Mass Ejections Continuously Out to 1 AU. *Astrophysical Journal Letters*, **710**, L82–L87.
- Lynch, B.J., Antiochos, S.K., MacNeice, P.J., Zurbuchen, T.H. & Fisk, L.A. (2004). Observable Properties of the Breakout Model for Coronal Mass Ejections. *Astrophysical Journal*, **617**, 589–599.
- Lynch, B.J., Antiochos, S.K., Devore, C.R., Luhmann, J.G. & Zurbuchen, T.H. (2008). Topological Evolution of a Fast Magnetic Breakout CME in Three Dimensions. *Astrophysical Journal*, **683**, 1192–1206.
- MACNEICE, P., ANTIOCHOS, S.K., PHILLIPS, A., SPICER, D.S., DEVORE,
 C.R. & OLSON, K. (2004). A Numerical Study of the Breakout Model for
 Coronal Mass Ejection Initiation. Astrophysical Journal, 614, 1028–1041.
- MACQUEEN, R.M., EDDY, J.A., GOSLING, J.T., HILDNER, E., MUNRO, R.H., NEWKIRK, G.A., JR., POLAND, A.I. & ROSS, C.L. (1974). The Outer Solar Corona as Observed from Skylab: Preliminary Results. *Astrophysical Journal Letters*, **187**, L85+.

- MACQUEEN, R.M., CSOEKE-POECKH, A., HILDNER, E., HOUSE, L., REYNOLDS, R., STANGER, A., TEPOEL, H. & WAGNER, W. (1980). The High Altitude Observatory Coronagraph/Polarimeter on the Solar Maximum Mission. *Solar Physics*, **65**, 91–107.
- MACQUEEN, R.M., HUNDHAUSEN, A.J. & CONOVER, C.W. (1986). The propagation of coronal mass ejection transients. *Journal of Geophysical Research*, **91**, 31–38.
- MALONEY, S.A., GALLAGHER, P.T. & MCATEER, R.T.J. (2009). Reconstructing the 3-D Trajectories of CMEs in the Inner Heliosphere. *Solar Physics*, **256**, 149–166.
- Maričić, D., Vršnak, B., Stanger, A.L. & Veronig, A. (2004). Coronal Mass Ejection of 15 May 2001: I. Evolution of Morphological Features of the Eruption. *Solar Physics*, **225**, 337–353.
- McComas, D.J., Acton, L.W., Balat-Pichelin, M., Bothmer, V., Dirling, R.B., Jr., Feldman, W.C., Gloeckler, G., Habbal, S.R., Hassler, D.M., Mann, I., Matthaeus, W.H., McNutt, R.L., Jr., Mewaldt, R.A., Murphy, N., Ofman, L., Sittler, E.C., Jr., Smith, C.W., Velli, M. & Zurbuchen, T.H. (2008). Solar Probe Plus: Report of the Science and Technology Definition Team. NASA/TM-2008-214161, NASA.
- MIERLA, M., DAVILA, J., THOMPSON, W., INHESTER, B., SRIVASTAVA, N., KRAMAR, M., ST. CYR, O.C., STENBORG, G. & HOWARD, R.A. (2008).

- A Quick Method for Estimating the Propagation Direction of Coronal Mass Ejections Using STEREO-COR1 Images. *Solar Physics*, **252**, 385–396.
- MINNAERT, M. (1930). On the continuous spectrum of the corona and its polarisation. With 3 figures. (Received July 30, 1930). Zeitschrift fur Astrophysik, 1, 209-+.
- Moon, Y.J., Choe, G.S., Wang, H., Park, Y.D., Gopalswamy, N., Yang, G. & Yashiro, S. (2002). A Statistical Study of Two Classes of Coronal Mass Ejections. *Astrophysical Journal*, **581**, 694–702.
- Moore, R.L. & Sterling, A.C. (2006). Initiation of Coronal Mass Ejections.

 Washington DC American Geophysical Union Geophysical Monograph Series,

 165, 43-+.
- MORAN, T.G. & DAVILA, J.M. (2004). Three-Dimensional Polarimetric Imaging of Coronal Mass Ejections. *Science*, **305**, 66–71.
- Moran, T.G., Davila, J.M. & Thompson, W.T. (2010). Three-Dimensional Polarimetric Coronal Mass Ejection Localization Tested Through Triangulation. *Astrophysical Journal*, **712**, 453–458.
- MORGAN, H., HABBAL, S.R. & WOO, R. (2006). The Depiction of Coronal Structure in White-Light Images. *Solar Physics*, **236**, 263–272.
- Murtagh, F., Starck, J. & Bijaoui, A. (1995). Image restoration with noise suppression using a multiresolution support. *Astronomy & Astrophysics Supplemental*, **112**, 179–+.

- Odstrčil, D. & Pizzo, V.J. (1999). Three-dimensional propagation of coronal mass ejections in a structured solar wind flow 2. CME launched adjacent to the streamer belt. *Journal of Geophysical Research*, **104**, 493–504.
- OLMEDO, O., ZHANG, J., WECHSLER, H., POLAND, A. & BORNE, K. (2008).

 Automatic Detection and Tracking of Coronal Mass Ejections in Coronagraph

 Time Series. Solar Physics, 248, 485–499.
- OSTERBROCK, D.E. (1961). The Heating of the Solar Chromosphere, Plages, and Corona by Magnetohydrodynamic Waves. *Astrophysical Journal*, **134**, 347–+.
- Parker, E.N. (1957). Sweet's Mechanism for Merging Magnetic Fields in Conducting Fluids. *Journal of Geophysical Research*, **62**, 509–520.
- PARKER, E.N. (1958). Dynamics of the Interplanetary Gas and Magnetic Fields.

 Astrophysical Journal, 128, 664-+.
- Parks, G.K. (2004). Physics of space plasmas: an introduction.
- Petschek, H.E. (1964). Magnetic Field Annihilation. NASA Special Publication, **50**, 425-+.
- Phillips, A.C. (1999). The Physics of Stars, 2nd Edition.
- PIDDINGTON, J.H. (1958). Interplanetary Magnetic Field and Its Control of Cosmic-Ray Variations. *Physical Review*, **112**, 589–596.
- Pizzo, V.J. & Biesecker, D.A. (2004). Geometric localization of STEREO CMEs. Geophysical Research Letters, 31, 21802—+.

- PNEUMAN, G.W. & KOPP, R.A. (1971). Gas-Magnetic Field Interactions in the Solar Corona. *Solar Physics*, **18**, 258–270.
- Prialnik, D. (2000). An Introduction to the Theory of Stellar Structure and Evolution.
- Priest, E. & Forbes, T. (2000). Magnetic Reconnection.
- PRIEST, E.R. & FORBES, T.G. (1990a). Magnetic field evolution during prominence eruptions and two-ribbon flares. *Solar Physics*, **126**, 319–350.
- PRIEST, E.R. & FORBES, T.G. (1990b). The evolution of coronal magnetic fields. *Solar Physics*, **130**, 399–402.
- PRIEST, E.R. & FORBES, T.G. (2002). The magnetic nature of solar flares.

 Astronomy & Astrophysics Review, 10, 313–377.
- RILEY, P. & CROOKER, N.U. (2004). Kinematic Treatment of Coronal Mass Ejection Evolution in the Solar Wind. *Astrophysical Journal*, **600**, 1035–1042.
- ROBBRECHT, E. & BERGHMANS, D. (2004). Automated recognition of coronal mass ejections (CMEs) in near-real-time data. *Astronomy & Astrophysics*, **425**, 1097–1106.
- SCHATZMAN, E. (1949). The heating of the solar corona and chromosphere. Annales d'Astrophysique, 12, 203-+.
- Schrank, H. (1961). Graphical construction of rays in an ideal luneburg lens.

 Antennas and Propagation, IRE Transactions on, 9, 410 –411.
- SCHRIJVER, C.J. & ZWAAN, C. (2000). Solar and Stellar Magnetic Activity.

- Schrijver, C.J., Elmore, C., Kliem, B., Török, T. & Title, A.M. (2008). Observations and Modeling of the Early Acceleration Phase of Erupting Filaments Involved in Coronal Mass Ejections. *Astrophysical Journal*, **674**, 586–595.
- SCHWARZSCHILD, M. (1948). On Noise Arising from the Solar Granulation. Astrophysical Journal, 107, 1–+.
- Schwenn, R., dal Lago, A., Huttunen, E. & Gonzalez, W.D. (2005). The association of coronal mass ejections with their effects near the Earth.

 Annales Geophysicae, 23, 1033–1059.
- Sheeley, N.R., Walters, J.H., Wang, Y. & Howard, R.A. (1999). Continuous tracking of coronal outflows: Two kinds of coronal mass ejections.

 Journal of Geophysical Research, 104, 24739–24768.
- Sheeley, N.R., Jr., Michels, D.J., Howard, R.A. & Koomen, M.J. (1980). Initial observations with the SOLWIND coronagraph. *Astrophysical Journal Letters*, **237**, L99–L101.
- SRIVASTAVA, N., INHESTER, B., MIERLA, M. & PODLIPNIK, B. (2009). 3D Reconstruction of the Leading Edge of the 20 May 2007 Partial Halo CME. Solar Physics, 259, 213–225.
- Stenborg, G. & Cobelli, P.J. (2003). A wavelet packets equalization technique to reveal the multiple spatial-scale nature of coronal structures. *Astronomy & Astrophysics*, **398**, 1185–1193.

- Stenborg, G., Vourlidas, A. & Howard, R.A. (2008). A Fresh View of the Extreme-Ultraviolet Corona from the Application of a New Image-Processing Technique. *Astrophysical Journal*, **674**, 1201–1206.
- SWEET, P.A. (1958). The Neutral Point Theory of Solar Flares. In B. Lehnert, ed., *Electromagnetic Phenomena in Cosmical Physics*, vol. 6 of *IAU Symposium*, 123–+.
- Tappin, S.J. (2006). The Deceleration of an Interplanetary Transient from the Sun to 5 Au. *Solar Physics*, **233**, 233–248.
- Temmer, M., Veronig, A.M., Vršnak, B., Rybák, J., Gömöry, P., Stoiser, S. & Maričić, D. (2008). Acceleration in Fast Halo CMEs and Synchronized Flare HXR Bursts. *Astrophysical Journal Letters*, **673**, L95–L98.
- TEMMER, M., PREISS, S. & VERONIG, A.M. (2009). CME Projection Effects Studied with STEREO/COR and SOHO/LASCO. *Solar Physics*, **256**, 183–199.
- THERNISIEN, A., VOURLIDAS, A. & HOWARD, R.A. (2009). Forward Modeling of Coronal Mass Ejections Using STEREO/SECCHI Data. *Solar Physics*, **256**, 111–130.
- THERNISIEN, A.F.R., HOWARD, R.A. & VOURLIDAS, A. (2006). Modeling of Flux Rope Coronal Mass Ejections. *Astrophysical Journal*, **652**, 763–773.
- THOMPSON, W.T. (2006). Coordinate systems for solar image data. *Astronomy & Astrophysics*, **449**, 791–803.

- Thompson, W.T., Davila, J.M., Fisher, R.R., Orwig, L.E., Mentzell, J.E., Hetherington, S.E., Derro, R.J., Federline, R.E., Clark, D.C., Chen, P.T.C., Tveekrem, J.L., Martino, A.J., Novello, J., Wesenberg, R.P., Stcyr, O.C., Reginald, N.L., Howard, R.A., Mehalick, K.I., Hersh, M.J., Newman, M.D., Thomas, D.L., Card, G.L. & Elmore, D.F. (2003). COR1 inner coronagraph for Stereo-Secchi. In S. L. Keil & S. V. Avakyan, ed., Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol. 4853 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 1–11.
- Tousey, R. & Koomen, M. (1972). Movement of a Bright Source in the White-Light Corona. In *Bulletin of the American Astronomical Society*, vol. 4 of *Bulletin of the American Astronomical Society*, 394—+.
- VAN DE HULST, H.C. (1950). The electron density of the solar corona. Bulletin of the Astronomical Institutes of the Netherlands, 11, 135-+.
- VAN DER HOLST, B., JACOBS, C. & POEDTS, S. (2007). Simulation of a Breakout Coronal Mass Ejection in the Solar Wind. *Astrophysical Journal Letters*, **671**, L77–L80.
- Vourlidas, A. & Howard, R.A. (2006). The Proper Treatment of Coronal Mass Ejection Brightness: A New Methodology and Implications for Observations. *Astrophysical Journal*, **642**, 1216–1221.
- Vrsnak, B. (1990). Eruptive instability of cylindrical prominences. *Solar Physics*, **129**, 295–312.

- VRŠNAK, B. (2001). Deceleration of Coronal Mass Ejections. *Solar Physics*, **202**, 173–189.
- Vršnak, B. & Gopalswamy, N. (2002). Influence of the aerodynamic drag on the motion of interplanetary ejecta. *Journal of Geophysical Research (Space Physics)*, **107**, 1019–+.
- Vršnak, B., Ruždjak, D., Sudar, D. & Gopalswamy, N. (2004). Kinematics of coronal mass ejections between 2 and 30 solar radii. What can be learned about forces governing the eruption? *Astronomy & Astrophysics*, 423, 717–728.
- Vršnak, B., Sudar, D., Ruždjak, D. & Žic, T. (2007). Projection effects in coronal mass ejections. *Astronomy & Astrophysics*, **469**, 339–346.
- Vršnak, B., Vrbanec, D., Čalogović, J. & Žic, T. (2009). The role of aerodynamic drag in dynamics of coronal mass ejections. In N. Gopalswamy & D. F. Webb, ed., *IAU Symposium*, vol. 257 of *IAU Symposium*, 271–277.
- WEN, Y., MAIA, D.J.F. & WANG, J. (2007). The CME Acceleration Problem: Error Estimates in LASCO Coronal Mass Ejection Measurements. Astrophysical Journal, 657, 1117–1126.
- Wood, B.E., Howard, R.A., Thernisien, A., Plunkett, S.P. & Socker, D.G. (2009). Reconstructing the 3D Morphology of the 17 May 2008 CME. Solar Physics, 259, 163–178.
- Wuelser, J., Lemen, J.R., Tarbell, T.D., Wolfson, C.J., Cannon, J.C., Carpenter, B.A., Duncan, D.W., Gradwohl, G.S.,

MEYER, S.B., MOORE, A.S., NAVARRO, R.L., PEARSON, J.D., ROSSI, G.R., SPRINGER, L.A., HOWARD, R.A., MOSES, J.D., NEWMARK, J.S., DELABOUDINIERE, J., ARTZNER, G.E., AUCHERE, F., BOUGNET, M., BOUYRIES, P., BRIDOU, F., CLOTAIRE, J., COLAS, G., DELMOTTE, F., JEROME, A., LAMARE, M., MERCIER, R., MULLOT, M., RAVET, M., SONG, X., BOTHMER, V. & DEUTSCH, W. (2004). EUVI: the STEREO-SECCHI extreme ultraviolet imager. In S. Fineschi & M. A. Gummin, ed., Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol. 5171 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 111–122.

- XIE, H., OFMAN, L. & LAWRENCE, G. (2004). Cone model for halo CMEs: Application to space weather forecasting. *Journal of Geophysical Research (Space Physics)*, **109**, 3109–+.
- XIE, H., St. Cyr, O.C., Gopalswamy, N., Yashiro, S., Krall, J., Kramar, M. & Davila, J. (2009). On the Origin, 3D Structure and Dynamic Evolution of CMEs Near Solar Minimum. *Solar Physics*, **259**, 143–161.
- Xue, X.H., Wang, C.B. & Dou, X.K. (2005). An ice-cream cone model for coronal mass ejections. *Journal of Geophysical Research (Space Physics)*, **110**, 8103–+.
- Yashiro, S., Gopalswamy, N., Michalek, G., St. Cyr, O.C., Plunkett, S.P., Rich, N.B. & Howard, R.A. (2004). A catalog of white light coronal mass ejections observed by the SOHO spacecraft. *Journal of Geophysical Research (Space Physics)*, **109**, 7105–+.

- Young, C.A. & Gallagher, P.T. (2008). Multiscale Edge Detection in the Corona. Solar Physics, 248, 457–469.
- ZHANG, J. & DERE, K.P. (2006). A Statistical Study of Main and Residual Accelerations of Coronal Mass Ejections. Astrophysical Journal, 649, 1100– 1109.
- ZHANG, J. & WANG, J. (2002). Are Homologous Flare-Coronal Mass Ejection Events Triggered by Moving Magnetic Features? Astrophysical Journal, 566, L117–L120.
- Zhao, X.P., Plunkett, S.P. & Liu, W. (2002). Determination of geometrical and kinematical properties of halo coronal mass ejections using the cone model.

 Journal of Geophysical Research (Space Physics), 107, 1223—+.
- ZIRIN, H. (1998). The Astrophysics of the Sun.